



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:27 PM UTC

PDB ID : 6IAM / pdb_00006iam
Title : Modulating Protein-Protein Interactions with Visible Light Peptide Backbone Switches
Authors : Werel, L.; Essen, L.-O.
Deposited on : 2018-11-27
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	16% 95% 5%
2	B	12	42% 75% 17% 8%
3	C	4	50% 50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5458 atoms, of which 2509 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WD repeat-containing protein 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	306	Total	C	H	N	O	S	0	4	0
			4765	1524	2374	400	456	11			

- Molecule 2 is a protein called SER-ALA-ARG-ALA-XY5-VAL-HIS-LEU-ARG-LYS-SER-ALA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	H	N	O	0	0	1
			202	63	102	24	13			

- Molecule 3 is a protein called Small ubiquitin-related modifier 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	4	Total	C	H	N	O	0	0	0
			63	19	33	5	6			

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	K	0	0
			3	3		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	395	Total 395	O 395	0	0
6	B	19	Total 19	O 19	0	0
6	C	9	Total 9	O 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: WD repeat-containing protein 5



- Molecule 2: SER-ALA-ARG-ALA-XY5-VAL-HIS-LEU-ARG-LYS-SER-ALA



- Molecule 3: Small ubiquitin-related modifier 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.53Å 46.56Å 66.17Å 90.00° 107.22° 90.00°	Depositor
Resolution (Å)	37.49 – 1.51 37.49 – 1.51	Depositor EDS
% Data completeness (in resolution range)	99.3 (37.49-1.51) 99.3 (37.49-1.51)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 1.51Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.148 , 0.170 0.208 , 0.212	Depositor DCC
R_{free} test set	2053 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	11.3	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5458	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XY5, NA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2460	0.64	0/3336
2	B	1.21	2/78 (2.6%)	1.97	2/101 (2.0%)
3	C	0.32	0/30	0.75	0/39
All	All	0.36	2/2568 (0.1%)	0.72	2/3476 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	VAL	CA-C	-8.41	1.35	1.52
2	B	6	VAL	CA-CB	5.59	1.69	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6	VAL	N-CA-C	14.63	151.97	111.00
2	B	6	VAL	CB-CA-C	-9.07	94.17	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	4	ALA	Peptide
2	B	5	XY5	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2391	2374	2373	12	0
2	B	100	102	88	1	0
3	C	30	33	33	1	0
4	A	3	0	0	0	0
5	A	2	0	0	0	0
6	A	395	0	0	8	2
6	B	19	0	0	0	0
6	C	9	0	0	0	0
All	All	2949	2509	2494	14	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:GLU:OE2	6:A:501:HOH:O	1.76	1.02
1:A:204:GLN:OE1	6:A:502:HOH:O	1.78	0.99
1:A:195[B]:CYS:SG	1:A:217:VAL:HG11	2.25	0.77
1:A:211:ASP:OD2	6:A:504:HOH:O	2.13	0.66
1:A:229:ILE:HD12	1:A:243:TYR:HB3	1.84	0.58
1:A:263:PHE:O	6:A:505:HOH:O	2.19	0.53
1:A:29:THR:O	1:A:31:VAL:HG13	2.13	0.48
1:A:204:GLN:HB3	6:A:503:HOH:O	2.13	0.48
1:A:204:GLN:NE2	6:A:503:HOH:O	2.05	0.48
1:A:29:THR:O	1:A:31:VAL:N	2.48	0.47
6:A:547:HOH:O	2:B:6:VAL:HG12	2.15	0.46
1:A:41:LEU:HB2	1:A:327:ILE:HB	1.99	0.44
3:C:3:LYS:O	3:C:4:PRO:C	2.61	0.44
1:A:57:GLY:HA2	6:A:630:HOH:O	2.21	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:659:HOH:O	6:A:803:HOH:O[2_444]	1.96	0.24
6:A:713:HOH:O	6:A:853:HOH:O[2_444]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/306 (101%)	296 (96%)	12 (4%)	0	100	100
2	B	9/12 (75%)	9 (100%)	0	0	100	100
3	C	2/4 (50%)	2 (100%)	0	0	100	100
All	All	319/322 (99%)	307 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/268 (102%)	271 (100%)	1 (0%)	84	71
2	B	8/8 (100%)	8 (100%)	0	100	100
3	C	3/3 (100%)	3 (100%)	0	100	100
All	All	283/279 (101%)	282 (100%)	1 (0%)	84	71

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/306 (100%)	0.81	50 (16%) 4 5	7, 13, 32, 102	3 (0%)
2	B	11/12 (91%)	1.72	5 (45%) 0 0	9, 30, 46, 51	0
3	C	4/4 (100%)	2.11	2 (50%) 0 0	25, 36, 48, 49	0
All	All	321/322 (99%)	0.86	57 (17%) 4 4	7, 13, 42, 102	3 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	LEU	10.7
1	A	191	TYR	8.9
1	A	190	SER	8.8
1	A	193	GLY	8.7
1	A	210	ILE	8.6
1	A	121	CYS	7.7
1	A	29	THR	7.1
1	A	95	TRP	7.1
1	A	187	VAL	7.0
1	A	74	ALA	6.6
1	A	192	ASP	6.1
1	A	188	SER	5.9
1	A	31	VAL	5.8
1	A	140[A]	GLN	5.2
1	A	138	ASN	5.2
2	B	12	ALA	5.2
1	A	189	SER	5.0
1	A	139	PRO	4.9
1	A	30	PRO	4.5
2	B	6	VAL	4.2
1	A	211	ASP	4.1
1	A	94	ALA	4.0
1	A	123	LYS	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	243	TYR	3.8
1	A	212	ASP	3.7
1	A	145	VAL	3.7
3	C	4	PRO	3.5
1	A	96	SER	3.5
1	A	53	PHE	3.4
1	A	159	LYS	3.4
3	C	2	ALA	3.2
1	A	209	LEU	3.2
1	A	186	ILE	3.0
1	A	32	LYS	2.9
1	A	178	HIS	2.9
2	B	11	SER	2.9
1	A	78	LYS	2.8
1	A	72	TRP	2.8
1	A	173	PRO	2.8
1	A	244	SER	2.7
1	A	107	ASP	2.6
1	A	137	PHE	2.6
1	A	247	LYS	2.5
1	A	213	ASP	2.4
1	A	129	SER	2.4
1	A	225	ASN	2.4
1	A	114	TRP	2.3
2	B	8	LEU	2.3
1	A	116	VAL	2.3
1	A	42	ALA	2.2
1	A	54	SER	2.1
1	A	245	LYS	2.1
1	A	47	ALA	2.0
1	A	128	HIS	2.0
1	A	323	ASN	2.0
2	B	9	ARG	2.0
1	A	179	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K	A	403	1/1	0.75	0.20	61,61,61,61	0
5	NA	A	405	1/1	0.75	0.31	36,36,36,36	0
4	K	A	402	1/1	0.83	0.20	45,45,45,45	0
4	K	A	401	1/1	0.92	0.28	38,38,38,38	0
5	NA	A	404	1/1	0.95	0.09	30,30,30,30	1

6.5 Other polymers [i](#)

There are no such residues in this entry.