



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:17 AM UTC

PDB ID : 6ICW / pdb_00006icw
Title : Crystal structure of H7 hemagglutinin mutant AH-SGTQ (A138S, V186G, P221T and L226Q) from the influenza virus A/Anhui/1/2013 (H7N9)
Authors : Gao, G.F.; Xu, Y.
Deposited on : 2018-09-07
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	
2	B	177	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2386	1478	433	460	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	128	SER	ALA	engineered mutation	UNP R4NN21
A	177	GLY	VAL	engineered mutation	UNP R4NN21
A	212	THR	PRO	engineered mutation	UNP R4NN21
A	217	GLN	LEU	engineered mutation	UNP R4NN21

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	163	Total	C	N	O	S	0	0	0
			1328	817	231	273	7			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total	O	0	0
			54	54		
4	B	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	115.61Å 115.61Å 293.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.53 – 2.60 37.53 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.53-2.60) 99.8 (37.53-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.258 , 0.283 0.258 , 0.280	Depositor DCC
R_{free} test set	1219 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.524	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 61.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.005 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.021 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.011 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3825	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/2431	0.34	0/3286
2	B	0.21	0/1351	1.07	8/1821 (0.4%)
All	All	0.16	0/3782	0.69	8/5107 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	462	PHE	N-CA-C	24.14	141.99	112.54
2	B	462	PHE	CA-C-N	19.77	151.35	122.09
2	B	462	PHE	C-N-CA	19.77	151.35	122.09
2	B	463	HIS	N-CA-C	15.48	135.97	109.96
2	B	463	HIS	N-CA-CB	-9.72	94.48	110.23
2	B	448	ARG	N-CA-C	9.40	123.36	111.24
2	B	461	ILE	N-CA-CB	-7.15	103.10	111.39
2	B	448	ARG	CB-CA-C	-6.39	100.31	110.86

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	202	GLN	Peptide
2	B	462	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2386	0	2328	33	0
2	B	1328	0	1222	11	1
3	A	28	0	26	0	0
3	B	14	0	13	0	0
4	A	54	0	0	1	0
4	B	15	0	0	1	0
All	All	3825	0	3589	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:THR:HA	1:A:236:PHE:O	1.92	0.69
1:A:169:ILE:O	1:A:225:TRP:HA	1.92	0.69
1:A:168:LEU:HB3	1:A:249:SER:HB2	1.81	0.63
2:B:461:ILE:HG23	2:B:461:ILE:O	2.02	0.60
2:B:444:LYS:NZ	4:B:601:HOH:O	2.31	0.59
1:A:309:THR:HG22	2:B:373:LEU:HD11	1.84	0.58
1:A:170:VAL:HG22	1:A:225:TRP:HB3	1.87	0.57
1:A:49:VAL:HG23	1:A:74:ALA:HB2	1.87	0.56
1:A:71:GLU:OE2	1:A:247:ARG:NH2	2.39	0.55
1:A:203:SER:O	1:A:203:SER:OG	2.21	0.54
1:A:194:THR:O	1:A:236:PHE:HA	2.07	0.54
1:A:106:GLY:HA2	1:A:255:SER:HB3	1.91	0.53
1:A:116:THR:O	1:A:157:LYS:NZ	2.41	0.52
2:B:466:ASP:OD1	2:B:467:ASP:N	2.43	0.52
1:A:7:HIS:HB2	1:A:311:MET:HE1	1.92	0.52
1:A:170:VAL:HA	1:A:224:HIS:O	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:463:HIS:HB3	2:B:487:ALA:HB2	1.93	0.51
1:A:202:GLN:HA	1:A:203:SER:CB	2.41	0.51
2:B:441:GLU:HB3	2:B:445:ARG:NH1	2.26	0.51
1:A:87:CYS:SG	1:A:88:TYR:N	2.83	0.51
1:A:148:ASP:O	1:A:149:ASN:C	2.54	0.51
1:A:143:LEU:HD12	1:A:242:PHE:HD2	1.77	0.50
1:A:44:LYS:HE3	1:A:269:GLU:HB2	1.94	0.49
1:A:171:TRP:CE2	1:A:195:VAL:HG21	2.48	0.48
1:A:120:ILE:HD11	1:A:153:PRO:HD2	1.96	0.47
2:B:349:ASN:ND2	2:B:467:ASP:OD1	2.47	0.46
1:A:135:SER:OG	1:A:136:PHE:N	2.48	0.46
1:A:95:GLU:HB3	1:A:99:ARG:NH1	2.31	0.45
1:A:192:LEU:N	1:A:239:ASN:OD1	2.42	0.45
1:A:40:ARG:HD2	1:A:265:ASP:HB2	1.99	0.44
2:B:442:ARG:NH1	2:B:446:GLN:OE1	2.51	0.44
1:A:31:GLU:OE1	1:A:304:SER:OG	2.36	0.43
1:A:174:HIS:ND1	1:A:186:TYR:OH	2.38	0.43
1:A:110:LYS:NZ	1:A:139:GLU:OE2	2.49	0.42
1:A:139:GLU:OE1	1:A:247:ARG:HD3	2.20	0.42
2:B:342:TRP:HB2	2:B:362:THR:HG22	2.01	0.42
2:B:366:ILE:O	2:B:370:THR:OG1	2.32	0.42
1:A:3:ILE:N	4:A:714:HOH:O	2.52	0.41
1:A:183:THR:HG22	1:A:188:SER:HA	2.02	0.41
1:A:89:PRO:HG3	1:A:217:GLN:HB3	2.04	0.40
1:A:200:TYR:CE2	1:A:202:GLN:HB2	2.56	0.40
1:A:105:SER:HB2	1:A:251:LEU:HD22	2.03	0.40
2:B:465:CYS:SG	2:B:470:MET:HG3	2.61	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:448:ARG:NH1	2:B:460:GLU:OE1[2_545]	1.94	0.26

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/321 (97%)	286 (92%)	25 (8%)	1 (0%)	36	58
2	B	161/177 (91%)	155 (96%)	6 (4%)	0	100	100
All	All	473/498 (95%)	441 (93%)	31 (7%)	1 (0%)	43	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/269 (97%)	256 (98%)	5 (2%)	50	75
2	B	141/152 (93%)	136 (96%)	5 (4%)	32	59
All	All	402/421 (96%)	392 (98%)	10 (2%)	42	69

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	CYS
1	A	145	SER
1	A	147	THR
1	A	176	SER
1	A	205	VAL
2	B	449	GLU
2	B	456	THR
2	B	460	GLU
2	B	463	HIS
2	B	469	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	A	217	GLN
2	B	480	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	601	1	14,14,15	0.26	0	17,19,21	0.43	0
3	NAG	B	501	2	14,14,15	0.28	0	17,19,21	0.47	0
3	NAG	A	602	1	14,14,15	0.20	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	3/6/23/26	0/1/1/1
3	NAG	B	501	2	-	0/6/23/26	0/1/1/1
3	NAG	A	602	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	NAG	O5-C5-C6-O6
3	A	601	NAG	O5-C5-C6-O6
3	A	602	NAG	C4-C5-C6-O6
3	A	601	NAG	C4-C5-C6-O6
3	A	601	NAG	C1-C2-N2-C7
3	A	602	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/321 (97%)	1.08	72 (22%) 2 2	29, 61, 109, 142	0
2	B	163/177 (92%)	1.41	45 (27%) 1 1	30, 85, 131, 157	0
All	All	477/498 (95%)	1.19	117 (24%) 2 1	29, 67, 126, 157	0

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	203	SER	7.7
2	B	328	ALA	6.3
2	B	452	GLU	5.7
1	A	149	ASN	5.4
1	A	222	ASP	5.3
1	A	207	SER	5.2
2	B	438	LYS	4.9
2	B	464	LYS	4.9
1	A	316	GLU	4.8
1	A	151	ALA	4.6
1	A	156	THR	4.6
1	A	183	THR	4.6
1	A	145	SER	4.3
1	A	195	VAL	4.3
2	B	459	PHE	4.2
1	A	205	VAL	4.1
1	A	180	ALA	4.1
1	A	190	ASN	4.1
2	B	460	GLU	4.0
2	B	451	ALA	4.0
1	A	237	SER	4.0
1	A	189	GLY	3.9
2	B	461	ILE	3.9
2	B	487	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	235	THR	3.8
2	B	350	ALA	3.8
1	A	120	ILE	3.7
1	A	194	THR	3.7
2	B	354	GLY	3.7
1	A	216	GLY	3.7
2	B	355	THR	3.6
1	A	236	PHE	3.6
2	B	388	ASP	3.6
1	A	199	ASN	3.6
1	A	186	TYR	3.5
1	A	215	ASN	3.5
1	A	193	VAL	3.3
1	A	5	LEU	3.3
1	A	210	ALA	3.3
2	B	329	GLY	3.3
1	A	191	LYS	3.2
1	A	242	PHE	3.2
1	A	196	GLY	3.2
1	A	311	MET	3.2
2	B	344	GLY	3.2
2	B	445	ARG	3.1
2	B	462	PHE	3.1
1	A	185	LEU	3.0
1	A	181	GLU	3.0
1	A	241	ALA	3.0
2	B	447	LEU	3.0
2	B	488	MET	3.0
1	A	147	THR	3.0
2	B	366	ILE	2.9
2	B	482	LYS	2.9
1	A	3	ILE	2.9
1	A	192	LEU	2.9
2	B	377	ILE	2.8
1	A	4	CYS	2.8
1	A	204	PHE	2.8
1	A	21	GLU	2.8
2	B	478	TYR	2.8
2	B	379	LYS	2.8
1	A	162	THR	2.8
1	A	154	GLN	2.7
1	A	187	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	198	SER	2.7
2	B	483	TYR	2.6
2	B	456	THR	2.6
2	B	375	ARG	2.6
2	B	443	VAL	2.6
1	A	240	GLY	2.6
2	B	359	TYR	2.6
2	B	485	GLU	2.5
2	B	380	THR	2.5
1	A	118	SER	2.5
1	A	178	SER	2.5
1	A	116	THR	2.5
1	A	122	THR	2.5
1	A	201	GLN	2.5
1	A	22	ARG	2.4
1	A	206	PRO	2.4
1	A	212	THR	2.4
1	A	179	THR	2.4
2	B	489	GLN	2.4
2	B	343	TYR	2.4
1	A	188	SER	2.4
1	A	155	MET	2.3
1	A	184	LYS	2.3
1	A	197	SER	2.3
2	B	455	GLY	2.3
2	B	347	HIS	2.3
2	B	433	ASP	2.3
1	A	119	GLY	2.3
1	A	213	GLN	2.3
2	B	454	ASP	2.3
1	A	200	TYR	2.3
1	A	182	GLN	2.3
2	B	449	GLU	2.2
2	B	490	ASN	2.2
2	B	337	GLY	2.2
1	A	152	PHE	2.2
1	A	208	PRO	2.2
2	B	338	LEU	2.2
1	A	117	TYR	2.2
2	B	348	GLN	2.2
1	A	239	ASN	2.2
1	A	172	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	238	PHE	2.2
1	A	129	CYS	2.1
2	B	360	LYS	2.1
1	A	12	ASN	2.1
2	B	352	GLY	2.1
1	A	150	ALA	2.1
2	B	470	MET	2.1
1	A	173	ILE	2.0
1	A	121	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	602	14/15	0.49	0.19	93,108,113,115	0
3	NAG	A	601	14/15	0.57	0.20	95,112,129,130	0
3	NAG	B	501	14/15	0.80	0.15	47,72,78,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.