



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:04 AM UTC

PDB ID : 3IDM / pdb_00003idm
Title : Crystal structure of the HIV-1 Cross Neutralizing Monoclonal Antibody 2F5 Fab' fragment in complex with gp41 Peptide analog ELD(Nrg)WAS
Authors : Julien, J.-P.; Bryson, S.; Pai, E.F.
Deposited on : 2009-07-21
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	11% 84% 15% .
2	B	237	9% 81% 14% ..
3	C	7	14% 86% 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3634 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2F5 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1644	1025	281	333	5			

- Molecule 2 is a protein called 2F5 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	227	Total	C	N	O	S	0	0	0
			1703	1082	289	325	7			

- Molecule 3 is a protein called gp41 MPER peptide analog.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	0	0	0
			65	38	12	15			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	99	Total	O	0	0
			99	99		
4	B	120	Total	O	0	0
			120	120		
4	C	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2F5 Fab light chain



- Molecule 2: 2F5 Fab heavy chain



- Molecule 3: gp41 MPER peptide analog



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.60Å 64.90Å 176.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.89 – 2.24 60.89 – 2.24	Depositor EDS
% Data completeness (in resolution range)	93.1 (60.89-2.24) 93.1 (60.89-2.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.66	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.22Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.229 , 0.236 0.227 , 0.236	Depositor DCC
R_{free} test set	1591 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	38.5	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3634	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NRG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	1/1681 (0.1%)	0.97	6/2284 (0.3%)
2	B	0.71	0/1744	1.03	6/2384 (0.3%)
3	C	0.71	0/51	1.11	0/66
All	All	0.69	1/3476 (0.0%)	1.00	12/4734 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	VAL	CA-CB	5.29	1.61	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	ASP	N-CA-C	9.27	123.09	111.69
2	B	144	ASP	N-CA-C	6.50	118.51	110.91
1	A	114	SER	N-CA-C	-5.79	99.09	108.41
1	A	97	THR	N-CA-C	5.74	118.55	110.23
1	A	133	VAL	N-CA-C	5.59	116.63	108.53
1	A	168	SER	N-CA-C	5.53	119.13	112.38
1	A	138	ASN	N-CA-C	5.51	117.91	111.02
2	B	67	LEU	N-CA-C	5.46	118.72	109.76
2	B	33	GLY	N-CA-C	5.35	122.65	115.59
1	A	208	SER	N-CA-C	5.28	117.00	109.14
2	B	24	PHE	N-CA-C	5.22	116.92	109.14
2	B	143	LYS	N-CA-C	5.11	118.40	110.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1644	0	1588	18	0
2	B	1703	0	1711	15	0
3	C	65	0	52	3	0
4	A	99	0	0	0	0
4	B	120	0	0	0	0
4	C	3	0	0	0	0
All	All	3634	0	3351	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:GLN:NE2	1:A:93:PHE:H	1.63	0.94
2:B:87:THR:HG22	2:B:111:ILE:H	1.29	0.94
1:A:90:GLN:HE22	1:A:93:PHE:H	0.97	0.90
1:A:106:VAL:H	1:A:166:GLN:HE22	1.35	0.74
1:A:90:GLN:HE22	1:A:93:PHE:N	1.81	0.72
1:A:90:GLN:HE21	1:A:92:HIS:H	1.45	0.62
2:B:100(H):ARG:O	2:B:100(K):VAL:HG22	2.01	0.60
1:A:90:GLN:NE2	1:A:92:HIS:H	2.00	0.59
1:A:6:GLN:HE21	1:A:99:GLY:HA3	1.68	0.58
2:B:12:VAL:HG21	2:B:18:LEU:HD13	1.87	0.55
1:A:90:GLN:NE2	1:A:93:PHE:N	2.46	0.54
2:B:54:ASP:OD1	3:C:4:NRG:NH2	2.41	0.53
2:B:38:ARG:HD2	2:B:48:LEU:HD21	1.91	0.53
1:A:155:GLN:HB3	1:A:158:ASN:HD21	1.77	0.49
2:B:84:PRO:O	2:B:87:THR:HG23	2.14	0.48
2:B:82(C):VAL:HB	2:B:111:ILE:HD13	1.96	0.48
2:B:150:VAL:HG11	2:B:178:LEU:HD11	1.95	0.48
1:A:124:GLN:NE2	1:A:131:SER:H	2.11	0.48
2:B:112:SER:OG	2:B:114:THR:HG22	2.14	0.47
1:A:124:GLN:HE22	1:A:131:SER:H	1.64	0.45
2:B:54:ASP:CG	3:C:4:NRG:NH2	2.74	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:LYS:H	2:B:16:GLN:NE2	2.15	0.45
1:A:21:ILE:HG23	1:A:102:THR:HG21	1.98	0.45
1:A:20:THR:HG22	1:A:74:THR:HB	1.98	0.44
1:A:61:ARG:HB2	1:A:76:SER:O	2.17	0.44
1:A:105:ASP:HB2	1:A:166:GLN:NE2	2.33	0.44
2:B:143:LYS:NZ	2:B:171:GLN:OE1	2.51	0.44
1:A:90:GLN:NE2	1:A:92:HIS:N	2.65	0.44
2:B:12:VAL:O	2:B:111:ILE:HA	2.17	0.43
3:C:4:NRG:NH2	3:C:4:NRG:O3	2.50	0.43
2:B:193:THR:HB	2:B:210:ARG:HE	1.84	0.42
2:B:142:VAL:HB	2:B:178:LEU:HD13	2.02	0.42
1:A:8:PRO:O	1:A:102:THR:HG22	2.19	0.42
1:A:38:GLN:O	1:A:84:ALA:HB1	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/214 (99%)	206 (97%)	6 (3%)	0	100	100
2	B	221/237 (93%)	212 (96%)	8 (4%)	1 (0%)	24	23
3	C	4/7 (57%)	4 (100%)	0	0	100	100
All	All	437/458 (95%)	422 (97%)	14 (3%)	1 (0%)	43	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	126	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/187 (100%)	176 (94%)	11 (6%)	18	16
2	B	197/207 (95%)	187 (95%)	10 (5%)	21	21
3	C	5/5 (100%)	5 (100%)	0	100	100
All	All	389/399 (98%)	368 (95%)	21 (5%)	20	19

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ARG
1	A	22	THR
1	A	33	LEU
1	A	54	LEU
1	A	74	THR
1	A	133	VAL
1	A	143	GLU
1	A	154	LEU
1	A	158	ASN
1	A	169	LYS
1	A	188	LYS
2	B	29	LEU
2	B	38	ARG
2	B	60	SER
2	B	107	ILE
2	B	138	LEU
2	B	159	LEU
2	B	170	LEU
2	B	178	LEU
2	B	195	THR
2	B	214	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	38	GLN
1	A	90	GLN
1	A	124	GLN
1	A	137	ASN
1	A	155	GLN
1	A	158	ASN
1	A	166	GLN
1	A	210	ASN
2	B	16	GLN
2	B	39	GLN
2	B	164	HIS
2	B	199	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NRG	C	4	3	7,13,14	1.56	1 (14%)	5,15,17	2.90	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NRG	C	4	3	-	2/9/13/15	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	4	NRG	CB-CA	-2.71	1.49	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	4	NRG	NE-CZ-NH2	-3.94	113.00	120.26
3	C	4	NRG	NH1-CZ-NE	3.47	126.92	117.58
3	C	4	NRG	CD-NE-CZ	3.41	129.71	123.46

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	4	NRG	NE-CD-CG-CB
3	C	4	NRG	CG-CD-NE-CZ

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	4	NRG	3	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	214/214 (100%)	0.65	23 (10%)	11 10	23, 36, 54, 61	0
2	B	227/237 (95%)	0.62	21 (9%)	14 13	21, 35, 58, 82	0
3	C	6/7 (85%)	0.74	1 (16%)	4 3	28, 40, 43, 59	0
All	All	447/458 (97%)	0.63	45 (10%)	12 11	21, 35, 57, 82	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	127	SER	11.0
2	B	216	CYS	10.1
2	B	126	PRO	6.2
2	B	215	SER	5.8
2	B	134	GLY	5.7
2	B	185	PRO	4.5
2	B	214	LYS	4.4
1	A	193	GLU	4.3
1	A	203	SER	4.0
1	A	212	GLY	3.8
1	A	170	ASP	3.7
2	B	114	THR	3.6
1	A	1	ALA	3.5
1	A	126	LYS	3.4
2	B	157	GLY	3.4
1	A	154	LEU	3.3
2	B	135	THR	3.1
2	B	191	THR	3.1
1	A	169	LYS	3.1
1	A	76	SER	3.0
2	B	213	PRO	3.0
2	B	192	GLN	2.9
1	A	152	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	53	SER	2.9
2	B	204	ASN	2.9
1	A	7	SER	2.8
1	A	77	THR	2.8
1	A	109	THR	2.7
2	B	188	SER	2.6
3	C	7	SER	2.6
2	B	158	ALA	2.6
2	B	187	SER	2.6
1	A	2	LEU	2.5
1	A	18	ARG	2.5
2	B	100(B)	PHE	2.3
1	A	70	GLU	2.3
1	A	22	THR	2.2
1	A	190	LYS	2.2
2	B	201	LYS	2.2
1	A	123	GLU	2.2
1	A	188	LYS	2.2
1	A	67	SER	2.1
1	A	12	SER	2.1
1	A	9	SER	2.0
2	B	161	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NRG	C	4	14/15	0.87	0.14	31,39,54,56	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.