



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:37 PM UTC

PDB ID : 6ID5 / pdb_00006id5
Title : Crystal structure of H7 hemagglutinin mutant H7-SVPL (A138S) from the influenza virus A/Anhui/1/2013 (H7N9)
Authors : Gao, G.F.; Xu, Y.; Qi, J.X.
Deposited on : 2018-09-08
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	
2	B	177	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2388	1483	432	458	15	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	128	SER	ALA	engineered mutation	UNP R4NN21

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	163	1328	817	231	273	7	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	14	Total	O	0	0
			14	14		
4	B	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	115.39Å 115.39Å 294.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.59 – 2.60 36.59 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.59-2.60) 99.5 (36.59-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.259 , 0.310 0.259 , 0.311	Depositor DCC
R_{free} test set	1155 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	64.8	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.016 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.008 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3786	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2434	0.37	1/3292 (0.0%)
2	B	0.35	0/1351	0.60	4/1821 (0.2%)
All	All	0.24	0/3785	0.47	5/5113 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	331	ILE	N-CA-C	-17.84	80.94	108.23
2	B	332	GLU	N-CA-C	7.96	122.42	112.87
2	B	331	ILE	CB-CA-C	7.43	119.95	110.96
2	B	333	ASN	CB-CA-C	5.43	121.23	110.42
1	A	202	GLN	CB-CA-C	5.32	125.80	113.19

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	202	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2337	56	1
2	B	1328	0	1222	33	0
3	A	28	0	26	0	0
3	B	14	0	13	0	0
4	A	14	0	0	10	0
4	B	14	0	0	12	0
All	All	3786	0	3598	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:423:MET:SD	4:B:611:HOH:O	2.00	1.15
2:B:423:MET:CB	4:B:611:HOH:O	2.02	1.08
1:A:191:LYS:O	1:A:205:VAL:HG13	1.54	1.06
1:A:202:GLN:HA	1:A:203:SER:HB3	1.37	1.05
1:A:169:ILE:HA	4:A:701:HOH:O	1.68	0.92
1:A:204:PHE:CE2	1:A:224:HIS:CE1	2.59	0.89
1:A:204:PHE:HE2	1:A:224:HIS:CE1	1.92	0.88
1:A:121:ARG:NH1	1:A:145:SER:O	2.06	0.87
2:B:329:GLY:N	4:B:602:HOH:O	1.98	0.87
2:B:418:GLU:OE1	4:B:601:HOH:O	1.93	0.86
2:B:423:MET:CG	4:B:611:HOH:O	2.11	0.83
1:A:205:VAL:HG12	1:A:206:PRO:HD2	1.64	0.80
1:A:202:GLN:HA	1:A:203:SER:CB	2.12	0.78
1:A:170:VAL:N	4:A:701:HOH:O	1.94	0.77
1:A:67:ASP:N	4:A:702:HOH:O	2.17	0.76
2:B:388:ASP:OD2	2:B:406:ARG:NH2	2.17	0.76
1:A:205:VAL:CG1	1:A:206:PRO:HD2	2.15	0.74
2:B:332:GLU:O	2:B:333:ASN:HB3	1.88	0.73
2:B:328:ALA:N	4:B:602:HOH:O	2.21	0.72
1:A:94:ASN:ND2	4:A:703:HOH:O	2.20	0.72
1:A:191:LYS:O	1:A:205:VAL:CG1	2.37	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:411:GLU:HB3	4:B:604:HOH:O	1.97	0.65
1:A:20:THR:HG22	1:A:21:GLU:HG3	1.79	0.65
2:B:375:ARG:NH2	2:B:424:GLU:OE2	2.30	0.65
2:B:346:ARG:HA	2:B:354:GLY:O	1.97	0.64
2:B:423:MET:HB2	4:B:611:HOH:O	1.79	0.63
1:A:202:GLN:CA	1:A:203:SER:HB3	2.20	0.63
2:B:380:THR:HG22	2:B:382:GLN:H	1.64	0.63
2:B:383:GLN:NE2	2:B:385:GLU:OE2	2.26	0.61
1:A:202:GLN:HB3	1:A:203:SER:OG	2.01	0.60
1:A:66:CYS:C	4:A:702:HOH:O	2.44	0.59
1:A:110:LYS:NZ	1:A:139:GLU:OE2	2.35	0.58
1:A:170:VAL:HG22	1:A:225:TRP:HB3	1.85	0.58
2:B:411:GLU:CG	4:B:604:HOH:O	2.52	0.57
2:B:450:ASN:ND2	2:B:478:TYR:OH	2.37	0.56
2:B:347:HIS:CD2	2:B:474:ARG:HH12	2.24	0.55
1:A:116:THR:HA	4:A:706:HOH:O	2.07	0.55
2:B:347:HIS:HD2	2:B:474:ARG:HH12	1.54	0.54
1:A:169:ILE:O	1:A:225:TRP:HA	2.08	0.53
1:A:121:ARG:NH1	1:A:146:ASN:O	2.43	0.52
1:A:202:GLN:CA	1:A:203:SER:CB	2.83	0.52
2:B:395:GLU:OE2	2:B:397:GLN:HB3	2.10	0.52
1:A:309:THR:HG22	2:B:373:LEU:HD11	1.92	0.51
2:B:328:ALA:CA	4:B:602:HOH:O	2.57	0.51
2:B:411:GLU:CB	4:B:604:HOH:O	2.56	0.51
1:A:131:ARG:NH1	1:A:136:PHE:O	2.39	0.51
1:A:44:LYS:HD3	1:A:269:GLU:HB2	1.92	0.50
1:A:168:LEU:HB3	1:A:249:SER:HB2	1.94	0.49
1:A:156:THR:HA	1:A:236:PHE:O	2.13	0.49
1:A:207:SER:O	1:A:211:ARG:NH2	2.38	0.49
1:A:117:TYR:N	4:A:706:HOH:O	2.33	0.48
1:A:5:LEU:HD21	2:B:440:TYR:HD1	1.78	0.47
1:A:106:GLY:HA2	1:A:255:SER:HB3	1.96	0.47
1:A:269:GLU:HB3	4:A:704:HOH:O	2.14	0.47
1:A:192:LEU:CD1	1:A:205:VAL:HG22	2.43	0.47
1:A:49:VAL:HG23	1:A:74:ALA:HB2	1.97	0.46
1:A:171:TRP:CZ2	1:A:195:VAL:HG21	2.50	0.46
2:B:347:HIS:NE2	2:B:354:GLY:HA3	2.30	0.46
1:A:269:GLU:CB	4:A:704:HOH:O	2.63	0.46
2:B:411:GLU:HG3	4:B:604:HOH:O	2.14	0.46
2:B:332:GLU:O	2:B:333:ASN:CB	2.59	0.45
1:A:193:VAL:O	1:A:204:PHE:N	2.38	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:ILE:HG22	2:B:461:ILE:HD11	1.97	0.45
1:A:170:VAL:O	1:A:245:PRO:HB3	2.17	0.45
1:A:155:MET:O	1:A:237:SER:HA	2.17	0.44
2:B:346:ARG:HH21	2:B:355:THR:HG21	1.83	0.44
2:B:450:ASN:N	2:B:450:ASN:OD1	2.49	0.44
2:B:453:GLU:HG2	2:B:459:PHE:HE2	1.82	0.44
1:A:166:PRO:HA	1:A:228:LEU:O	2.18	0.44
2:B:462:PHE:HB3	2:B:490:ASN:HB2	1.99	0.44
1:A:135:SER:OG	1:A:136:PHE:N	2.46	0.44
2:B:395:GLU:HB3	2:B:398:ILE:HG22	2.00	0.43
1:A:183:THR:HG22	1:A:188:SER:HA	2.01	0.43
1:A:193:VAL:N	1:A:204:PHE:O	2.30	0.43
1:A:169:ILE:HB	1:A:226:LEU:HD23	2.02	0.42
1:A:79:GLU:OE2	1:A:103:ARG:NH2	2.50	0.42
1:A:116:THR:O	1:A:157:LYS:NZ	2.51	0.42
1:A:191:LYS:HA	1:A:239:ASN:HD21	1.84	0.42
1:A:287:ASN:ND2	1:A:300:VAL:O	2.44	0.42
1:A:192:LEU:HD12	1:A:205:VAL:HG22	2.01	0.41
1:A:269:GLU:OE1	4:A:704:HOH:O	2.22	0.41
1:A:305:LEU:HB3	2:B:421:VAL:HG21	2.03	0.41
1:A:37:ASN:OD1	1:A:274:HIS:NE2	2.37	0.41
1:A:171:TRP:CE2	1:A:195:VAL:HG21	2.55	0.41
1:A:172:GLY:HA2	1:A:222:ASP:O	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:LYS:NZ	1:A:68:GLN:O[6_555]	1.97	0.23

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/321 (97%)	295 (95%)	16 (5%)	1 (0%)	36	58
2	B	161/177 (91%)	150 (93%)	11 (7%)	0	100	100
All	All	473/498 (95%)	445 (94%)	27 (6%)	1 (0%)	43	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/270 (97%)	253 (97%)	9 (3%)	32	60
2	B	141/152 (93%)	136 (96%)	5 (4%)	32	59
All	All	403/422 (96%)	389 (96%)	14 (4%)	32	59

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	A	19	LEU
1	A	127	SER
1	A	146	ASN
1	A	155	MET
1	A	205	VAL
1	A	226	LEU
1	A	227	MET
1	A	312	LYS
2	B	331	ILE
2	B	332	GLU
2	B	450	ASN
2	B	454	ASP
2	B	469	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	374	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	601	1	14,14,15	0.26	0	17,19,21	0.44	0
3	NAG	A	602	1	14,14,15	0.29	0	17,19,21	0.51	0
3	NAG	B	501	2	14,14,15	0.27	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	1/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	602	1	-	2/6/23/26	0/1/1/1
3	NAG	B	501	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	NAG	O5-C5-C6-O6
3	A	602	NAG	C4-C5-C6-O6
3	A	601	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/321 (97%)	1.01	53 (16%) 4 3	43, 79, 136, 175	0
2	B	163/177 (92%)	1.28	34 (20%) 2 2	40, 104, 157, 190	0
All	All	477/498 (95%)	1.10	87 (18%) 3 3	40, 86, 149, 190	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	240	GLY	4.9
2	B	447	LEU	4.8
2	B	480	HIS	4.6
1	A	118	SER	4.6
2	B	437	ASP	4.5
1	A	195	VAL	4.4
2	B	380	THR	4.2
1	A	117	TYR	4.2
1	A	186	TYR	4.2
2	B	438	LYS	4.2
2	B	451	ALA	4.2
2	B	483	TYR	4.1
1	A	180	ALA	4.1
1	A	158	SER	4.0
2	B	350	ALA	4.0
1	A	241	ALA	3.9
1	A	236	PHE	3.7
1	A	149	ASN	3.7
2	B	328	ALA	3.7
2	B	488	MET	3.6
1	A	183	THR	3.5
1	A	6	GLY	3.5
1	A	207	SER	3.5
2	B	379	LYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	156	THR	3.2
1	A	128	SER	3.2
1	A	203	SER	3.2
2	B	360	LYS	3.2
2	B	472	SER	3.1
1	A	192	LEU	3.1
1	A	189	GLY	3.0
1	A	114	GLY	3.0
2	B	473	ILE	3.0
1	A	119	GLY	3.0
1	A	190	ASN	2.9
2	B	455	GLY	2.9
1	A	196	GLY	2.8
1	A	122	THR	2.8
1	A	143	LEU	2.8
2	B	373	LEU	2.8
1	A	129	CYS	2.8
1	A	145	SER	2.8
1	A	237	SER	2.7
1	A	261	GLY	2.7
1	A	212	PRO	2.7
1	A	193	VAL	2.7
2	B	433	ASP	2.7
2	B	378	GLU	2.7
2	B	376	LEU	2.6
1	A	172	GLY	2.6
1	A	316	GLU	2.6
2	B	358	ASP	2.6
2	B	388	ASP	2.6
2	B	375	ARG	2.5
1	A	242	PHE	2.5
1	A	199	ASN	2.5
2	B	475	ASN	2.5
2	B	457	GLY	2.5
1	A	146	ASN	2.5
2	B	347	HIS	2.5
2	B	460	GLU	2.5
1	A	3	ILE	2.4
1	A	201	GLN	2.4
1	A	215	ASN	2.4
1	A	150	ALA	2.4
2	B	490	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	181	GLU	2.3
2	B	477	THR	2.3
2	B	474	ARG	2.3
1	A	144	LEU	2.3
2	B	461	ILE	2.3
1	A	200	TYR	2.3
2	B	478	TYR	2.3
1	A	176	SER	2.3
1	A	314	VAL	2.2
2	B	353	GLU	2.2
1	A	120	ILE	2.2
1	A	239	ASN	2.2
1	A	116	THR	2.1
1	A	234	VAL	2.1
1	A	148	ASP	2.1
2	B	345	PHE	2.1
1	A	235	THR	2.1
1	A	159	TYR	2.1
1	A	211	ARG	2.0
1	A	194	THR	2.0
2	B	377	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	602	14/15	0.43	0.14	123,128,133,134	0
3	NAG	A	601	14/15	0.58	0.20	122,129,145,147	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	501	14/15	0.80	0.14	74,84,92,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.