



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:37 PM UTC

PDB ID : 8IDI / pdb_00008idi
Title : Crystal structure of nanobody VHH-T71 with MERS-CoV RBD
Authors : Wang, X.; Tian, L.
Deposited on : 2023-02-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	5% 88% 6% 6%
1	C	137	4% 88% 7% ...
2	B	229	7% 86% 5% 9%
2	D	229	7% 83% 7% 9%
3	E	5	60% 40%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	5	 A horizontal bar chart showing the quality of chain F. The bar is divided into two segments: a green segment on the left representing 20% and a yellow segment on the right representing 80%. The labels '20%' and '80%' are positioned below their respective segments.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VHH-T71.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			979	610	168	198	3			
1	C	131	Total	C	N	O	S	0	0	0
			994	618	171	202	3			

- Molecule 2 is a protein called Spike protein S1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	0	1	0
			1612	1029	256	316	11			
2	D	208	Total	C	N	O	S	0	3	0
			1619	1033	257	318	11			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	589	GLU	-	expression tag	UNP R9UQ53
B	590	HIS	-	expression tag	UNP R9UQ53
B	591	HIS	-	expression tag	UNP R9UQ53
B	592	HIS	-	expression tag	UNP R9UQ53
B	593	HIS	-	expression tag	UNP R9UQ53
B	594	HIS	-	expression tag	UNP R9UQ53
B	595	HIS	-	expression tag	UNP R9UQ53
D	589	GLU	-	expression tag	UNP R9UQ53
D	590	HIS	-	expression tag	UNP R9UQ53
D	591	HIS	-	expression tag	UNP R9UQ53
D	592	HIS	-	expression tag	UNP R9UQ53
D	593	HIS	-	expression tag	UNP R9UQ53
D	594	HIS	-	expression tag	UNP R9UQ53
D	595	HIS	-	expression tag	UNP R9UQ53

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b

eta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	5	Total	C	N	O	0	0	0
			61	34	2	25			
3	F	5	Total	C	N	O	0	0	0
			61	34	2	25			



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	125	Total	O	0	0
			125	125		
4	B	267	Total	O	0	0
			267	267		
4	C	176	Total	O	0	0
			176	176		
4	D	250	Total	O	0	0
			250	250		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: VHH-T71





- Molecule 1: VHH-T71



- Molecule 2: Spike protein S1



- Molecule 2: Spike protein S1



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  60% 40%

MAN1	MAN2	MAN3	MAN4	MAN5
------	------	------	------	------

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  20% 80%

MAN1	MAN2	MAN3	MAN4	MAN5
------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	238.55Å 46.84Å 95.95Å 90.00° 112.52° 90.00°	Depositor
Resolution (Å)	36.55 – 1.90 36.55 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.55-1.90) 99.8 (36.55-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.173 , 0.207 0.176 , 0.209	Depositor DCC
R_{free} test set	3939 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.732	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6144	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/1000 (0.1%)	0.48	0/1357
1	C	0.41	1/1015 (0.1%)	0.50	0/1377
2	B	0.30	0/1654	0.49	0/2259
2	D	0.28	0/1667	0.48	0/2276
All	All	0.35	2/5336 (0.0%)	0.49	0/7269

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	10	GLY	C-O	-6.22	1.15	1.23
1	C	2	VAL	C-O	-5.85	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	979	0	921	3	0
1	C	994	0	937	6	0
2	B	1612	0	1580	8	0
2	D	1619	0	1589	10	0
3	E	61	0	52	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	61	0	52	2	0
4	A	125	0	0	2	0
4	B	267	0	0	2	0
4	C	176	0	0	2	0
4	D	250	0	0	3	0
All	All	6144	0	5131	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:2:NAG:O6	3:F:4:MAN:O2	1.93	0.79
2:B:455:ASP:O	4:B:601:HOH:O	2.06	0.74
2:D:455:ASP:OD2	4:D:601:HOH:O	2.15	0.64
2:D:535:TRP:CD1	2:D:536:GLU:HG2	2.38	0.59
2:D:410:ASN:ND2	4:D:603:HOH:O	2.26	0.51
1:C:47:GLU:OE1	4:C:201:HOH:O	2.20	0.49
1:A:44:LYS:NZ	4:A:203:HOH:O	2.38	0.49
2:B:467:PHE:O	2:B:524:SER:HB2	2.13	0.48
2:B:525:PRO:O	4:B:602:HOH:O	2.20	0.47
1:C:4:LEU:HD11	1:C:99:THR:HG22	1.98	0.46
1:A:4:LEU:O	4:A:201:HOH:O	2.21	0.44
2:D:583:SER:OG	4:D:602:HOH:O	2.19	0.44
2:D:524:SER:HB3	2:D:527:VAL:HG13	1.99	0.44
2:B:414:LEU:HA	1:C:104:ILE:HD11	1.98	0.44
2:D:535:TRP:HD1	2:D:536:GLU:HG2	1.81	0.43
2:B:524:SER:HB3	2:B:527:VAL:HG13	2.01	0.43
1:C:122:GLN:NE2	4:C:203:HOH:O	2.27	0.43
1:C:104:ILE:HG23	1:C:105:ASP:OD1	2.19	0.42
2:B:523:TYR:HB3	2:B:527:VAL:HG11	2.00	0.42
2:D:498:SER:HB3	2:D:534:VAL:HG23	2.00	0.41
2:D:410:ASN:OD1	2:D:412:THR:HG22	2.19	0.41
2:B:407:CYS:O	2:B:584:VAL:HA	2.20	0.41
3:F:2:NAG:HO6	3:F:4:MAN:C2	2.29	0.41
2:D:408:ASN:HA	2:D:585:CYS:O	2.21	0.41
2:D:407:CYS:O	2:D:584:VAL:HA	2.21	0.41
2:B:391:GLY:HA2	1:C:32:HIS:CD2	2.56	0.40
1:A:104:ILE:HG23	1:A:105:ASP:OD1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/137 (93%)	123 (97%)	4 (3%)	0	100	100
1	C	129/137 (94%)	128 (99%)	1 (1%)	0	100	100
2	B	207/229 (90%)	205 (99%)	1 (0%)	1 (0%)	24	16
2	D	209/229 (91%)	203 (97%)	5 (2%)	1 (0%)	24	16
All	All	672/732 (92%)	659 (98%)	11 (2%)	2 (0%)	36	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	428	ILE
2	D	428	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	103/111 (93%)	100 (97%)	3 (3%)	37	31
1	C	105/111 (95%)	101 (96%)	4 (4%)	29	21
2	B	191/207 (92%)	190 (100%)	1 (0%)	81	84
2	D	193/207 (93%)	191 (99%)	2 (1%)	68	70
All	All	592/636 (93%)	582 (98%)	10 (2%)	53	52

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	VAL
1	A	5	GLN
1	A	27	GLU
2	B	579	THR
1	C	0	SER
1	C	1	GLN
1	C	66	LYS
1	C	104	ILE
2	D	533	THR
2	D	579	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
2	B	408	ASN
2	B	436	ASN
2	B	475	ASN
2	B	516	GLN
2	B	566	GLN
2	D	408	ASN
2	D	436	ASN
2	D	566	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	E	1	2,3	14,14,15	0.43	0	17,19,21	0.40	0
3	NAG	E	2	3	14,14,15	0.22	0	17,19,21	0.46	0
3	BMA	E	3	3	11,11,12	0.62	0	15,15,17	0.76	0
3	MAN	E	4	3	11,11,12	0.81	1 (9%)	15,15,17	1.09	2 (13%)
3	MAN	E	5	3	11,11,12	0.77	0	15,15,17	0.99	1 (6%)
3	NAG	F	1	2,3	14,14,15	0.74	1 (7%)	17,19,21	1.39	2 (11%)
3	NAG	F	2	3	14,14,15	0.31	0	17,19,21	0.37	0
3	BMA	F	3	3	11,11,12	0.66	0	15,15,17	0.73	0
3	MAN	F	4	3	11,11,12	0.35	0	15,15,17	1.06	0
3	MAN	F	5	3	11,11,12	0.78	0	15,15,17	1.03	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1
3	MAN	E	5	3	-	1/2/19/22	0/1/1/1
3	NAG	F	1	2,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	BMA	F	3	3	-	1/2/19/22	0/1/1/1
3	MAN	F	4	3	-	2/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	1	NAG	C1-C2	2.32	1.55	1.52
3	E	4	MAN	O5-C1	-2.00	1.40	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	NAG	O3-C3-C2	-3.68	101.75	109.40
3	E	4	MAN	C1-O5-C5	2.95	116.14	112.19
3	E	5	MAN	C1-O5-C5	2.54	115.59	112.19
3	F	1	NAG	C1-O5-C5	2.35	115.33	112.19
3	F	5	MAN	C1-O5-C5	2.23	115.18	112.19
3	E	4	MAN	O2-C2-C3	-2.12	105.75	110.15
3	F	5	MAN	O2-C2-C3	-2.11	105.79	110.15

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	4	MAN	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6
3	F	3	BMA	O5-C5-C6-O6
3	F	4	MAN	C4-C5-C6-O6
3	F	1	NAG	C4-C5-C6-O6
3	E	5	MAN	O5-C5-C6-O6
3	F	1	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	2	NAG	2	0
3	F	4	MAN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/137 (94%)	0.33	7 (5%) 31 33	20, 27, 50, 93	0
1	C	131/137 (95%)	-0.07	6 (4%) 37 40	15, 22, 42, 67	0
2	B	208/229 (90%)	0.02	17 (8%) 17 18	13, 20, 44, 100	1 (0%)
2	D	208/229 (90%)	0.07	15 (7%) 21 23	11, 23, 45, 113	3 (1%)
All	All	676/732 (92%)	0.08	45 (6%) 24 25	11, 23, 46, 113	4 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	579	THR	7.7
1	A	2	VAL	6.6
2	B	460	SER	6.0
2	B	473	PHE	5.8
2	D	578	GLY	5.6
2	D	580	ASP	5.5
1	C	0	SER	5.4
2	B	580	ASP	5.2
2	B	579	THR	5.0
2	B	458	VAL	4.4
1	A	28	TYR	4.1
2	D	512	THR	4.1
2	D	581	THR	4.1
2	D	511	ARG	3.6
2	B	511	ARG	3.5
1	C	27	GLU	3.5
1	C	130	SER	3.4
1	C	1	GLN	3.4
2	D	510	ASP	3.4
2	D	535	TRP	3.3
2	B	381	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	26	SER	2.9
2	B	456	LEU	2.9
2	B	588	LEU	2.8
2	D	583	SER	2.7
2	D	540	TYR	2.7
2	D	506	LEU	2.7
1	A	27	GLU	2.7
1	A	66	LYS	2.6
2	B	583	SER	2.6
2	D	508	SER	2.6
2	B	486	HIS	2.6
2	D	588	LEU	2.6
1	C	26	SER	2.5
2	B	581	THR	2.5
2	D	381	VAL	2.4
2	D	582	ASN	2.4
1	C	129	SER	2.3
1	A	3	GLN	2.3
2	B	457	SER	2.2
2	B	461	ALA	2.2
2	B	547	PRO	2.2
1	A	130	SER	2.1
2	B	548	LEU	2.1
2	B	578	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	F	3	11/12	0.26	0.17	100,105,107,109	0
3	MAN	F	4	11/12	0.44	0.33	30,30,30,30	0
3	BMA	E	3	11/12	0.48	0.17	97,101,103,106	0
3	MAN	E	4	11/12	0.51	0.27	64,96,110,116	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAN	F	5	11/12	0.51	0.24	86,96,101,105	0
3	MAN	E	5	11/12	0.59	0.21	80,87,99,104	0
3	NAG	E	2	14/15	0.66	0.18	72,78,86,93	0
3	NAG	F	2	14/15	0.69	0.17	73,80,90,99	0
3	NAG	F	1	14/15	0.83	0.14	26,42,73,81	0
3	NAG	E	1	14/15	0.90	0.10	22,40,54,73	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.