



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 07:00 PM UTC

PDB ID : 5IFM / pdb_00005ifm
Title : Human NONO (p54nrb) Homodimer
Authors : Knott, G.J.; Bond, C.S.
Deposited on : 2016-02-26
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	
1	B	261	
1	C	261	
1	D	261	
1	E	261	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	261	
1	G	261	
1	H	261	
1	I	261	
1	J	261	
1	K	261	
1	L	261	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	D	401	-	-	X	-
5	GOL	B	402	-	-	X	-
5	GOL	C	405	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 25317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-POU domain-containing octamer-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			2065	1301	369	386	9			
1	B	259	Total	C	N	O	S	0	1	0
			2114	1333	380	390	11			
1	C	256	Total	C	N	O	S	0	1	0
			2090	1316	374	391	9			
1	D	257	Total	C	N	O	S	0	1	0
			2099	1320	379	391	9			
1	E	256	Total	C	N	O	S	0	0	0
			2084	1312	374	389	9			
1	F	255	Total	C	N	O	S	0	0	0
			2075	1307	373	386	9			
1	G	255	Total	C	N	O	S	0	1	0
			2084	1312	373	390	9			
1	H	257	Total	C	N	O	S	0	0	0
			2090	1317	375	388	10			
1	I	255	Total	C	N	O	S	0	0	0
			2075	1307	372	387	9			
1	J	259	Total	C	N	O	S	0	0	0
			2106	1328	377	390	11			
1	K	259	Total	C	N	O	S	0	0	0
			2106	1328	377	390	11			
1	L	255	Total	C	N	O	S	0	0	0
			2075	1307	372	387	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	GLY	-	expression tag	UNP Q15233
B	52	GLY	-	expression tag	UNP Q15233
C	52	GLY	-	expression tag	UNP Q15233
D	52	GLY	-	expression tag	UNP Q15233
E	52	GLY	-	expression tag	UNP Q15233

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	52	GLY	-	expression tag	UNP Q15233
G	52	GLY	-	expression tag	UNP Q15233
H	52	GLY	-	expression tag	UNP Q15233
I	52	GLY	-	expression tag	UNP Q15233
J	52	GLY	-	expression tag	UNP Q15233
K	52	GLY	-	expression tag	UNP Q15233
L	52	GLY	-	expression tag	UNP Q15233

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	C	2	Total Cl 2 2	0	0
2	D	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0
2	F	2	Total Cl 2 2	0	0
2	G	1	Total Cl 1 1	0	0
2	H	1	Total Cl 1 1	0	0
2	L	1	Total Cl 1 1	0	0

- Molecule 3 is PROLINE (CCD ID: PRO) (formula: C₅H₉NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	5	1	2		
3	C	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		
4	B	1	Total	K	0	0
			1	1		
4	C	1	Total	K	0	0
			1	1		
4	D	1	Total	K	0	0
			1	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	33	Total	O	0	0
			33	33		
6	B	26	Total	O	0	0
			26	26		
6	C	24	Total	O	0	0
			24	24		
6	D	37	Total	O	0	0
			37	37		
6	E	15	Total	O	0	0
			15	15		
6	F	7	Total	O	0	0
			7	7		
6	G	19	Total	O	0	0
			19	19		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	10	Total 10	O 10	0	0
6	I	5	Total 5	O 5	0	0
6	J	4	Total 4	O 4	0	0
6	K	7	Total 7	O 7	0	0
6	L	7	Total 7	O 7	0	0

- Molecule 1: Non-POU domain-containing octamer-binding protein









- Molecule 1: Non-POU domain-containing octamer-binding protein




- Molecule 1: Non-POU domain-containing octamer-binding protein




- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein





- Molecule 1: Non-POU domain-containing octamer-binding protein









- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.15Å 407.18Å 68.96Å 90.00° 97.75° 90.00°	Depositor
Resolution (Å)	48.15 – 2.60 48.15 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (48.15-2.60) 98.8 (48.15-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.61Å)	Xtriage
Refinement program	BUSTER-TNT 2.10.0	Depositor
R, R_{free}	0.197 , 0.234 0.213 , 0.254	Depositor DCC
R_{free} test set	5521 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	48.7	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.038 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	25317	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, K, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/2104	1.35	9/2823 (0.3%)
1	B	0.91	1/2157 (0.0%)	1.34	10/2893 (0.3%)
1	C	0.91	1/2133 (0.0%)	1.43	19/2862 (0.7%)
1	D	0.96	1/2139 (0.0%)	1.43	19/2869 (0.7%)
1	E	0.82	0/2124	1.38	11/2850 (0.4%)
1	F	0.82	0/2115	1.36	11/2838 (0.4%)
1	G	0.83	0/2124	1.36	5/2850 (0.2%)
1	H	0.83	0/2130	1.43	17/2858 (0.6%)
1	I	0.80	0/2115	1.36	8/2838 (0.3%)
1	J	0.78	0/2146	1.33	7/2879 (0.2%)
1	K	0.82	0/2146	1.38	13/2879 (0.5%)
1	L	0.83	0/2115	1.39	8/2838 (0.3%)
All	All	0.85	3/25548 (0.0%)	1.38	137/34277 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	304	ARG	C-N	5.89	1.41	1.33
1	D	300	MET	SD-CE	-5.59	1.65	1.79
1	B	82	PRO	CA-C	5.46	1.54	1.51

The worst 5 of 137 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	109	LYS	N-CA-C	7.52	122.49	113.16
1	K	58	ASP	CA-CB-CG	7.19	119.79	112.60
1	B	84	ASP	CA-CB-CG	6.87	119.47	112.60
1	D	109	LYS	N-CA-C	6.76	121.38	113.20
1	D	182	ASP	CA-C-N	6.73	130.21	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	2064	23	0
1	B	2114	0	2124	37	0
1	C	2090	0	2085	33	0
1	D	2099	0	2094	33	0
1	E	2084	0	2079	25	0
1	F	2075	0	2073	20	0
1	G	2084	0	2076	22	0
1	H	2090	0	2091	25	0
1	I	2075	0	2071	20	0
1	J	2106	0	2111	14	0
1	K	2106	0	2111	17	0
1	L	2075	0	2071	18	0
2	A	1	0	0	1	0
2	C	2	0	0	1	0
2	D	1	0	0	2	0
2	E	1	0	0	1	0
2	F	2	0	0	1	0
2	G	1	0	0	1	0
2	H	1	0	0	1	0
2	L	1	0	0	0	0
3	A	8	0	7	2	0
3	C	8	0	7	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	12	0	16	8	0
5	C	12	0	16	7	0
5	D	6	0	8	2	0
6	A	33	0	0	1	0
6	B	26	0	0	0	0
6	C	24	0	0	1	0
6	D	37	0	0	0	0
6	E	15	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	F	7	0	0	0	0
6	G	19	0	0	0	0
6	H	10	0	0	0	0
6	I	5	0	0	0	0
6	J	4	0	0	0	0
6	K	7	0	0	0	0
6	L	7	0	0	0	0
All	All	25317	0	25104	208	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:285:VAL:HG21	1:L:285:VAL:HG21	1.43	0.98
1:B:84:ASP:HB2	1:B:135:ARG:HH22	1.31	0.95
1:C:61:ASN:HA	5:C:405:GOL:H32	1.55	0.88
1:I:285:VAL:HG21	1:J:285:VAL:HG21	1.57	0.84
1:L:156:PRO:HD2	1:L:159:VAL:HG21	1.58	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/261 (97%)	242 (96%)	10 (4%)	0	100	100
1	B	258/261 (99%)	249 (96%)	9 (4%)	0	100	100
1	C	255/261 (98%)	248 (97%)	5 (2%)	2 (1%)	16	34
1	D	256/261 (98%)	248 (97%)	6 (2%)	2 (1%)	16	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	254/261 (97%)	249 (98%)	5 (2%)	0	100	100
1	F	253/261 (97%)	246 (97%)	7 (3%)	0	100	100
1	G	254/261 (97%)	246 (97%)	8 (3%)	0	100	100
1	H	255/261 (98%)	246 (96%)	9 (4%)	0	100	100
1	I	253/261 (97%)	245 (97%)	8 (3%)	0	100	100
1	J	257/261 (98%)	250 (97%)	6 (2%)	1 (0%)	30	51
1	K	257/261 (98%)	249 (97%)	7 (3%)	1 (0%)	30	51
1	L	253/261 (97%)	244 (96%)	9 (4%)	0	100	100
All	All	3057/3132 (98%)	2962 (97%)	89 (3%)	6 (0%)	43	66

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	85	ILE
1	D	53	GLU
1	J	107	LYS
1	C	107	LYS
1	D	107	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/228 (97%)	213 (96%)	9 (4%)	27	54
1	B	228/228 (100%)	222 (97%)	6 (3%)	40	68
1	C	225/228 (99%)	212 (94%)	13 (6%)	18	39
1	D	225/228 (99%)	206 (92%)	19 (8%)	10	23
1	E	224/228 (98%)	214 (96%)	10 (4%)	24	50
1	F	223/228 (98%)	206 (92%)	17 (8%)	12	27
1	G	224/228 (98%)	205 (92%)	19 (8%)	10	22
1	H	225/228 (99%)	206 (92%)	19 (8%)	10	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	223/228 (98%)	207 (93%)	16 (7%)	13	30
1	J	227/228 (100%)	212 (93%)	15 (7%)	15	34
1	K	227/228 (100%)	207 (91%)	20 (9%)	9	21
1	L	223/228 (98%)	209 (94%)	14 (6%)	16	36
All	All	2696/2736 (98%)	2519 (93%)	177 (7%)	15	34

5 of 177 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	88	GLU
1	K	58	ASP
1	I	115	ARG
1	J	90	MET
1	K	107	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	61	ASN
1	F	173	GLN
1	L	154	ASN
1	I	61	ASN
1	J	61	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 21 ligands modelled in this entry, 14 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	D	402	-	5,5,5	0.25	0	5,5,5	0.44	0
5	GOL	C	404	-	5,5,5	0.31	0	5,5,5	0.70	0
3	PRO	C	403	-	8,8,8	0.89	0	10,10,10	1.01	0
5	GOL	B	401	-	5,5,5	0.39	0	5,5,5	0.92	0
5	GOL	C	405	-	5,5,5	0.25	0	5,5,5	1.05	0
3	PRO	A	402	-	8,8,8	0.96	0	10,10,10	1.16	1 (10%)
5	GOL	B	402	-	5,5,5	0.30	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	D	402	-	-	4/4/4/4	-
5	GOL	C	404	-	-	4/4/4/4	-
3	PRO	C	403	-	-	2/4/11/11	0/1/1/1
5	GOL	B	401	-	-	2/4/4/4	-
5	GOL	C	405	-	-	2/4/4/4	-
3	PRO	A	402	-	-	2/4/11/11	0/1/1/1
5	GOL	B	402	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	PRO	C-CA-N	2.02	114.73	106.84

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	402	GOL	O1-C1-C2-C3
5	C	405	GOL	C1-C2-C3-O3
5	D	402	GOL	O1-C1-C2-C3
5	D	402	GOL	C1-C2-C3-O3
5	C	405	GOL	O2-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	402	GOL	2	0
5	C	404	GOL	3	0
3	C	403	PRO	1	0
5	B	401	GOL	1	0
5	C	405	GOL	4	0
3	A	402	PRO	2	0
5	B	402	GOL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/261 (97%)	-0.08	1 (0%) 88 86	22, 50, 81, 113	0
1	B	259/261 (99%)	0.14	11 (4%) 40 35	20, 49, 96, 148	1 (0%)
1	C	256/261 (98%)	-0.09	6 (2%) 61 55	20, 46, 86, 132	1 (0%)
1	D	257/261 (98%)	-0.09	5 (1%) 66 61	16, 46, 100, 148	1 (0%)
1	E	256/261 (98%)	0.18	5 (1%) 65 60	28, 63, 116, 138	0
1	F	255/261 (97%)	0.21	4 (1%) 70 66	35, 64, 119, 140	0
1	G	255/261 (97%)	0.08	6 (2%) 59 54	20, 55, 103, 129	1 (0%)
1	H	257/261 (98%)	0.20	5 (1%) 66 61	35, 63, 103, 126	0
1	I	255/261 (97%)	0.36	4 (1%) 70 66	41, 71, 123, 136	0
1	J	259/261 (99%)	0.68	15 (5%) 29 23	45, 80, 143, 165	0
1	K	259/261 (99%)	0.43	10 (3%) 43 38	37, 66, 108, 124	0
1	L	255/261 (97%)	0.33	7 (2%) 56 50	38, 72, 109, 141	0
All	All	3077/3132 (98%)	0.20	79 (2%) 57 51	16, 60, 115, 165	4 (0%)

The worst 5 of 79 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	309	VAL	5.0
1	B	311	LEU	4.8
1	B	310	MET	4.7
1	B	307	HIS	4.6
1	J	311	LEU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PRO	C	403	8/8	0.77	0.17	64,67,107,128	0
5	GOL	C	404	6/6	0.80	0.19	49,51,57,61	0
5	GOL	B	402	6/6	0.85	0.20	48,55,58,59	0
2	CL	L	401	1/1	0.88	0.11	73,73,73,73	0
5	GOL	B	401	6/6	0.88	0.22	38,41,46,50	0
5	GOL	D	402	6/6	0.88	0.17	56,69,71,72	0
3	PRO	A	402	8/8	0.89	0.17	45,47,62,82	0
4	K	C	406	1/1	0.89	0.12	76,76,76,76	0
4	K	A	403	1/1	0.90	0.10	73,73,73,73	0
5	GOL	C	405	6/6	0.90	0.17	30,42,46,47	0
4	K	D	403	1/1	0.90	0.10	71,71,71,71	0
2	CL	D	401	1/1	0.91	0.13	90,90,90,90	0
4	K	B	403	1/1	0.93	0.10	71,71,71,71	0
2	CL	H	401	1/1	0.93	0.08	63,63,63,63	0
2	CL	G	401	1/1	0.94	0.10	70,70,70,70	0
2	CL	F	402	1/1	0.94	0.11	79,79,79,79	0
2	CL	F	401	1/1	0.95	0.14	64,64,64,64	0
2	CL	C	402	1/1	0.96	0.07	59,59,59,59	0
2	CL	C	401	1/1	0.96	0.08	56,56,56,56	0
2	CL	A	401	1/1	0.97	0.09	48,48,48,48	0
2	CL	E	401	1/1	0.98	0.06	52,52,52,52	0

6.5 Other polymers ⓘ

There are no such residues in this entry.