



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 07:43 PM UTC

PDB ID : 3IGV / pdb_00003igv
Title : Crystal structure of HCV NS5B polymerase with a novel monocyclic dihydro-pyridinone inhibitor
Authors : Zhao, Q.; Showalter, R.E.; Han, Q.; Kissinger, C.R.
Deposited on : 2009-07-28
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	578	
1	B	578	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			
1	B	557	Total	C	N	O	S	0	0	0
			4334	2735	764	804	31			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	544	GLN	ARG	engineered mutation	UNP P26663
A	571	LEU	-	expression tag	UNP P26663
A	572	GLU	-	expression tag	UNP P26663
A	573	HIS	-	expression tag	UNP P26663
A	574	HIS	-	expression tag	UNP P26663
A	575	HIS	-	expression tag	UNP P26663
A	576	HIS	-	expression tag	UNP P26663
A	577	HIS	-	expression tag	UNP P26663
A	578	HIS	-	expression tag	UNP P26663
B	544	GLN	ARG	engineered mutation	UNP P26663
B	571	LEU	-	expression tag	UNP P26663
B	572	GLU	-	expression tag	UNP P26663
B	573	HIS	-	expression tag	UNP P26663
B	574	HIS	-	expression tag	UNP P26663
B	575	HIS	-	expression tag	UNP P26663
B	576	HIS	-	expression tag	UNP P26663
B	577	HIS	-	expression tag	UNP P26663
B	578	HIS	-	expression tag	UNP P26663

- Molecule 2 is N-{3-[(6S)-6-ethyl-1-(4-fluorobenzyl)-4-hydroxy-2-oxo-1,2,5,6-tetrahydropyridin-3-yl]-1,1-dioxido-2H-1,2,4-benzothiadiazin-7-yl}methanesulfonamide (CCD ID: B80) (formula: C₂₂H₂₃FN₄O₆S₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			35	22	1	4	6	2		
2	B	1	Total	C	F	N	O	S	0	0
			35	22	1	4	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		
3	B	63	Total	O	0	0
			63	63		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-DIRECTED RNA POLYMERASE



• Molecule 1: RNA-DIRECTED RNA POLYMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.91Å 107.22Å 126.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.55 – 2.60 28.55 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.8 (28.55-2.60) 96.8 (28.55-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.245 , 0.301 0.245 , 0.298	Depositor DCC
R_{free} test set	1787 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 20.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8860	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2398e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B80

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/4428	0.79	2/6009 (0.0%)
1	B	0.46	0/4428	0.79	0/6009
All	All	0.46	0/8856	0.79	2/12018 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	539	ILE	CA-C-N	5.09	126.20	119.84
1	A	539	ILE	C-N-CA	5.09	126.20	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4334	0	4354	23	0
1	B	4334	0	4354	25	0
2	A	35	0	22	1	0
2	B	35	0	22	1	0
3	A	59	0	0	2	0
3	B	63	0	0	2	0
All	All	8860	0	8752	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:THR:HG21	1:B:270:LYS:HE2	1.65	0.77
1:B:201:VAL:HG23	1:B:384:LEU:HG	1.75	0.69
1:A:144:VAL:HB	1:A:394:ARG:HG2	1.79	0.63
1:B:201:VAL:CG2	1:B:384:LEU:HG	2.29	0.62
1:A:451:CYS:HB2	3:A:636:HOH:O	2.00	0.61
1:A:539:ILE:HB	1:A:542:ALA:HB2	1.84	0.59
1:A:58:GLN:HG2	1:A:60:LEU:HD21	1.83	0.59
1:B:114:LYS:HE2	3:B:617:HOH:O	2.03	0.59
1:A:187:MET:HE1	1:A:292:THR:HG22	1.88	0.56
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.89	0.55
1:B:314:LEU:HB3	1:B:321:VAL:HG13	1.88	0.54
1:A:201:VAL:HG22	1:A:384:LEU:HG	1.90	0.53
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.91	0.52
1:B:187:MET:HE2	1:B:293:LEU:HD23	1.93	0.50
1:B:445:CYS:SG	1:B:454:ILE:HD12	2.51	0.50
1:A:77:THR:O	1:B:24:ASN:HB2	2.11	0.50
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.94	0.50
1:B:119:ILE:HD13	1:B:169:VAL:HG11	1.93	0.49
1:B:472:PHE:HE1	1:B:525:LEU:HD23	1.76	0.49
1:A:5:THR:O	1:A:275:GLY:HA3	2.13	0.49
1:A:119:ILE:HD13	1:A:169:VAL:HG11	1.95	0.48
1:B:11:ILE:HD12	1:B:45:ALA:HB1	1.95	0.47
1:A:223:CYS:HB3	3:A:623:HOH:O	2.14	0.46
1:B:60:LEU:HD13	1:B:64:TYR:CE2	2.50	0.46
1:B:314:LEU:HB3	1:B:321:VAL:CG1	2.46	0.45
1:A:12:THR:HG21	1:A:270:LYS:HD2	1.99	0.45
1:A:466:LEU:HD22	1:A:551:PHE:HE2	1.82	0.45
1:B:217:PHE:HA	3:B:632:HOH:O	2.16	0.45
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.99	0.44
1:A:414:MET:HE2	2:A:579:B80:H17	2.00	0.44
1:B:30:LEU:HD23	1:B:396:ALA:HA	2.00	0.44
1:A:496:PRO:HG2	1:A:499:VAL:HG23	2.00	0.43
1:A:314:LEU:HB3	1:A:321:VAL:HG13	2.00	0.43
1:A:314:LEU:HB3	1:A:321:VAL:CG1	2.48	0.43
2:B:579:B80:O32	2:B:579:B80:N24	2.52	0.43
1:B:227:THR:HB	1:B:347:SER:O	2.19	0.43
1:B:215:MET:HB2	1:B:326:SER:HB2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ASN:HD22	1:A:117:ASN:HA	1.69	0.42
1:B:5:THR:O	1:B:275:GLY:HA3	2.19	0.42
1:B:230:GLU:HG3	1:B:262:ILE:HG23	2.01	0.41
1:B:319:ASP:CG	1:B:366:CYS:H	2.29	0.41
1:A:390:THR:HB	1:A:391:PRO:HD3	2.03	0.41
1:B:390:THR:HB	1:B:391:PRO:HD3	2.03	0.41
1:A:458:ASP:HB3	1:A:545:LEU:HD11	2.03	0.41
1:B:17:GLU:HB3	1:B:41:THR:HG22	2.03	0.41
1:A:537:THR:HB	1:A:538:PRO:HD2	2.02	0.40
1:B:191:TYR:CZ	1:B:193:PHE:HB2	2.57	0.40
1:A:201:VAL:CG2	1:A:384:LEU:HG	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/578 (96%)	540 (98%)	13 (2%)	0	100	100
1	B	553/578 (96%)	538 (97%)	15 (3%)	0	100	100
All	All	1106/1156 (96%)	1078 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/493 (96%)	470 (99%)	5 (1%)	65	84
1	B	475/493 (96%)	471 (99%)	4 (1%)	73	88
All	All	950/986 (96%)	941 (99%)	9 (1%)	70	87

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	148	GLN
1	A	262	ILE
1	A	352	ASP
1	A	381	VAL
1	B	159	LEU
1	B	201	VAL
1	B	352	ASP
1	B	461	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	117	ASN
1	A	142	ASN
1	A	272	GLN
1	A	330	GLN
1	A	355	GLN
1	A	374	HIS
1	A	411	ASN
1	A	483	ASN
1	B	110	ASN
1	B	117	ASN
1	B	142	ASN
1	B	374	HIS
1	B	411	ASN
1	B	436	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	B80	B	579	-	38,38,38	3.50	12 (31%)	46,58,58	2.67	8 (17%)
2	B80	A	579	-	38,38,38	3.37	10 (26%)	46,58,58	2.56	11 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B80	B	579	-	-	0/11/50/50	0/4/4/4
2	B80	A	579	-	-	1/11/50/50	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	579	B80	C12-S34	-17.01	1.60	1.75
2	A	579	B80	C12-S34	-16.21	1.61	1.75
2	B	579	B80	S35-N26	-7.71	1.53	1.63
2	A	579	B80	S35-N26	-7.25	1.53	1.63
2	B	579	B80	O29-S34	4.29	1.48	1.43
2	B	579	B80	O28-S34	4.06	1.48	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	579	B80	C20-S35	-3.85	1.66	1.75
2	A	579	B80	O28-S34	3.83	1.48	1.43
2	A	579	B80	C10-N26	-3.80	1.37	1.43
2	B	579	B80	C20-S35	-3.79	1.66	1.75
2	B	579	B80	C10-N26	-3.74	1.37	1.43
2	A	579	B80	O29-S34	3.54	1.47	1.43
2	A	579	B80	O31-S35	3.32	1.49	1.43
2	A	579	B80	O30-S35	3.08	1.48	1.43
2	B	579	B80	O31-S35	3.03	1.48	1.43
2	B	579	B80	O30-S35	2.98	1.48	1.43
2	A	579	B80	C13-C15	2.82	1.51	1.47
2	A	579	B80	C15-N23	2.70	1.36	1.30
2	B	579	B80	C9-N23	-2.34	1.35	1.39
2	B	579	B80	S34-N24	-2.33	1.60	1.62
2	B	579	B80	C13-C15	2.24	1.50	1.47
2	B	579	B80	C15-N23	2.21	1.35	1.30

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	579	B80	O29-S34-O28	-9.44	108.84	118.44
2	B	579	B80	O29-S34-O28	-9.32	108.95	118.44
2	B	579	B80	C12-S34-N24	8.79	109.62	102.13
2	B	579	B80	O30-S35-O31	-7.80	108.42	118.87
2	A	579	B80	O30-S35-O31	-7.76	108.47	118.87
2	A	579	B80	C12-S34-N24	7.53	108.54	102.13
2	B	579	B80	C9-N23-C15	4.57	125.38	117.58
2	A	579	B80	C9-N23-C15	4.02	124.44	117.58
2	B	579	B80	O32-C14-C17	3.64	119.28	113.29
2	A	579	B80	O32-C14-C17	3.09	118.37	113.29
2	B	579	B80	C8-C21-N25	-2.85	108.13	113.55
2	B	579	B80	C20-S35-N26	2.61	109.50	106.56
2	A	579	B80	O31-S35-N26	2.55	112.19	107.10
2	A	579	B80	C22-C18-C17	-2.28	110.41	113.02
2	A	579	B80	C8-C21-N25	-2.14	109.48	113.55
2	A	579	B80	O27-C16-C13	-2.10	120.67	125.16
2	A	579	B80	C7-C12-S34	2.08	121.48	119.82
2	A	579	B80	O29-S34-N24	2.07	109.94	108.26
2	B	579	B80	C7-C12-S34	2.05	121.46	119.82

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	579	B80	C17-C18-C22-C19


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	579	B80	1	0
2	A	579	B80	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/578 (96%)	0.40	17 (3%) 51 45	33, 48, 62, 94	0
1	B	557/578 (96%)	0.45	20 (3%) 46 40	37, 49, 67, 93	0
All	All	1114/1156 (96%)	0.42	37 (3%) 49 43	33, 49, 65, 94	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	532	THR	4.3
1	B	534	LEU	4.2
1	A	532	THR	4.1
1	A	549	GLY	4.0
1	B	535	LYS	3.7
1	B	542	ALA	3.5
1	A	541	ALA	3.3
1	B	213	ASN	3.3
1	B	541	ALA	3.1
1	B	549	GLY	3.1
1	B	148	GLN	3.1
1	B	533	LYS	3.0
1	B	536	LEU	3.0
1	B	352	ASP	2.9
1	A	533	LYS	2.9
1	B	545	LEU	2.8
1	A	545	LEU	2.8
1	A	535	LYS	2.8
1	A	543	SER	2.7
1	A	16	ALA	2.7
1	B	544	GLN	2.6
1	B	531	LYS	2.5
1	A	542	ALA	2.3
1	A	536	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	270	LYS	2.3
1	B	15	ALA	2.2
1	B	353	PRO	2.2
1	A	223	CYS	2.2
1	B	539	ILE	2.1
1	B	116	VAL	2.1
1	A	148	GLN	2.0
1	B	95	HIS	2.0
1	B	538	PRO	2.0
1	A	355	GLN	2.0
1	A	534	LEU	2.0
1	A	95	HIS	2.0
1	A	402	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	B80	B	579	35/35	0.94	0.10	37,40,41,42	0
2	B80	A	579	35/35	0.95	0.08	33,42,43,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B80 B 579:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around B80 A 579:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.