



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 08:25 AM UTC

PDB ID : 4IGQ / pdb_00004igq
Title : Histone H3 Lysine 4 Demethylating Rice JMJ703 in complex with methylated H3K4 substrate
Authors : Chen, Q.F.; Chen, X.S.; Wang, Q.; Zhang, F.B.; Lou, Z.Y.; Zhang, Q.F.; Zhou, D.X.
Deposited on : 2012-12-17
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	4% 56% 16% • 24%
2	B	3	67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	M3L	B	4	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Os05g0196500 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	0	0
			2228	1445	366	407	10			


- Molecule 2 is a protein called mathylated H3K4 substrate.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			28	18	5	5			

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is N-OXALYLGLYCINE (CCD ID: OGA) (formula: C₄H₅NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			10	4	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	141	Total	O	0	0
			141	141		

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	91.42Å 91.42Å 75.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.59 – 2.35 39.59 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.59-2.35) 99.8 (39.59-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.27Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.181 , 0.235 0.180 , 0.232	Depositor DCC
R_{free} test set	843 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2408	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: M3L, FE, OGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/2300	0.89	5/3126 (0.2%)
2	B	1.54	1/15 (6.7%)	4.44	2/19 (10.5%)
All	All	0.57	1/2315 (0.0%)	0.95	7/3145 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	M3L	C-N	5.44	1.41	1.33

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	M3L	CA-C-N	-12.99	98.32	121.70
2	B	4	M3L	C-N-CA	-12.99	98.32	121.70
1	A	351	GLN	CB-CA-C	-10.10	90.92	110.10
1	A	140	LYS	N-CA-C	7.53	120.52	109.07
1	A	141	TRP	N-CA-C	6.66	120.15	110.42
1	A	351	GLN	N-CA-C	6.20	128.35	111.00
1	A	352	SER	N-CA-C	-6.10	100.20	109.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2105	82	0
2	B	28	0	32	39	0
3	A	1	0	0	0	0
4	A	10	0	3	0	0
5	A	141	0	0	6	0
All	All	2408	0	2140	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:381:TRP:CE3	2:B:3:THR:HG22	1.54	1.40
1:A:381:TRP:CE3	2:B:3:THR:CG2	2.16	1.25
1:A:381:TRP:CD2	2:B:3:THR:HG22	1.85	1.12
1:A:381:TRP:CD2	2:B:3:THR:CG2	2.41	1.03
1:A:494:ALA:HB1	2:B:4:M3L:HM23	1.42	0.99
1:A:381:TRP:CZ3	2:B:3:THR:HG22	2.02	0.94
1:A:381:TRP:CG	2:B:3:THR:HB	2.05	0.92
1:A:381:TRP:CE3	2:B:3:THR:HG21	2.05	0.91
2:B:4:M3L:O	2:B:5:GLN:C	2.10	0.89
1:A:194:PRO:O	5:A:692:HOH:O	1.91	0.87
1:A:494:ALA:HB1	2:B:4:M3L:CM2	2.05	0.85
1:A:350:ALA:N	1:A:351:GLN:HB2	1.92	0.84
1:A:324:ASP:HA	2:B:3:THR:CG2	2.12	0.80
1:A:494:ALA:CB	2:B:4:M3L:HM23	2.13	0.77
1:A:324:ASP:HA	2:B:3:THR:OG1	1.83	0.77
2:B:4:M3L:HB2	2:B:4:M3L:HM13	1.71	0.71
1:A:361:ARG:HB3	1:A:361:ARG:HH11	1.55	0.70
1:A:287:PHE:O	5:A:613:HOH:O	2.10	0.69
1:A:379:VAL:HG22	1:A:380:PRO:HD2	1.74	0.69
1:A:435:GLU:HG3	5:A:694:HOH:O	1.91	0.69
1:A:162:GLU:O	1:A:165:GLU:HG2	1.95	0.67
1:A:326:GLU:HB3	1:A:381:TRP:HZ3	1.62	0.65
1:A:402:SER:OG	2:B:4:M3L:HM22	1.98	0.64
1:A:164:PHE:HE2	1:A:192:TRP:CD1	2.15	0.64
1:A:324:ASP:CA	2:B:3:THR:HG23	2.28	0.64
1:A:275:PHE:HE1	1:A:320:ILE:HG22	1.63	0.63
1:A:161:GLU:OE2	1:A:191:SER:HB3	1.98	0.63
1:A:381:TRP:CD2	2:B:3:THR:CB	2.82	0.63
2:B:3:THR:O	2:B:4:M3L:C	2.47	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:TRP:CH2	1:A:194:PRO:HG3	2.35	0.62
1:A:401:TYR:HB2	1:A:497:VAL:HG13	1.81	0.61
1:A:361:ARG:HB3	1:A:361:ARG:NH1	2.15	0.61
1:A:383:TYR:OH	2:B:4:M3L:HD2	2.01	0.60
1:A:381:TRP:CD2	2:B:3:THR:HB	2.37	0.60
1:A:192:TRP:HH2	1:A:405:TYR:CZ	2.20	0.59
1:A:162:GLU:H	1:A:162:GLU:CD	2.11	0.58
1:A:324:ASP:HA	2:B:3:THR:HG23	1.85	0.58
1:A:324:ASP:CA	2:B:3:THR:CG2	2.82	0.58
1:A:381:TRP:CG	2:B:3:THR:CB	2.85	0.57
1:A:159:PRO:HB3	1:A:170:TYR:CZ	2.39	0.57
1:A:402:SER:OG	2:B:4:M3L:CM2	2.52	0.57
1:A:494:ALA:HB3	2:B:4:M3L:HE3	1.85	0.57
1:A:201:LYS:HB3	1:A:201:LYS:NZ	2.19	0.57
1:A:192:TRP:CH2	1:A:405:TYR:CZ	2.94	0.56
1:A:350:ALA:CA	1:A:351:GLN:HB2	2.36	0.56
1:A:423:VAL:O	1:A:427:SER:HB3	2.06	0.55
2:B:4:M3L:HM13	2:B:4:M3L:CB	2.35	0.55
1:A:350:ALA:N	1:A:352:SER:N	2.55	0.55
1:A:496:ASN:OD1	2:B:4:M3L:HM31	2.08	0.54
1:A:326:GLU:HB3	1:A:381:TRP:CZ3	2.44	0.53
1:A:211:THR:HB	1:A:320:ILE:HG23	1.91	0.52
1:A:275:PHE:CE1	1:A:320:ILE:HG22	2.44	0.52
1:A:381:TRP:CD1	2:B:3:THR:HB	2.46	0.50
1:A:284:LYS:HB2	1:A:284:LYS:NZ	2.27	0.50
1:A:324:ASP:O	2:B:3:THR:HG23	2.10	0.50
1:A:430:ARG:NH1	5:A:680:HOH:O	2.44	0.50
1:A:324:ASP:HA	2:B:3:THR:CB	2.42	0.49
1:A:163:GLU:HB3	1:A:170:TYR:CD1	2.48	0.49
1:A:183:ILE:HG12	1:A:184:CYS:N	2.28	0.48
1:A:496:ASN:N	2:B:4:M3L:HM11	2.28	0.48
1:A:300:GLU:HG3	5:A:672:HOH:O	2.14	0.48
1:A:324:ASP:C	2:B:3:THR:HG23	2.39	0.48
1:A:192:TRP:CH2	1:A:405:TYR:CE1	3.02	0.47
1:A:316:GLU:C	1:A:317:ILE:HD12	2.39	0.47
1:A:159:PRO:HB3	1:A:170:TYR:CE1	2.50	0.47
1:A:408:TRP:C	1:A:408:TRP:CD1	2.92	0.47
1:A:160:THR:OG1	1:A:162:GLU:HG2	2.15	0.47
1:A:319:VAL:HG22	1:A:386:MET:HE2	1.98	0.46
1:A:350:ALA:N	1:A:351:GLN:CB	2.74	0.45
1:A:495:VAL:HA	2:B:4:M3L:HM12	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:M3L:HG2	2:B:4:M3L:HM32	1.98	0.45
1:A:402:SER:HG	2:B:4:M3L:HM22	1.82	0.45
1:A:175:ARG:HB3	1:A:176:PRO:HD3	1.99	0.44
1:A:404:ASN:O	1:A:473:PHE:HA	2.18	0.43
1:A:281:ASP:O	1:A:285:GLN:HG3	2.17	0.43
1:A:495:VAL:C	2:B:4:M3L:HM11	2.43	0.43
1:A:432:HIS:HE1	1:A:459:GLU:OE2	2.02	0.42
1:A:379:VAL:HG22	1:A:380:PRO:CD	2.48	0.42
1:A:282:PHE:CD2	1:A:282:PHE:C	2.98	0.41
1:A:296:VAL:N	5:A:698:HOH:O	2.53	0.41
1:A:324:ASP:C	2:B:3:THR:CG2	2.93	0.41
1:A:350:ALA:HB3	1:A:351:GLN:CG	2.51	0.41
1:A:324:ASP:OD1	2:B:3:THR:OG1	2.35	0.41
1:A:324:ASP:HA	2:B:3:THR:HG21	2.00	0.41
1:A:327:THR:HA	1:A:328:GLY:HA2	1.61	0.41
1:A:359:LEU:N	1:A:360:PRO:HD2	2.35	0.40
1:A:312:VAL:O	1:A:312:VAL:HG12	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/360 (72%)	253 (97%)	7 (3%)	1 (0%)	30	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	361	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/313 (76%)	217 (91%)	22 (9%)	8	9
2	B	2/2 (100%)	2 (100%)	0	100	100
All	All	241/315 (76%)	219 (91%)	22 (9%)	9	9

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	LYS
1	A	161	GLU
1	A	162	GLU
1	A	168	LEU
1	A	169	LYS
1	A	183	ILE
1	A	191	SER
1	A	277	LYS
1	A	284	LYS
1	A	310	VAL
1	A	319	VAL
1	A	320	ILE
1	A	361	ARG
1	A	362	LEU
1	A	399	HIS
1	A	413	LEU
1	A	425	LEU
1	A	427	SER
1	A	435	GLU
1	A	448	VAL
1	A	455	LEU
1	A	497	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	432	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	M3L	B	4	2	10,11,12	1.04	1 (10%)	9,14,16	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M3L	B	4	2	-	8/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	M3L	CM1-NZ	-2.09	1.43	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	M3L	O-C-CA-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	4	M3L	CG-CD-CE-NZ
2	B	4	M3L	CA-CB-CG-CD
2	B	4	M3L	CE-CD-CG-CB
2	B	4	M3L	CD-CE-NZ-CM1
2	B	4	M3L	CD-CE-NZ-CM3
2	B	4	M3L	CD-CE-NZ-CM2
2	B	4	M3L	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	4	M3L	17	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	OGA	A	502	3	9,9,9	3.20	4 (44%)	8,11,11	1.95	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OGA	A	502	3	-	2/8/9/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	OGA	O1-C1	6.09	1.37	1.22
4	A	502	OGA	C2-N1	5.24	1.42	1.33
4	A	502	OGA	C2-C1	-3.62	1.50	1.54
4	A	502	OGA	O2-C1	2.93	1.38	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	OGA	O2-C1-O1	-4.09	114.17	123.90
4	A	502	OGA	O3-C5-C4	2.28	121.46	112.81

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	502	OGA	O1-C1-C2-N1
4	A	502	OGA	O1-C1-C2-O2'

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/360 (75%)	0.09	13 (4%) 35 41	21, 35, 70, 100	0
2	B	2/3 (66%)	1.75	0 100 100	72, 72, 72, 79	0
All	All	275/363 (75%)	0.10	13 (4%) 36 42	21, 35, 72, 100	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	ALA	4.6
1	A	324	ASP	3.6
1	A	408	TRP	3.4
1	A	328	GLY	3.4
1	A	262	PHE	3.3
1	A	381	TRP	2.9
1	A	498	ALA	2.8
1	A	194	PRO	2.4
1	A	161	GLU	2.3
1	A	444	LEU	2.2
1	A	350	ALA	2.1
1	A	351	GLN	2.0
1	A	315	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	M3L	B	4	12/13	0.67	0.19	44,62,85,90	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	OGA	A	502	10/10	0.97	0.06	26,29,31,34	0
3	FE	A	501	1/1	1.00	0.03	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.