



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:33 PM UTC

PDB ID : 4IH7 / pdb_00004ih7
Title : Hepatitis C Virus polymerase NS5B (BK) with fragment-based compounds
Authors : Harris, S.F.; Wong, A.
Deposited on : 2012-12-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	
1	B	570	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	6	0
			4382	2764	774	809	35			
1	B	557	Total	C	N	O	S	0	7	0
			4390	2768	775	812	35			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	expression tag	UNP P26663
B	1	HIS	-	expression tag	UNP P26663

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 3-(3-tert-butylphenyl)pyridin-2(1H)-one (CCD ID: 1ER) (formula: C₁₅H₁₇NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	15	1	1		
3	B	1	Total	C	N	O	0	0
			17	15	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	191	Total	O	0	0
			191	191		
4	B	181	Total	O	0	0
			181	181		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase



• Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.25Å 104.92Å 125.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.89 – 2.30 39.89 – 2.30	Depositor EDS
% Data completeness (in resolution range)	83.4 (39.89-2.30) 83.4 (39.89-2.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.29Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.11.4	Depositor
R, R_{free}	0.223 , 0.264 0.236 , 0.276	Depositor DCC
R_{free} test set	2164 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.913	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9180	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.26 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2583e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 1ER, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	1/4477 (0.0%)	1.33	10/6074 (0.2%)
1	B	0.78	1/4485 (0.0%)	1.32	14/6085 (0.2%)
All	All	0.78	2/8962 (0.0%)	1.33	24/12159 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	187	MET	SD-CE	-9.98	1.54	1.79
1	B	187	MET	SD-CE	-9.74	1.55	1.79

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ASP	CA-CB-CG	6.10	118.70	112.60
1	B	129	ASP	CA-CB-CG	6.08	118.69	112.60
1	B	552	VAL	N-CA-C	-5.96	106.92	111.62
1	B	303	CYS	N-CA-C	-5.50	105.18	111.07
1	B	170	CYS	CA-C-N	5.48	127.56	120.44
1	B	170	CYS	C-N-CA	5.48	127.56	120.44
1	B	86	GLU	CA-C-N	5.46	127.60	120.28
1	B	86	GLU	C-N-CA	5.46	127.60	120.28
1	B	418	THR	CA-C-N	5.38	127.43	120.44
1	B	418	THR	C-N-CA	5.38	127.43	120.44
1	A	418	THR	CA-C-N	5.37	127.42	120.44
1	A	418	THR	C-N-CA	5.37	127.42	120.44
1	A	197	PRO	CA-C-N	5.36	125.79	119.94
1	A	197	PRO	C-N-CA	5.36	125.79	119.94
1	A	427	THR	CA-C-N	5.34	127.38	120.44
1	A	427	THR	C-N-CA	5.34	127.38	120.44
1	A	201	VAL	N-CA-CB	5.20	117.62	110.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	196	SER	N-CA-C	-5.17	102.52	110.07
1	B	85	VAL	CA-C-N	5.16	127.97	120.79
1	B	85	VAL	C-N-CA	5.16	127.97	120.79
1	A	303	CYS	N-CA-C	-5.10	105.61	111.07
1	B	427	THR	CA-C-N	5.05	127.36	120.54
1	B	427	THR	C-N-CA	5.05	127.36	120.54
1	A	321	VAL	N-CA-CB	5.03	118.35	111.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4382	0	4397	20	0
1	B	4390	0	4400	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	17	0	17	0	0
3	B	17	0	17	0	0
4	A	191	0	0	1	0
4	B	181	0	0	2	0
All	All	9180	0	8831	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:477:TYR:HB3	4:B:879:HOH:O	1.70	0.89
1:A:314:LEU:HB3	1:A:321:VAL:HG23	1.76	0.67
1:B:314:LEU:HB3	1:B:321:VAL:HG23	1.78	0.65
1:A:433:LEU:HB3	1:A:439:LEU:HD12	1.80	0.64
1:B:433:LEU:HB3	1:B:439:LEU:HD12	1.80	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:481:GLU:HB3	4:B:879:HOH:O	2.01	0.60
1:B:48:ARG:HG2	1:B:159:LEU:HD13	1.85	0.59
1:A:201:VAL:HG13	1:A:384:LEU:HG	1.84	0.58
1:B:201:VAL:HG13	1:B:384:LEU:HG	1.84	0.57
1:B:545:LEU:HB3	1:B:547:LEU:HD13	1.86	0.57
1:B:315:VAL:HG22	1:B:320:LEU:HD12	1.87	0.56
1:B:308:LEU:HB3	1:B:311:CYS:SG	2.49	0.52
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.49	0.51
1:A:215:MET:HB2	1:A:326:SER:HB2	1.94	0.50
1:B:461:GLN:HG2	1:B:541:ALA:HB3	1.93	0.50
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.92	0.50
1:B:187:MET:HE2	1:B:296:TYR:CB	2.42	0.50
1:B:215:MET:HB2	1:B:326:SER:HB2	1.94	0.49
1:A:187:MET:HE2	1:A:296:TYR:CB	2.43	0.49
1:A:461:GLN:HG2	1:A:541:ALA:HB3	1.93	0.49
1:A:465:ARG:HG3	1:A:545:LEU:HD12	1.95	0.49
1:B:465:ARG:HD2	1:B:542:ALA:O	2.14	0.48
1:B:83:LEU:HB2	1:B:173[A]:MET:HA	1.95	0.48
1:B:466:LEU:HD22	1:B:551:PHE:HE2	1.78	0.48
1:B:172:LYS:HE3	1:B:560:ILE:HD13	1.94	0.47
1:A:187:MET:CE	1:A:293:LEU:HA	2.45	0.46
1:B:187:MET:CE	1:B:293:LEU:HA	2.45	0.46
1:B:466:LEU:HD22	1:B:551:PHE:CE2	2.51	0.45
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.98	0.45
1:A:531:LYS:HG3	4:A:854:HOH:O	2.17	0.45
1:B:67:VAL:HG11	1:B:300:SER:HB2	2.00	0.44
1:A:172:LYS:HE3	1:A:560:ILE:HD13	1.99	0.44
1:B:187:MET:HG2	1:B:296:TYR:CD1	2.53	0.43
1:A:187:MET:HE2	1:A:296:TYR:HB3	2.01	0.42
1:B:187:MET:HE2	1:B:296:TYR:HB3	2.00	0.42
1:B:5:THR:HG23	1:B:278:ARG:HH12	1.84	0.42
1:B:359:ASP:HB3	1:B:362:LEU:HD12	2.01	0.42
1:A:67:VAL:HG11	1:A:300:SER:HB2	2.02	0.42
1:B:527:ASN:HD22	1:B:536:LEU:HD11	1.84	0.42
1:A:86:GLU:HA	1:A:111:LEU:HD21	2.02	0.42
1:A:5:THR:HG23	1:A:278:ARG:HH12	1.84	0.42
1:A:187:MET:HG2	1:A:296:TYR:CD1	2.55	0.41
1:A:359:ASP:HB3	1:A:362:LEU:HD12	2.02	0.41
1:A:42:SER:HA	1:A:140:ALA:CB	2.50	0.41
1:A:144:VAL:HB	1:A:394:ARG:HG2	2.02	0.41
1:B:176:TYR:OH	1:B:562:HIS:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/570 (98%)	541 (97%)	18 (3%)	0	100	100
1	B	560/570 (98%)	546 (98%)	14 (2%)	0	100	100
All	All	1119/1140 (98%)	1087 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	481/485 (99%)	459 (95%)	22 (5%)	24	36
1	B	482/485 (99%)	462 (96%)	20 (4%)	27	41
All	All	963/970 (99%)	921 (96%)	42 (4%)	25	38

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	CYS
1	A	26	LEU
1	A	31	LEU
1	A	57	LEU
1	A	159	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	VAL
1	A	262	ILE
1	A	303	CYS
1	A	308	LEU
1	A	310	ASP
1	A	315	VAL
1	A	321	VAL
1	A	330	GLN
1	A	381	VAL
1	A	419	LEU
1	A	453	SER
1	A	461	GLN
1	A	466	LEU
1	A	473	SER
1	A	531	LYS
1	A	547	LEU
1	A	556	SER
1	B	14	CYS
1	B	26	LEU
1	B	31	LEU
1	B	32	ARG
1	B	57	LEU
1	B	141	LYS
1	B	159	LEU
1	B	201	VAL
1	B	262	ILE
1	B	303	CYS
1	B	308	LEU
1	B	310	ASP
1	B	315	VAL
1	B	321	VAL
1	B	419	LEU
1	B	453	SER
1	B	461	GLN
1	B	473	SER
1	B	534	LEU
1	B	556	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	241	GLN
1	A	273	ASN
1	A	402	HIS
1	A	461	GLN
1	A	514	GLN
1	A	562	HIS
1	B	241	GLN
1	B	251	GLN
1	B	273	ASN
1	B	514	GLN
1	B	562	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1ER	B	602	-	18,18,18	0.48	0	26,26,26	0.45	0
3	1ER	A	602	-	18,18,18	0.47	0	26,26,26	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1ER	B	602	-	-	0/10/10/10	0/2/2/2
3	1ER	A	602	-	-	6/10/10/10	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	1ER	C-C1-C4-C14
3	A	602	1ER	C-C1-C4-C5
3	A	602	1ER	C3-C1-C4-C14
3	A	602	1ER	C3-C1-C4-C5
3	A	602	1ER	C2-C1-C4-C14
3	A	602	1ER	C2-C1-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	557/570 (97%)	0.41	15 (2%) 56 58	18, 37, 59, 88	6 (1%)
1	B	557/570 (97%)	0.49	15 (2%) 56 58	17, 39, 59, 83	7 (1%)
All	All	1114/1140 (97%)	0.45	30 (2%) 56 58	17, 38, 59, 88	13 (1%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	148	GLN	3.9
1	B	148	GLN	3.1
1	A	548	SER	2.8
1	A	311	CYS	2.7
1	A	111	LEU	2.6
1	A	376	ALA	2.6
1	B	541	ALA	2.5
1	A	102	GLY	2.5
1	B	102	GLY	2.5
1	B	14	CYS	2.4
1	A	106	LYS	2.3
1	A	542	ALA	2.3
1	B	270	LYS	2.3
1	B	531	LYS	2.3
1	A	116	VAL	2.3
1	A	158	ARG	2.2
1	B	382	TYR	2.2
1	A	545	LEU	2.2
1	B	222	ARG	2.2
1	B	562	HIS	2.2
1	B	158	ARG	2.1
1	A	531	LYS	2.1
1	B	303	CYS	2.1
1	B	16	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	402	HIS	2.1
1	A	381	VAL	2.1
1	A	550	TRP	2.1
1	B	542	ALA	2.0
1	B	32	ARG	2.0
1	A	95	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	B	601	1/1	0.90	0.08	77,77,77,77	0
2	ZN	A	601	1/1	0.91	0.12	89,89,89,89	0
3	1ER	A	602	17/17	0.92	0.09	24,30,35,37	0
3	1ER	B	602	17/17	0.94	0.08	29,31,33,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.