



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:12 PM UTC

PDB ID : 9IIE / pdb_00009iie
Title : Cryogenic Temperature Crystal Structure of Fc Fragment of Human IgG1 from Biosimilar VEGF-Trap
Authors : Destan, E.; DeMirici, H.
Deposited on : 2024-06-20
Resolution : 3.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2351 (3.18-3.10)
Clashscore	190562	2452 (3.18-3.10)
Ramachandran outliers	187476	2324 (3.18-3.10)
Sidechain outliers	187428	2324 (3.18-3.10)
RSRZ outliers	180081	2351 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	21% 72% 25% .
1	B	208	6% 82% 15% .
2	C	9	33% 67%
3	D	7	43% 43% 14%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunoglobulin gamma-1 heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	0
			1660	1057	279	318	6			
1	B	207	Total	C	N	O	S	0	0	0
			1658	1056	279	317	6			

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	0	0	0
			110	62	4	44			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	7	Total	C	N	O	0	0	0
			85	48	3	34			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	15	Total 15	O 15	0	0
4	B	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Immunoglobulin gamma-1 heavy chain




- Molecule 1: Immunoglobulin gamma-1 heavy chain



- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.92Å 80.48Å 137.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.84 – 3.14 23.84 – 3.14	Depositor EDS
% Data completeness (in resolution range)	79.5 (23.84-3.14) 79.3 (23.84-3.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 3.17Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.257 , 0.309 0.257 , 0.307	Depositor DCC
R_{free} test set	806 reflections (7.94%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.465	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.80	EDS
Total number of atoms	3543	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUL, MAN, BMA, FUC, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/1706	0.28	0/2324
1	B	0.08	0/1704	0.24	0/2322
All	All	0.11	0/3410	0.26	0/4646

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1660	0	1628	38	0
1	B	1658	0	1625	20	0
2	C	110	0	94	5	0
3	D	85	0	73	2	0
4	A	15	0	0	1	0
4	B	15	0	0	0	0
All	All	3543	0	3420	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:GLN:HE21	1:A:286:ARG:NH2	1.79	0.81
1:A:286:ARG:NH1	2:C:2:NAG:H81	1.97	0.79
1:B:331:PRO:HB3	1:B:357:PHE:HB3	1.63	0.78
1:A:280:GLN:NE2	1:A:286:ARG:NH2	2.34	0.75
1:A:279:GLU:HA	1:A:285:TYR:HD1	1.55	0.71
1:B:261:ASN:HB2	1:B:307:LYS:HB3	1.72	0.71
1:B:303:GLU:HA	1:B:322:SER:HB3	1.75	0.68
1:A:331:PRO:HB3	1:A:357:PHE:HB3	1.76	0.66
1:A:230:PRO:HD3	1:A:244:VAL:HG22	1.78	0.64
1:A:227:LEU:HD13	1:A:319:LYS:HB3	1.82	0.62
1:A:332:GLN:OE1	1:B:345:LYS:NZ	2.25	0.61
1:A:273:LYS:N	1:A:290:VAL:O	2.31	0.61
1:A:227:LEU:HD11	1:A:306:CYS:HB3	1.83	0.59
1:A:350:LEU:HD12	1:A:395:LEU:HD23	1.85	0.59
1:A:286:ARG:NH1	2:C:2:NAG:C8	2.64	0.58
1:A:261:ASN:ND2	1:A:268:GLU:OE2	2.35	0.58
1:A:336:LEU:HB2	1:A:351:THR:HB	1.86	0.57
1:A:355:LYS:NZ	1:B:349:SER:OG	2.38	0.55
1:B:398:ASP:H	1:B:401:ARG:HH12	1.55	0.55
1:B:335:THR:HB	1:B:426:LEU:HD22	1.89	0.54
1:A:296:GLN:NE2	1:A:300:ASN:OD1	2.41	0.54
1:B:398:ASP:H	1:B:401:ARG:NH1	2.07	0.52
1:B:399:LYS:O	1:B:403:GLN:HG2	2.09	0.51
1:A:223:PRO:HB2	1:A:313:LEU:HD21	1.92	0.51
1:A:362:ILE:HG13	1:A:414:HIS:HD2	1.77	0.50
1:A:252:SER:HB2	1:A:255:ASP:HB2	1.94	0.50
1:A:398:ASP:N	1:A:398:ASP:OD1	2.42	0.49
1:A:228:PHE:CE1	2:C:3:BMA:H62	2.49	0.48
1:A:243:GLU:HG3	1:A:290:VAL:HG13	1.97	0.46
1:A:353:LEU:HD13	1:A:392:TYR:CZ	2.49	0.46
1:B:364:VAL:HG22	1:B:412:VAL:HG22	1.97	0.46
3:D:4:MAN:H61	3:D:5:NAG:H82	1.97	0.45
1:A:253:HIS:NE2	1:A:283:SER:O	2.50	0.45
1:B:362:ILE:HG21	1:B:391:LEU:HD21	1.98	0.44
1:A:249:VAL:HG11	2:C:2:NAG:H2	2.00	0.44
1:A:341:ASP:N	1:A:341:ASP:OD1	2.47	0.44
1:B:346:ASN:N	1:B:346:ASN:OD1	2.51	0.44
1:A:340:ARG:HA	1:A:343:LEU:HD23	2.00	0.44
1:A:362:ILE:HG12	1:A:363:ALA:H	1.81	0.43
1:B:263:TYR:HB2	1:B:305:LYS:HB3	2.00	0.43
1:A:424:LYS:NZ	4:A:502:HOH:O	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:384:ASP:OD1	1:B:385:SER:N	2.45	0.43
1:B:348:VAL:HG23	1:B:399:LYS:HG2	1.99	0.43
1:A:235:THR:HG21	1:A:298:TRP:CD1	2.54	0.42
1:A:228:PHE:HE1	2:C:3:BMA:H62	1.83	0.42
1:B:264:VAL:HG22	1:B:304:TYR:HD1	1.85	0.42
1:A:279:GLU:HA	1:A:285:TYR:CD1	2.45	0.42
1:A:282:ASN:OD1	1:A:284:THR:OG1	2.31	0.42
1:B:345:LYS:HB3	1:B:345:LYS:HE2	1.83	0.42
1:A:280:GLN:NE2	1:A:286:ARG:HH22	2.16	0.41
1:B:248:VAL:HB	1:B:287:VAL:HG23	2.03	0.41
1:B:352:CYS:HB2	1:B:366:TRP:CZ2	2.55	0.41
1:A:367:GLU:HG2	1:A:372:PRO:HA	2.03	0.41
1:A:303:GLU:HG2	1:A:322:SER:HB3	2.01	0.41
1:A:253:HIS:ND1	1:A:253:HIS:N	2.68	0.41
1:B:228:PHE:HE1	3:D:3:BMA:H62	1.85	0.41
1:A:417:LEU:HD13	1:A:422:THR:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/208 (99%)	195 (95%)	10 (5%)	0	100	100
1	B	205/208 (99%)	202 (98%)	3 (2%)	0	100	100
All	All	410/416 (99%)	397 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/194 (100%)	181 (93%)	13 (7%)	15	39
1	B	193/194 (100%)	182 (94%)	11 (6%)	18	45
All	All	387/388 (100%)	363 (94%)	24 (6%)	16	42

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	238	ILE
1	A	253	HIS
1	A	261	ASN
1	A	286	ARG
1	A	290	VAL
1	A	317	ILE
1	A	341	ASP
1	A	346	ASN
1	A	362	ILE
1	A	369	ASN
1	A	398	ASP
1	A	404	GLN
1	A	428	LEU
1	B	237	MET
1	B	241	THR
1	B	254	GLU
1	B	273	LYS
1	B	277	ARG
1	B	287	VAL
1	B	345	LYS
1	B	346	ASN
1	B	399	LYS
1	B	401	ARG
1	B	423	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	280	GLN
1	A	371	GLN
1	A	374	ASN
1	A	420	HIS
1	B	419	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.23	0	17,19,21	0.52	0
2	NAG	C	2	2	14,14,15	0.29	0	17,19,21	0.61	0
2	BMA	C	3	2	11,11,12	0.59	0	15,15,17	0.95	0
2	MAN	C	4	2	11,11,12	1.29	1 (9%)	15,15,17	1.26	2 (13%)
2	NAG	C	5	2	14,14,15	0.88	1 (7%)	17,19,21	1.36	1 (5%)
2	GAL	C	6	2	11,11,12	0.62	0	15,15,17	0.92	0
2	MAN	C	7	2	11,11,12	0.83	0	15,15,17	1.41	2 (13%)
2	NAG	C	8	2	14,14,15	0.25	0	17,19,21	0.35	0
2	FUL	C	9	2	10,10,11	1.79	3 (30%)	14,14,16	1.04	1 (7%)
3	NAG	D	1	3,1	14,14,15	0.35	0	17,19,21	0.38	0
3	NAG	D	2	3	14,14,15	0.31	0	17,19,21	0.42	0
3	BMA	D	3	3	11,11,12	0.69	0	15,15,17	0.82	0
3	MAN	D	4	3	11,11,12	0.66	0	15,15,17	1.08	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	5	3	14,14,15	0.26	0	17,19,21	0.43	0
3	MAN	D	6	3	11,11,12	0.67	0	15,15,17	0.92	2 (13%)
3	FUC	D	7	3	10,10,11	0.69	0	14,14,16	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
2	BMA	C	3	2	-	2/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	4/6/23/26	0/1/1/1
2	GAL	C	6	2	-	1/2/19/22	0/1/1/1
2	MAN	C	7	2	-	0/2/19/22	1/1/1/1
2	NAG	C	8	2	-	1/6/23/26	0/1/1/1
2	FUL	C	9	2	-	-	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	NAG	D	5	3	-	2/6/23/26	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	FUC	D	7	3	-	-	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	9	FUL	O5-C1	4.37	1.51	1.43
2	C	5	NAG	O5-C1	3.08	1.48	1.43
2	C	9	FUL	C2-C3	-2.36	1.48	1.52
2	C	4	MAN	C2-C3	2.19	1.55	1.52
2	C	9	FUL	O5-C5	2.09	1.47	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	5	NAG	C1-O5-C5	5.19	119.14	112.19
2	C	7	MAN	C1-O5-C5	3.62	117.04	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7	MAN	C1-C2-C3	-2.79	105.58	109.64
3	D	4	MAN	C1-O5-C5	2.58	115.64	112.19
3	D	4	MAN	O2-C2-C3	-2.54	104.89	110.15
2	C	4	MAN	C1-O5-C5	2.44	115.46	112.19
2	C	4	MAN	O2-C2-C1	2.40	114.72	109.22
3	D	6	MAN	C1-O5-C5	2.21	115.15	112.19
2	C	9	FUL	C6-C5-C4	-2.15	109.15	113.08
3	D	6	MAN	O2-C2-C3	-2.06	105.89	110.15

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	BMA	O5-C5-C6-O6
3	D	5	NAG	O5-C5-C6-O6
2	C	5	NAG	C4-C5-C6-O6
2	C	5	NAG	O5-C5-C6-O6
2	C	3	BMA	C4-C5-C6-O6
3	D	5	NAG	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	C	6	GAL	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	C	8	NAG	O5-C5-C6-O6
2	C	1	NAG	C1-C2-N2-C7
2	C	5	NAG	C3-C2-N2-C7
2	C	5	NAG	C1-C2-N2-C7
2	C	1	NAG	C3-C2-N2-C7


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	7	MAN	C1-C2-C3-C4-C5-O5

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	3	BMA	1	0
2	C	3	BMA	2	0
2	C	2	NAG	3	0
3	D	5	NAG	1	0
3	D	4	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/208 (99%)	1.19	43 (20%) 2 1	16, 41, 110, 144	0
1	B	207/208 (99%)	0.72	12 (5%) 29 15	15, 31, 55, 76	0
All	All	414/416 (99%)	0.95	55 (13%) 7 3	15, 34, 102, 144	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	VAL	4.6
1	A	252	SER	4.5
1	A	270	HIS	4.2
1	A	276	PRO	4.1
1	A	281	TYR	3.9
1	A	309	SER	3.7
1	B	403	GLN	3.6
1	A	306	CYS	3.6
1	A	287	VAL	3.5
1	A	279	GLU	3.4
1	A	285	TYR	3.4
1	A	260	PHE	3.4
1	A	258	VAL	3.3
1	A	310	ASN	3.3
1	A	289	SER	3.2
1	A	268	GLU	3.2
1	A	254	GLU	3.1
1	A	249	VAL	3.1
1	A	272	ALA	2.9
1	A	269	VAL	2.8
1	A	312	ALA	2.8
1	A	253	HIS	2.7
1	A	280	GLN	2.6
1	A	313	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	244	VAL	2.6
1	A	256	PRO	2.6
1	A	257	GLU	2.4
1	A	291	LEU	2.4
1	A	278	GLU	2.4
1	A	286	ARG	2.3
1	A	263	TYR	2.3
1	A	247	VAL	2.3
1	B	301	GLY	2.2
1	B	324	ALA	2.2
1	B	414	HIS	2.2
1	A	267	VAL	2.2
1	B	348	VAL	2.2
1	B	252	SER	2.2
1	A	311	LYS	2.2
1	B	404	GLN	2.1
1	A	290	VAL	2.1
1	A	283	SER	2.1
1	A	246	CYS	2.1
1	B	405	GLY	2.1
1	A	264	VAL	2.1
1	B	251	VAL	2.1
1	A	271	ASN	2.1
1	A	275	LYS	2.1
1	A	250	ASP	2.1
1	B	225	VAL	2.0
1	A	315	ALA	2.0
1	B	347	GLN	2.0
1	A	277	ARG	2.0
1	A	284	THR	2.0
1	B	281	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FUL	C	9	10/11	0.35	0.23	92,102,108,108	0
2	NAG	C	1	14/15	0.44	0.26	95,108,113,114	0
2	MAN	C	7	11/12	0.58	0.16	42,54,58,61	0
2	NAG	C	2	14/15	0.60	0.18	51,76,88,89	0
2	MAN	C	4	11/12	0.64	0.15	42,45,47,50	0
2	BMA	C	3	11/12	0.67	0.15	49,57,61,64	0
2	NAG	C	5	14/15	0.72	0.17	38,44,51,55	0
2	NAG	C	8	14/15	0.73	0.15	51,59,66,76	0
3	MAN	D	6	11/12	0.76	0.15	36,39,45,45	0
3	NAG	D	1	14/15	0.77	0.14	27,30,34,34	0
2	GAL	C	6	11/12	0.77	0.14	40,45,48,52	0
3	FUC	D	7	10/11	0.79	0.16	28,31,34,36	0
3	MAN	D	4	11/12	0.82	0.12	33,38,40,40	0
3	NAG	D	5	14/15	0.84	0.15	27,37,41,53	0
3	BMA	D	3	11/12	0.85	0.11	30,36,40,42	0
3	NAG	D	2	14/15	0.86	0.12	24,28,34,34	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**6.4 Ligands** [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.