



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 07:59 PM UTC

PDB ID : 5IJD / pdb_00005ijd
Title : The crystal structure of mouse TLR4/MD-2/lipid A complex
Authors : Wang, Y.; Su, L.; Morin, M.D.; Jones, B.T.; Whitby, L.R.; Surakattula, M.; Huang, H.; Shi, H.; Choi, J.H.; Wang, K.; Moresco, E.M.; Berger, M.; Zhan, X.; Zhang, H.; Boger, D.L.; Beutler, B.
Deposited on : 2016-03-01
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	13% 66% 31% ..
1	B	594	17% 69% 27% .
2	C	150	3% 64% 25% .. 9%
2	D	150	5% 57% 30% . 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	2	 100%
3	F	2	 100%
3	G	2	 50% 50%
3	H	2	 100%
3	I	2	 100%
3	J	2	 100%
3	K	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	E	2	X	-	-	-
3	NAG	G	1	X	-	-	-
3	NAG	H	2	X	-	-	-
3	NAG	J	1	X	-	-	-
3	NAG	J	2	X	-	-	-
3	NAG	K	2	X	-	-	-
4	NAG	B	705	X	-	-	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 12385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 4, Variable lymphocyte receptor B chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	590	Total	C	N	O	S	0	1	0
			4708	3011	784	889	24			
1	B	592	Total	C	N	O	S	0	0	0
			4713	3014	783	891	25			


- Molecule 2 is a protein called Lymphocyte antigen 96.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	137	Total	C	N	O	S	0	0	0
			1112	718	188	199	7			
2	D	137	Total	C	N	O	S	0	0	0
			1112	718	188	199	7			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	161	LYS	-	cloning artifact	UNP Q9JHF9
C	162	GLY	-	cloning artifact	UNP Q9JHF9
C	163	GLU	-	cloning artifact	UNP Q9JHF9
C	164	ASN	-	cloning artifact	UNP Q9JHF9
C	165	LEU	-	cloning artifact	UNP Q9JHF9
C	166	TYR	-	cloning artifact	UNP Q9JHF9
C	167	PHE	-	cloning artifact	UNP Q9JHF9
C	168	GLN	-	cloning artifact	UNP Q9JHF9
D	161	LYS	-	cloning artifact	UNP Q9JHF9
D	162	GLY	-	cloning artifact	UNP Q9JHF9
D	163	GLU	-	cloning artifact	UNP Q9JHF9
D	164	ASN	-	cloning artifact	UNP Q9JHF9
D	165	LEU	-	cloning artifact	UNP Q9JHF9
D	166	TYR	-	cloning artifact	UNP Q9JHF9
D	167	PHE	-	cloning artifact	UNP Q9JHF9
D	168	GLN	-	cloning artifact	UNP Q9JHF9

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	J	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	K	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is 2-deoxy-3-O-[(3R)-3-hydroxytetradecanoyl]-2-([(3R)-3-hydroxytetradecanoyl] amino)-4-O-phosphono-beta-D-glucopyranose (CCD ID: LP4) (formula: C₃₄H₆₆NO₁₂P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	0	0
			45	32	1	11	1		
5	D	1	Total	C	N	O	P	0	0
			45	32	1	11	1		

- Molecule 6 is (R)-((2R,3S,4R,5R,6R)-3-HYDROXY-2-(HYDROXYMETHYL)-5-((R)-3-HYDROXYTETRADECANAMIDO)-6-(PHOSPHONOXY)TETRAHYDRO-2H-PYRAN-4-YL) 3-HYDROXYTETRADECANOATE (CCD ID: LP5) (formula: C₃₄H₆₆NO₁₂P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total 48	C 34	N 1	O 12	P 1	0	0
6	D	1	Total 48	C 34	N 1	O 12	P 1	0	0

- Molecule 7 is LAURIC ACID (CCD ID: DAO) (formula: C₁₂H₂₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			13	12	1		
7	D	1	Total	C	O	0	0
			13	12	1		

- Molecule 8 is MYRISTIC ACID (CCD ID: MYR) (formula: $C_{14}H_{28}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	C	O	0	0
			15	14	1		
8	D	1	Total	C	O	0	0
			15	14	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	92	Total	O	0	0
			92	92		
9	C	29	Total	O	0	0
			29	29		
9	D	22	Total	O	0	0
			22	22		
9	B	89	Total	O	0	0
			89	89		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Toll-like receptor 4, Variable lymphocyte receptor B chimera




- Molecule 1: Toll-like receptor 4, Variable lymphocyte receptor B chimera






- Molecule 2: Lymphocyte antigen 96



- Molecule 2: Lymphocyte antigen 96



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 50%


MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%


MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50% 50%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.84Å 145.74Å 136.19Å 90.00° 100.67° 90.00°	Depositor
Resolution (Å)	39.34 – 2.70 39.34 – 2.70	Depositor EDS
% Data completeness (in resolution range)	77.2 (39.34-2.70) 77.7 (39.34-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.215 , 0.285 0.217 , 0.284	Depositor DCC
R_{free} test set	1978 reflections (2.78%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12385	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LP5, NAG, LP4, MYR, DAO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.92	0/4811	1.04	7/6519 (0.1%)
1	B	0.87	1/4814 (0.0%)	1.07	13/6524 (0.2%)
2	C	0.93	0/1143	1.05	1/1544 (0.1%)
2	D	0.90	0/1143	1.01	4/1544 (0.3%)
All	All	0.90	1/11911 (0.0%)	1.05	25/16131 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
2	C	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	432	LEU	CA-C	-6.37	1.44	1.52

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	31	VAL	N-CA-C	15.08	125.47	111.45
1	B	32	VAL	N-CA-C	-9.71	87.90	108.88
1	B	272	GLU	CA-C-N	6.93	127.22	119.32
1	B	272	GLU	C-N-CA	6.93	127.22	119.32
1	A	249	LEU	N-CA-C	-6.15	105.81	113.38

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	278	GLY	Peptide
1	A	361	GLY	Peptide
1	B	361	GLY	Peptide
2	C	132	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4708	0	4698	145	2
1	B	4713	0	4694	127	2
2	C	1112	0	1075	31	0
2	D	1112	0	1075	37	0
3	E	28	0	25	0	0
3	F	28	0	25	0	0
3	G	28	0	25	3	0
3	H	28	0	25	0	0
3	I	28	0	25	0	0
3	J	28	0	25	0	0
3	K	28	0	25	1	0
4	A	28	0	26	4	0
4	B	14	0	13	1	0
4	C	14	0	13	0	0
4	D	14	0	13	0	0
5	C	45	0	53	0	0
5	D	45	0	53	1	0
6	C	48	0	63	2	0
6	D	48	0	63	1	0
7	C	13	0	23	0	0
7	D	13	0	23	0	0
8	C	15	0	27	1	0
8	D	15	0	27	2	0
9	A	92	0	0	5	0
9	B	89	0	0	5	0
9	C	29	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	D	22	0	0	3	0
All	All	12385	0	12114	342	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 342 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:804:LP4:C1	6:D:805:LP5:O6	1.64	1.44
1:A:117:SER:O	1:A:119:GLY:N	2.04	0.90
1:A:191:VAL:HG12	1:A:221:ALA:HA	1.55	0.85
1:B:261:GLU:OE2	1:B:267:ASN:HB3	1.77	0.83
1:B:255:HIS:ND1	1:B:285:ASP:OD2	2.12	0.83

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:564:ARG:NH1	1:B:542:TYR:CD1[1_655]	2.15	0.05
1:A:540:GLN:O	1:B:519:ARG:NH2[1_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/594 (99%)	516 (88%)	72 (12%)	1 (0%)	43	68
1	B	590/594 (99%)	521 (88%)	64 (11%)	5 (1%)	16	37
2	C	135/150 (90%)	119 (88%)	16 (12%)	0	100	100
2	D	135/150 (90%)	114 (84%)	20 (15%)	1 (1%)	18	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1449/1488 (97%)	1270 (88%)	172 (12%)	7 (0%)	24	48

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	196	PHE
1	B	195	GLN
1	A	118	PRO
1	B	515	ASP
1	B	93	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	548/551 (100%)	522 (95%)	26 (5%)	23	51
1	B	549/551 (100%)	513 (93%)	36 (7%)	15	36
2	C	124/136 (91%)	119 (96%)	5 (4%)	28	56
2	D	124/136 (91%)	112 (90%)	12 (10%)	8	20
All	All	1345/1374 (98%)	1266 (94%)	79 (6%)	18	42

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	209	MET
1	B	380	ARG
1	B	268	LEU
1	B	340	LEU
1	B	423	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	235	ASN
1	B	247	GLN
1	B	359	ASN
2	C	21	GLN
2	C	98	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

19 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	K	1	1,3	14,14,15	0.98	1 (7%)	17,19,21	2.03	4 (23%)
3	NAG	K	2	3	14,14,15	0.73	0	17,19,21	1.51	2 (11%)
3	NAG	E	1	1,3	14,14,15	0.56	0	17,19,21	1.36	3 (17%)
3	NAG	F	1	1,3	14,14,15	0.72	0	17,19,21	1.98	4 (23%)
3	NAG	G	1	2,3	14,14,15	0.71	0	17,19,21	2.22	7 (41%)
3	NAG	E	2	3	14,14,15	0.47	0	17,19,21	1.64	4 (23%)
4	NAG	A	703	1	14,14,15	0.59	0	17,19,21	2.36	7 (41%)
3	NAG	J	1	1,3	14,14,15	0.79	0	17,19,21	2.01	5 (29%)
4	NAG	D	803	2	14,14,15	0.98	0	17,19,21	2.99	7 (41%)
4	NAG	A	704	1	14,14,15	0.74	0	17,19,21	2.09	5 (29%)
3	NAG	H	1	2,3	14,14,15	0.75	0	17,19,21	3.08	7 (41%)
3	NAG	G	2	3	14,14,15	0.55	0	17,19,21	1.91	5 (29%)
3	NAG	J	2	3	14,14,15	0.81	1 (7%)	17,19,21	1.55	3 (17%)
3	NAG	I	1	1,3	14,14,15	0.51	0	17,19,21	2.03	4 (23%)
4	NAG	B	705	1	14,14,15	1.34	3 (21%)	17,19,21	2.43	3 (17%)
3	NAG	I	2	3	14,14,15	0.80	1 (7%)	17,19,21	1.57	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	201	2	14,14,15	0.67	0	17,19,21	1.64	4 (23%)
3	NAG	F	2	3	14,14,15	0.62	0	17,19,21	1.17	2 (11%)
3	NAG	H	2	3	14,14,15	0.90	1 (7%)	17,19,21	1.52	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	K	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	K	2	3	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	1	2,3	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	E	2	3	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	A	703	1	-	3/6/23/26	0/1/1/1
3	NAG	J	1	1,3	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	D	803	2	-	3/6/23/26	0/1/1/1
4	NAG	A	704	1	-	3/6/23/26	0/1/1/1
3	NAG	H	1	2,3	-	3/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	NAG	J	2	3	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
4	NAG	B	705	1	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	I	2	3	-	3/6/23/26	0/1/1/1
4	NAG	C	201	2	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	3/6/23/26	0/1/1/1
3	NAG	H	2	3	1/1/5/7	3/6/23/26	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	705	NAG	C1-C2	3.23	1.56	1.52
3	K	1	NAG	C1-C2	2.96	1.56	1.52
4	B	705	NAG	C3-C2	2.58	1.57	1.52
3	I	2	NAG	C1-C2	2.46	1.55	1.52
3	H	2	NAG	C1-C2	2.35	1.55	1.52

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	1	NAG	C2-N2-C7	8.03	133.67	122.90
4	B	705	NAG	O5-C1-C2	-7.28	100.02	111.29
3	I	1	NAG	C1-O5-C5	6.62	121.05	112.19
4	D	803	NAG	C4-C3-C2	-5.86	102.42	111.02
3	H	1	NAG	C8-C7-N2	5.85	125.81	116.12

5 of 7 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	2	NAG	C1
3	G	1	NAG	C1
3	H	2	NAG	C1
3	J	1	NAG	C1
3	J	2	NAG	C1

5 of 43 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	H	1	NAG	C3-C2-N2-C7
4	D	803	NAG	C3-C2-N2-C7
4	C	201	NAG	C4-C5-C6-O6
4	D	803	NAG	C4-C5-C6-O6
3	J	2	NAG	C4-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	K	1	NAG	1	0
3	G	1	NAG	3	0
4	A	703	NAG	1	0
4	A	704	NAG	3	0
4	B	705	NAG	1	0

5.5 Carbohydrates

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	E	1	1,3	14,14,15	0.56	0	17,19,21	1.36	3 (17%)
3	NAG	E	2	3	14,14,15	0.47	0	17,19,21	1.64	4 (23%)
3	NAG	F	1	1,3	14,14,15	0.72	0	17,19,21	1.98	4 (23%)
3	NAG	F	2	3	14,14,15	0.62	0	17,19,21	1.17	2 (11%)
3	NAG	G	1	2,3	14,14,15	0.71	0	17,19,21	2.22	7 (41%)
3	NAG	G	2	3	14,14,15	0.55	0	17,19,21	1.91	5 (29%)
3	NAG	H	1	2,3	14,14,15	0.75	0	17,19,21	3.08	7 (41%)
3	NAG	H	2	3	14,14,15	0.90	1 (7%)	17,19,21	1.52	2 (11%)
3	NAG	I	1	1,3	14,14,15	0.51	0	17,19,21	2.03	4 (23%)
3	NAG	I	2	3	14,14,15	0.80	1 (7%)	17,19,21	1.57	3 (17%)
3	NAG	J	1	1,3	14,14,15	0.79	0	17,19,21	2.01	5 (29%)
3	NAG	J	2	3	14,14,15	0.81	1 (7%)	17,19,21	1.55	3 (17%)
3	NAG	K	1	1,3	14,14,15	0.98	1 (7%)	17,19,21	2.03	4 (23%)
3	NAG	K	2	3	14,14,15	0.73	0	17,19,21	1.51	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	E	2	3	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	3/6/23/26	0/1/1/1
3	NAG	G	1	2,3	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	NAG	H	1	2,3	-	3/6/23/26	0/1/1/1
3	NAG	H	2	3	1/1/5/7	3/6/23/26	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	3/6/23/26	0/1/1/1
3	NAG	J	1	1,3	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	J	2	3	1/1/5/7	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	K	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	K	2	3	1/1/5/7	3/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	1	NAG	C1-C2	2.96	1.56	1.52
3	I	2	NAG	C1-C2	2.46	1.55	1.52
3	H	2	NAG	C1-C2	2.35	1.55	1.52
3	J	2	NAG	C1-C2	2.02	1.55	1.52

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	1	NAG	C2-N2-C7	8.03	133.67	122.90
3	I	1	NAG	C1-O5-C5	6.62	121.05	112.19
3	H	1	NAG	C8-C7-N2	5.85	125.81	116.12
3	K	1	NAG	C8-C7-N2	4.99	124.39	116.12
3	F	1	NAG	C1-O5-C5	4.98	118.86	112.19

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	2	NAG	C1
3	G	1	NAG	C1
3	H	2	NAG	C1
3	J	1	NAG	C1
3	J	2	NAG	C1

5 of 29 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	H	1	NAG	C3-C2-N2-C7
3	J	2	NAG	C4-C5-C6-O6
3	K	2	NAG	O5-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
3	J	1	NAG	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	3	0
3	K	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	LP4	D	804	-	45,45,48	0.89	2 (4%)	54,56,60	1.64	8 (14%)
4	NAG	B	705	1	14,14,15	1.34	3 (21%)	17,19,21	2.43	3 (17%)
4	NAG	A	703	1	14,14,15	0.59	0	17,19,21	2.36	7 (41%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	D	803	2	14,14,15	0.98	0	17,19,21	2.99	7 (41%)
4	NAG	C	201	2	14,14,15	0.67	0	17,19,21	1.64	4 (23%)
6	LP5	D	805	-	47,48,48	0.76	2 (4%)	58,60,60	1.33	7 (12%)
4	NAG	A	704	1	14,14,15	0.74	0	17,19,21	2.09	5 (29%)
8	MYR	D	807	-	13,14,15	0.35	0	12,13,15	0.92	0
8	MYR	C	207	-	13,14,15	0.35	0	12,13,15	0.93	0
6	LP5	C	205	-	47,48,48	0.77	2 (4%)	58,60,60	1.29	5 (8%)
7	DAO	D	806	-	11,12,13	0.50	0	10,11,13	0.42	0
7	DAO	C	206	-	11,12,13	0.49	0	10,11,13	0.73	0
5	LP4	C	204	-	45,45,48	0.80	2 (4%)	54,56,60	1.66	12 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LP4	D	804	-	-	12/43/60/65	0/1/1/1
4	NAG	A	703	1	-	3/6/23/26	0/1/1/1
4	NAG	B	705	1	1/1/5/7	3/6/23/26	0/1/1/1
4	NAG	D	803	2	-	3/6/23/26	0/1/1/1
4	NAG	C	201	2	-	2/6/23/26	0/1/1/1
6	LP5	D	805	-	-	14/44/65/65	0/1/1/1
4	NAG	A	704	1	-	3/6/23/26	0/1/1/1
8	MYR	D	807	-	-	6/12/12/13	-
8	MYR	C	207	-	-	3/12/12/13	-
6	LP5	C	205	-	-	15/44/65/65	0/1/1/1
7	DAO	D	806	-	-	4/10/10/11	-
7	DAO	C	206	-	-	7/10/10/11	-
5	LP4	C	204	-	-	14/43/60/65	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	804	LP4	O3-C28	3.93	1.45	1.34
5	C	204	LP4	O3-C28	3.62	1.44	1.34
6	C	205	LP5	O3-C28	3.54	1.44	1.34
6	D	805	LP5	O3-C28	3.51	1.44	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	705	NAG	C1-C2	3.23	1.56	1.52

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	705	NAG	O5-C1-C2	-7.28	100.02	111.29
4	D	803	NAG	C4-C3-C2	-5.86	102.42	111.02
4	D	803	NAG	O5-C1-C2	-5.38	102.96	111.29
5	D	804	LP4	C1-C2-N2	-5.38	101.96	110.43
4	A	703	NAG	C1-O5-C5	5.09	119.01	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	705	NAG	C1

5 of 89 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	803	NAG	C3-C2-N2-C7
5	C	204	LP4	C17-C16-C8-C7
5	C	204	LP4	O44-C16-C8-C7
5	C	204	LP4	C29-C30-C31-C32
5	C	204	LP4	O43-C30-C31-C32


There are no ring outliers.


8 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	804	LP4	1	0
4	B	705	NAG	1	0
4	A	703	NAG	1	0
6	D	805	LP5	1	0
4	A	704	NAG	3	0
8	D	807	MYR	2	0
8	C	207	MYR	1	0
6	C	205	LP5	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	590/594 (99%)	0.50	77 (13%) 7 6	9, 33, 95, 142	1 (0%)
1	B	592/594 (99%)	0.75	102 (17%) 4 3	10, 43, 99, 132	0
2	C	137/150 (91%)	0.19	5 (3%) 46 42	20, 35, 55, 76	0
2	D	137/150 (91%)	0.58	7 (5%) 33 29	23, 42, 65, 79	0
All	All	1456/1488 (97%)	0.58	191 (13%) 7 6	9, 37, 95, 142	1 (0%)

The worst 5 of 191 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	121	PHE	7.7
1	B	27	PRO	7.1
1	A	100	LEU	6.8
1	A	80	GLN	6.0
1	A	102	HIS	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	J	2	14/15	0.43	0.21	95,108,117,122	0
3	NAG	I	2	14/15	0.59	0.19	70,82,103,104	0
4	NAG	A	703	14/15	0.62	0.19	52,76,83,84	0
3	NAG	J	1	14/15	0.65	0.28	98,106,115,118	0
4	NAG	B	705	14/15	0.65	0.19	67,75,83,85	0
4	NAG	A	704	14/15	0.71	0.16	66,72,77,78	0
3	NAG	F	2	14/15	0.72	0.14	39,52,57,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	H	2	14/15	0.73	0.14	78,86,88,90	0
4	NAG	D	803	14/15	0.73	0.17	68,74,77,77	0
3	NAG	E	2	14/15	0.76	0.15	64,76,82,86	0
3	NAG	I	1	14/15	0.77	0.18	84,88,91,94	0
3	NAG	G	2	14/15	0.78	0.13	65,74,85,87	0
3	NAG	E	1	14/15	0.79	0.16	59,65,74,75	0
4	NAG	C	201	14/15	0.79	0.14	66,74,83,92	0
3	NAG	K	2	14/15	0.79	0.13	42,48,52,56	0
3	NAG	G	1	14/15	0.83	0.13	45,48,53,62	0
3	NAG	H	1	14/15	0.86	0.11	47,52,59,70	0
3	NAG	K	1	14/15	0.93	0.09	24,27,33,37	0
3	NAG	F	1	14/15	0.96	0.07	26,30,33,38	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	J	2	14/15	0.43	0.21	95,108,117,122	0
3	NAG	I	2	14/15	0.59	0.19	70,82,103,104	0
3	NAG	J	1	14/15	0.65	0.28	98,106,115,118	0
3	NAG	F	2	14/15	0.72	0.14	39,52,57,60	0
3	NAG	H	2	14/15	0.73	0.14	78,86,88,90	0
3	NAG	E	2	14/15	0.76	0.15	64,76,82,86	0
3	NAG	I	1	14/15	0.77	0.18	84,88,91,94	0
3	NAG	G	2	14/15	0.78	0.13	65,74,85,87	0
3	NAG	E	1	14/15	0.79	0.16	59,65,74,75	0
3	NAG	K	2	14/15	0.79	0.13	42,48,52,56	0
3	NAG	G	1	14/15	0.83	0.13	45,48,53,62	0
3	NAG	H	1	14/15	0.86	0.11	47,52,59,70	0
3	NAG	K	1	14/15	0.93	0.09	24,27,33,37	0
3	NAG	F	1	14/15	0.96	0.07	26,30,33,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	703	14/15	0.62	0.19	52,76,83,84	0
4	NAG	B	705	14/15	0.65	0.19	67,75,83,85	0
4	NAG	A	704	14/15	0.71	0.16	66,72,77,78	0
4	NAG	D	803	14/15	0.73	0.17	68,74,77,77	0
4	NAG	C	201	14/15	0.79	0.14	66,74,83,92	0
8	MYR	C	207	15/16	0.83	0.25	52,62,77,89	0
8	MYR	D	807	15/16	0.83	0.20	46,50,65,66	0
5	LP4	D	804	45/48	0.88	0.12	35,50,59,68	0
7	DAO	C	206	13/14	0.89	0.15	39,44,59,65	0
5	LP4	C	204	45/48	0.90	0.12	28,37,52,56	0
7	DAO	D	806	13/14	0.92	0.13	45,51,63,69	0
6	LP5	D	805	48/48	0.92	0.11	31,45,63,67	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	LP5	C	205	48/48	0.92	0.11	38,46,56,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LP4 D 804:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LP4 C 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LP5 D 805:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LP5 C 205:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.