



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:53 AM UTC

PDB ID : 9IJU / pdb_00009iju
Title : Sertraline enhances the deubiquitinase activity of USP7 by binding to its switching loop region
Authors : Shi, L.; Xu, Z.; Chen, X.; Xiong, B.; Zhang, N.
Deposited on : 2024-06-25
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	4% 86% 6% 8%
1	C	353	3% 83% 8% 8%
1	E	353	2% 85% 6% 8%
1	G	353	6% 78% 6% 16%
2	B	75	% 92% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	75	99%
2	F	75	91% 9%
2	H	75	% 97%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2511	1596	414	485	16			
1	C	325	Total	C	N	O	S	0	0	0
			2454	1553	410	476	15			
1	E	323	Total	C	N	O	S	0	0	0
			2427	1534	403	474	16			
1	G	298	Total	C	N	O	S	0	0	0
			2118	1325	359	421	13			

- Molecule 2 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	75	Total	C	N	O	S	0	0	0
			587	370	100	116	1			
2	D	75	Total	C	N	O	S	0	0	0
			584	368	99	116	1			
2	F	75	Total	C	N	O	S	0	0	0
			590	371	102	116	1			
2	H	75	Total	C	N	O	S	0	0	0
			581	365	100	115	1			

- Molecule 3 is (1S,4S)-4-(3,4-dichlorophenyl)-N-methyl-1,2,3,4-tetrahydronaphthalen-1-amine (CCD ID: SRE) (formula: C₁₇H₁₇Cl₂N) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	Cl	N	0	0
			20	17	2	1		
3	G	1	Total	C	Cl	N	0	0
			20	17	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total	O	0	0
			12	12		
4	B	9	Total	O	0	0
			9	9		
4	C	12	Total	O	0	0
			12	12		
4	D	2	Total	O	0	0
			2	2		
4	E	10	Total	O	0	0
			10	10		
4	F	5	Total	O	0	0
			5	5		
4	G	8	Total	O	0	0
			8	8		
4	H	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7




- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7




- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7





- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 7






● Molecule 2: Ubiquitin



● Molecule 2: Ubiquitin



● Molecule 2: Ubiquitin



● Molecule 2: Ubiquitin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.64Å 84.86Å 106.78Å 90.00° 90.21° 90.00°	Depositor
Resolution (Å)	20.58 – 2.46 20.58 – 2.46	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.58-2.46) 99.6 (20.58-2.46)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.47Å)	Xtriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.218 , 0.270 0.232 , 0.271	Depositor DCC
R_{free} test set	3263 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	58.2	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11957	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SRE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2564	0.88	1/3476 (0.0%)
1	C	0.62	0/2504	0.91	1/3391 (0.0%)
1	E	0.64	0/2477	0.96	2/3360 (0.1%)
1	G	0.64	0/2157	0.88	0/2921
2	B	0.54	0/593	0.80	0/800
2	D	0.58	0/590	0.86	0/797
2	F	0.67	0/596	0.96	0/804
2	H	0.62	0/587	0.85	0/793
All	All	0.62	0/12068	0.90	4/16342 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	399	PRO	N-CA-C	5.64	117.58	110.70
1	C	235	THR	CB-CA-C	5.35	120.38	112.03
1	A	470	ASN	CA-CB-CG	5.10	117.70	112.60
1	E	470	ASN	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2511	0	2310	9	0
1	C	2454	0	2249	11	0
1	E	2427	0	2194	8	0
1	G	2118	0	1855	10	0
2	B	587	0	604	3	0
2	D	584	0	592	0	0
2	F	590	0	606	2	0
2	H	581	0	588	2	0
3	C	20	0	17	3	0
3	G	20	0	17	3	0
4	A	12	0	0	0	0
4	B	9	0	0	0	0
4	C	12	0	0	1	0
4	D	2	0	0	0	0
4	E	10	0	0	1	0
4	F	5	0	0	0	0
4	G	8	0	0	0	0
4	H	7	0	0	0	0
All	All	11957	0	11032	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:452:ALA:HB3	1:G:517:VAL:HB	1.75	0.69
1:C:533:ASP:HA	1:C:536:ILE:HD12	1.79	0.65
1:A:354:ILE:HD11	1:A:406:LEU:HB3	1.83	0.60
1:A:452:ALA:HB3	1:A:517:VAL:HB	1.85	0.57
2:F:6:LYS:HG2	2:F:12:THR:HG22	1.86	0.57
1:G:285:TRP:CZ2	3:G:601:SRE:H14	2.40	0.57
1:G:302:VAL:HG11	3:G:601:SRE:H18	1.85	0.56
1:C:342:ASP:HB3	1:C:344:ARG:HH22	1.72	0.55
1:A:216:LEU:HG	1:A:267:LEU:HD21	1.91	0.53
1:G:411:TYR:HA	1:G:418:ASN:HA	1.92	0.52
1:G:409:PHE:HB2	2:H:71:LEU:HD12	1.92	0.52
1:C:223:CYS:HB2	1:C:465:TYR:HB2	1.91	0.51
1:C:448:TYR:HB3	1:C:518:TYR:HB3	1.93	0.51
1:C:224:TYR:CE2	3:C:601:SRE:CL20	3.02	0.50
1:E:418:ASN:N	4:E:601:HOH:O	2.30	0.49
1:C:490:LYS:NZ	4:C:701:HOH:O	2.43	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:379:TYR:OH	2:H:64:GLU:OE2	2.30	0.48
1:C:222:THR:HB	1:C:225:MET:HE2	1.96	0.48
1:E:448:TYR:HB3	1:E:518:TYR:HB3	1.96	0.48
1:E:337:VAL:HG21	1:E:384:HIS:ND1	2.29	0.47
1:A:343:ARG:HD3	1:A:343:ARG:HA	1.73	0.46
3:C:601:SRE:H12A	3:C:601:SRE:H3A	1.97	0.45
1:G:285:TRP:HB3	1:G:289:ASP:HB2	1.99	0.45
1:C:224:TYR:HE2	3:C:601:SRE:CL20	2.36	0.44
1:A:346:ASP:O	2:B:46:ALA:HA	2.17	0.44
1:C:470:ASN:HD22	1:C:475:GLY:H	1.66	0.44
2:F:73:LEU:HD23	2:F:73:LEU:HA	1.80	0.44
1:A:214:VAL:HG23	1:A:267:LEU:HD23	1.99	0.44
1:A:347:TYR:OH	1:A:398:LEU:HD23	2.18	0.43
1:E:375:GLY:C	1:E:377:ASN:H	2.27	0.43
1:A:343:ARG:HH22	2:B:62:GLN:HG3	1.84	0.43
1:A:410:MET:HE2	1:A:421:ILE:HG12	2.02	0.42
1:G:302:VAL:HG21	3:G:601:SRE:H17	2.02	0.41
1:G:454:LEU:HB2	1:G:515:MET:HB3	2.03	0.41
1:E:452:ALA:HB3	1:E:517:VAL:HB	2.02	0.41
1:E:469:LEU:O	1:E:471:PRO:HD3	2.21	0.41
1:G:478:CYS:HA	1:G:487:ARG:HA	2.03	0.41
1:E:255:SER:HB2	1:E:257:PRO:HD2	2.01	0.41
1:C:354:ILE:HD11	1:C:406:LEU:HB3	2.03	0.41
2:B:26:VAL:HG21	2:B:56:LEU:HD21	2.03	0.40
1:C:318:GLY:HA2	1:C:321:PRO:HG2	2.03	0.40
1:E:304:LEU:HD22	1:E:320:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/353 (90%)	302 (95%)	16 (5%)	0	100	100
1	C	317/353 (90%)	305 (96%)	11 (4%)	1 (0%)	36	45
1	E	315/353 (89%)	302 (96%)	12 (4%)	1 (0%)	36	45
1	G	288/353 (82%)	268 (93%)	20 (7%)	0	100	100
2	B	73/75 (97%)	70 (96%)	3 (4%)	0	100	100
2	D	73/75 (97%)	69 (94%)	4 (6%)	0	100	100
2	F	73/75 (97%)	70 (96%)	3 (4%)	0	100	100
2	H	73/75 (97%)	69 (94%)	4 (6%)	0	100	100
All	All	1530/1712 (89%)	1455 (95%)	73 (5%)	2 (0%)	48	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	549	ARG
1	E	273	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/320 (81%)	247 (96%)	11 (4%)	26	38
1	C	250/320 (78%)	236 (94%)	14 (6%)	19	27
1	E	246/320 (77%)	236 (96%)	10 (4%)	27	40
1	G	206/320 (64%)	197 (96%)	9 (4%)	25	37
2	B	66/68 (97%)	63 (96%)	3 (4%)	24	37
2	D	64/68 (94%)	63 (98%)	1 (2%)	55	69
2	F	66/68 (97%)	62 (94%)	4 (6%)	17	24
2	H	64/68 (94%)	64 (100%)	0	100	100
All	All	1220/1552 (79%)	1168 (96%)	52 (4%)	26	38

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	223	CYS
1	A	277	LYS
1	A	337	VAL
1	A	343	ARG
1	A	353	SER
1	A	383	GLU
1	A	469	LEU
1	A	470	ASN
1	A	524	LEU
1	A	536	ILE
1	A	539	GLN
2	B	57	SER
2	B	62	GLN
2	B	73	LEU
1	C	217	LYS
1	C	223	CYS
1	C	225	MET
1	C	253	SER
1	C	260	LEU
1	C	320	ILE
1	C	323	LEU
1	C	328	MET
1	C	329	VAL
1	C	336	GLU
1	C	409	PHE
1	C	415	THR
1	C	469	LEU
1	C	483	ASP
2	D	57	SER
1	E	223	CYS
1	E	228	LEU
1	E	257	PRO
1	E	260	LEU
1	E	273	PRO
1	E	360	ILE
1	E	394	LYS
1	E	409	PHE
1	E	469	LEU
1	E	470	ASN
2	F	13	ILE
2	F	33	LYS
2	F	36	ILE
2	F	72	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	214	VAL
1	G	260	LEU
1	G	320	ILE
1	G	340	ARG
1	G	342	ASP
1	G	409	PHE
1	G	449	ILE
1	G	460	ASN
1	G	485	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	HIS
1	A	219	GLN
1	A	294	HIS
1	A	351	GLN
1	A	384	HIS
1	A	405	GLN
2	B	60	ASN
1	C	294	HIS
1	C	447	ASN
1	E	226	ASN
1	E	269	HIS
1	E	438	GLN
1	E	447	ASN
1	E	529	GLN
1	G	460	ASN
1	G	529	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SRE	C	601	-	22,22,22	0.32	0	31,31,31	0.75	0
3	SRE	G	601	-	22,22,22	0.27	0	31,31,31	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SRE	C	601	-	-	0/5/19/19	0/3/3/3
3	SRE	G	601	-	-	1/5/19/19	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	601	SRE	C2-C1-N11-C12

There are no ring outliers.

2 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	601	SRE	3	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	601	SRE	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/353 (92%)	0.33	13 (3%) 42 41	30, 58, 81, 99	0
1	C	325/353 (92%)	0.33	11 (3%) 48 48	44, 62, 81, 92	0
1	E	323/353 (91%)	0.30	7 (2%) 62 64	47, 63, 87, 100	0
1	G	298/353 (84%)	0.75	22 (7%) 20 18	30, 72, 90, 106	0
2	B	75/75 (100%)	-0.08	1 (1%) 75 77	43, 51, 62, 70	0
2	D	75/75 (100%)	0.09	0 100 100	45, 57, 68, 72	0
2	F	75/75 (100%)	0.04	0 100 100	47, 55, 65, 69	0
2	H	75/75 (100%)	0.01	1 (1%) 75 77	48, 56, 67, 70	0
All	All	1572/1712 (91%)	0.34	55 (3%) 47 47	30, 62, 85, 106	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	460	ASN	4.3
1	C	552	ALA	4.1
1	G	342	ASP	3.6
1	G	315	CYS	3.4
1	A	418	ASN	3.3
1	G	248	GLU	3.1
1	E	250	ASP	3.0
1	C	460	ASN	3.0
1	E	509	HIS	2.9
1	G	418	ASN	2.8
1	G	489	THR	2.8
1	A	274	VAL	2.7
1	C	535	ASP	2.7
1	E	460	ASN	2.7
1	G	221	ALA	2.6
1	C	549	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	218	ASN	2.6
1	G	219	GLN	2.6
1	A	442	PRO	2.6
1	A	214	VAL	2.6
1	G	530	ALA	2.5
1	E	508	ARG	2.5
1	E	417	GLN	2.5
1	G	268	GLN	2.5
1	G	475	GLY	2.5
1	A	459	ASP	2.5
1	A	287	THR	2.4
1	G	449	ILE	2.4
1	G	509	HIS	2.4
1	A	285	TRP	2.4
1	C	374	ASP	2.4
1	A	252	SER	2.3
1	C	253	SER	2.3
1	A	221	ALA	2.3
1	E	361	PHE	2.3
1	G	480	PHE	2.3
2	B	75	GLY	2.3
1	G	477	TRP	2.2
1	G	510	CYS	2.2
1	C	221	ALA	2.2
1	G	253	SER	2.2
1	C	534	HIS	2.2
1	G	493	ALA	2.1
2	H	9	THR	2.1
1	C	532	THR	2.1
1	A	248	GLU	2.1
1	G	410	MET	2.1
1	C	314	THR	2.1
1	A	460	ASN	2.1
1	E	211	THR	2.0
1	C	411	TYR	2.0
1	A	210	HIS	2.0
1	G	277	LYS	2.0
1	A	410	MET	2.0
1	G	461	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SRE	C	601	20/20	0.87	0.12	59,64,70,71	0
3	SRE	G	601	20/20	0.87	0.12	59,71,85,90	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SRE C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SRE G 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.