



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:45 AM UTC

PDB ID : 5IKP / pdb_00005ikp
Title : Crystal structure of human brain glycogen phosphorylase bound to AMP
Authors : Mathieu, C.; Li de la Sierra-Gallay, I.; Xu, X.; Haouz, A.; Rodrigues-Lima, F.
Deposited on : 2016-03-03
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	846	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycogen phosphorylase, brain form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	806	6520	4169	1126	1194	31	0	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P11216
A	-2	SER	-	expression tag	UNP P11216
A	-1	HIS	-	expression tag	UNP P11216

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0

- Molecule 3 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: $C_{10}H_{14}N_5O_7P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen phosphorylase, brain form



4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, α , β , γ	172.04Å 172.04Å 123.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.56 – 3.40 47.56 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.56-3.40) 99.8 (47.56-3.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 3.40Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.202 , 0.295 0.203 , 0.295	Depositor DCC
R_{free} test set	768 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	112.6	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 89.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6558	wwPDB-VP
Average B, all atoms (Å ²)	119.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	2/6669 (0.0%)	1.22	6/9028 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	657	ILE	CA-C	5.75	1.57	1.52
1	A	657	ILE	CA-CB	5.06	1.57	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	299	VAL	N-CA-CB	6.11	117.56	110.65
1	A	656	VAL	CA-C-N	5.90	123.96	120.24
1	A	656	VAL	C-N-CA	5.90	123.96	120.24
1	A	288	GLY	N-CA-C	-5.31	106.48	115.34
1	A	288	GLY	CA-C-N	5.30	127.65	120.44
1	A	288	GLY	C-N-CA	5.30	127.65	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6520	0	6468	61	0
2	A	15	0	6	0	0
3	A	23	0	12	0	0
All	All	6558	0	6486	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:THR:HG22	1:A:233:TYR:H	1.29	0.95
1:A:274:ASN:HA	1:A:277:ARG:HB2	1.61	0.81
1:A:168:GLN:HG3	1:A:175:GLN:HG3	1.81	0.61
1:A:95:LEU:HD23	1:A:123:GLU:HG3	1.87	0.56
1:A:196:TYR:HD2	1:A:242:ARG:HH22	1.53	0.56
1:A:582:HIS:HA	1:A:781:MET:HE3	1.88	0.55
1:A:87:LEU:HD13	1:A:341:HIS:HB3	1.89	0.55
1:A:26:GLU:HA	1:A:29:LYS:HD2	1.88	0.55
1:A:702:GLU:HG3	1:A:807:CYS:HB3	1.89	0.53
1:A:828:GLU:HB3	1:A:831:ASP:HB2	1.91	0.52
1:A:233:TYR:HB2	1:A:489:ARG:HD3	1.91	0.52
1:A:162:GLU:HA	1:A:183:LEU:HD12	1.91	0.52
1:A:482:LYS:HE3	1:A:823:GLU:HB2	1.91	0.52
1:A:308:ILE:HD13	1:A:352:ILE:HG21	1.92	0.51
1:A:583:VAL:HG11	1:A:642:VAL:HG22	1.92	0.50
1:A:374:TYR:HB3	1:A:452:VAL:HG22	1.94	0.50
1:A:281:PRO:HG2	1:A:291:LEU:HD23	1.94	0.49
1:A:388:PRO:HD2	1:A:391:MET:HE3	1.96	0.48
1:A:70:THR:HA	1:A:834:ILE:HG12	1.96	0.48
1:A:459:HIS:HA	1:A:462:ILE:HD12	1.95	0.48
1:A:69:ARG:HE	1:A:837:PRO:HB3	1.79	0.47
1:A:81:ARG:HB3	1:A:155:TYR:HE1	1.79	0.47
1:A:490:ARG:HG3	1:A:494:LEU:HD12	1.96	0.47
1:A:329:PHE:HB3	1:A:330:PRO:HD3	1.95	0.47
1:A:578:LEU:HB3	1:A:666:ILE:HD12	1.97	0.47
1:A:567:VAL:HG12	1:A:606:GLY:HA3	1.97	0.47
1:A:503:ILE:HG23	1:A:507:ILE:HD12	1.97	0.46
1:A:144:LEU:HA	1:A:147:MET:HE2	1.97	0.46
1:A:95:LEU:O	1:A:99:MET:HG2	2.16	0.46
1:A:578:LEU:HG	1:A:780:TYR:CD2	2.51	0.45
1:A:662:LEU:HD21	1:A:689:ILE:CD1	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:ALA:HA	1:A:367:ILE:HD12	1.99	0.45
1:A:492:LEU:HD13	1:A:683:LEU:HG	1.99	0.45
1:A:610:ALA:HB3	1:A:613:TYR:HB2	1.98	0.45
1:A:799:LYS:O	1:A:802:ILE:HG22	2.17	0.45
1:A:446:VAL:HG22	1:A:452:VAL:HG21	1.99	0.44
1:A:69:ARG:HA	1:A:72:GLN:HG2	1.99	0.44
1:A:455:VAL:O	1:A:483:THR:HA	2.17	0.43
1:A:772:LYS:HB3	1:A:775:ALA:HB3	1.99	0.43
1:A:381:PRO:HG3	1:A:467:VAL:HA	1.99	0.43
1:A:380:LEU:HA	1:A:381:PRO:HD3	1.95	0.43
1:A:424:ARG:HH12	1:A:473:GLU:CD	2.27	0.43
1:A:530:PHE:CE2	1:A:802:ILE:HD11	2.53	0.43
1:A:231:PRO:HA	1:A:238:VAL:HG22	2.02	0.42
1:A:732:TYR:CG	1:A:766:MET:HE1	2.54	0.42
1:A:731:TYR:CE2	1:A:775:ALA:HA	2.55	0.42
1:A:834:ILE:HA	1:A:835:PRO:HD3	1.93	0.42
1:A:579:ASN:O	1:A:583:VAL:HG23	2.19	0.42
1:A:507:ILE:HD11	1:A:521:LEU:HB3	2.02	0.41
1:A:662:LEU:HD21	1:A:689:ILE:HD12	2.02	0.41
1:A:489:ARG:HH22	1:A:814:ASP:CG	2.29	0.41
1:A:63:LEU:HD23	1:A:63:LEU:HA	1.92	0.41
1:A:101:ASN:HD22	1:A:231:PRO:HG2	1.86	0.41
1:A:89:PHE:O	1:A:131:LEU:HB2	2.21	0.40
1:A:138:ARG:HD2	1:A:138:ARG:HA	1.91	0.40
1:A:272:ALA:HA	1:A:275:ILE:HD12	2.03	0.40
1:A:738:LEU:HB2	1:A:777:TYR:CE2	2.56	0.40
1:A:82:ILE:HA	1:A:334:ALA:HB3	2.03	0.40
1:A:369:LYS:O	1:A:450:HIS:HB3	2.22	0.40
1:A:449:SER:O	1:A:478:LYS:HD2	2.20	0.40
1:A:562:MET:HB3	1:A:601:ARG:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	798/846 (94%)	721 (90%)	69 (9%)	8 (1%)	12	40

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	736	PRO
1	A	772	LYS
1	A	78	ASP
1	A	166	PHE
1	A	380	LEU
1	A	492	LEU
1	A	509	GLU
1	A	794	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	703/737 (95%)	640 (91%)	63 (9%)	9	30

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LEU
1	A	34	HIS
1	A	35	LEU
1	A	39	LEU
1	A	40	VAL
1	A	63	LEU
1	A	64	VAL
1	A	85	LEU
1	A	90	TYR
1	A	96	GLN
1	A	105	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	123	GLU
1	A	169	LYS
1	A	170	ILE
1	A	216	LEU
1	A	224	MET
1	A	228	THR
1	A	234	LYS
1	A	245	SER
1	A	274	ASN
1	A	277	ARG
1	A	299	VAL
1	A	316	PHE
1	A	327	GLU
1	A	328	THR
1	A	340	THR
1	A	363	LYS
1	A	380	LEU
1	A	392	PHE
1	A	431	ILE
1	A	458	ILE
1	A	461	GLU
1	A	474	LEU
1	A	486	ILE
1	A	489	ARG
1	A	502	THR
1	A	505	GLU
1	A	538	LYS
1	A	549	LEU
1	A	560	SER
1	A	568	LYS
1	A	583	VAL
1	A	586	LEU
1	A	591	LYS
1	A	636	VAL
1	A	652	LEU
1	A	667	SER
1	A	678	ASN
1	A	729	ARG
1	A	730	GLU
1	A	735	LEU
1	A	746	SER
1	A	754	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	759	LYS
1	A	781	MET
1	A	790	LEU
1	A	800	LYS
1	A	802	ILE
1	A	813	SER
1	A	816	THR
1	A	828	GLU
1	A	830	SER
1	A	839	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	97	ASN
1	A	101	ASN
1	A	114	GLN
1	A	235	ASN
1	A	282	ASN
1	A	376	ASN
1	A	410	HIS
1	A	459	HIS
1	A	558	ASN
1	A	566	HIS
1	A	665	GLN
1	A	678	ASN
1	A	734	HIS
1	A	782	GLN
1	A	838	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	901	1	15,15,16	2.88	3 (20%)	21,22,23	1.28	2 (9%)
3	AMP	A	902	-	25,25,25	1.56	4 (16%)	37,38,38	2.04	10 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	901	1	-	0/6/6/8	0/1/1/1
3	AMP	A	902	-	-	3/10/26/26	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	PLP	C5-C4	7.94	1.49	1.40
2	A	901	PLP	C3-C2	6.65	1.47	1.41
3	A	902	AMP	C5-C4	5.26	1.48	1.39
2	A	901	PLP	C3-C4	3.25	1.46	1.40
3	A	902	AMP	C5-C6	3.01	1.49	1.41
3	A	902	AMP	C8-N7	2.44	1.36	1.31
3	A	902	AMP	C5-N7	-2.16	1.35	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	AMP	C5-C4-N3	-5.84	118.68	126.72
3	A	902	AMP	N3-C4-N9	4.61	135.00	127.17
3	A	902	AMP	C2-N3-C4	3.83	121.17	111.83
3	A	902	AMP	C4-C5-N7	-3.71	106.34	110.58
3	A	902	AMP	N3-C2-N1	-3.57	123.17	128.58
3	A	902	AMP	C5-N7-C8	2.70	107.70	103.45
3	A	902	AMP	C4-N9-C8	2.49	108.35	105.74
3	A	902	AMP	C3'-C2'-C1'	2.44	106.08	101.46
3	A	902	AMP	C6-C5-N7	2.31	136.55	132.09
2	A	901	PLP	C6-C5-C4	2.21	119.91	118.10
3	A	902	AMP	C2-N1-C6	2.08	122.15	118.73
2	A	901	PLP	O3P-P-O2P	2.01	115.33	107.80

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	AMP	O4'-C4'-C5'-O5'
3	A	902	AMP	C3'-C4'-C5'-O5'
3	A	902	AMP	C4'-C5'-O5'-P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	806/846 (95%)	-0.49	1 (0%) 92 91	78, 117, 154, 190	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	383	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	AMP	A	902	23/23	0.92	0.07	123,130,136,145	0
2	PLP	A	901	15/16	0.95	0.07	101,111,127,139	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AMP A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.