



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:21 PM UTC

PDB ID : 5IKR / pdb_00005ikr
Title : The Structure of Mefenamic Acid Bound to Human Cyclooxygenase-2
Authors : Orlando, B.J.; Malkowski, M.G.
Deposited on : 2016-03-03
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	2% 94% 5%
1	B	551	3% 93% 6%
2	C	3	33% 67%
2	D	3	33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	D	2	-	-	X	-
4	COH	B	603	X	-	-	-

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 18781 atoms, of which 9037 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin G/H synthase 2.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	551	Total	C	H	N	O	S	77	0	0
			8825	2883	4355	755	806	26			
1	B	551	Total	C	H	N	O	S	243	0	0
			8826	2883	4356	755	806	26			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	3	Total	C	H	N	O	0	0	0
			78	22	39	2	15			
2	D	3	Total	C	H	N	O	0	0	0
			78	22	39	2	15			

- Molecule 3 is 2-[(2,3-DIMETHYLPHENYL)AMINO]BENZOIC ACID (CCD ID: ID8) (formula: C₁₅H₁₅NO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			32	15	14	1	2		
3	B	1	Total	C	H	N	O	0	0
			32	15	14	1	2		

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING CO (CCD ID: COH) (formula: $C_{34}H_{32}CoN_4O_4$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	Co	H	N	O	0	0
			73	34	1	30	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	Co	H	N	O	0	0
			73	34	1	30	4	4		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 6 is octyl beta-D-glucopyranoside (CCD ID: BOG) (formula: $C_{14}H_{28}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			48	14	28	6		
6	A	1	Total	C	H	O	0	0
			48	14	28	6		
6	A	1	Total	C	H	O	0	0
			48	14	28	6		
6	B	1	Total	C	H	O	0	0
			48	14	28	6		

- Molecule 7 is AMMONIUM ION (CCD ID: NH4) (formula: H_4N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	H	N	0	0
			5	4	1		
7	A	1	Total	H	N	0	0
			5	4	1		
7	A	1	Total	H	N	0	0
			5	4	1		
7	B	1	Total	H	N	0	0
			5	4	1		
7	B	1	Total	H	N	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	285	Total	O	0	0
			285	285		
8	B	206	Total	O	0	0
			206	206		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Prostaglandin G/H synthase 2



- Molecule 1: Prostaglandin G/H synthase 2



- Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	127.67Å 149.69Å 188.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.34 20.00 – 2.34	Depositor EDS
% Data completeness (in resolution range)	95.4 (20.00-2.34) 95.2 (20.00-2.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.35Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.211 0.191 , 0.214	Depositor DCC
R_{free} test set	3580 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18781	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COH, BOG, NAG, ID8, NH4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/4602	0.65	0/6241
1	B	0.38	1/4602 (0.0%)	0.66	0/6241
All	All	0.36	1/9204 (0.0%)	0.65	0/12482

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	385	TYR	C-O	-5.50	1.16	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4470	4355	4353	20	0
1	B	4470	4356	4353	33	0
2	C	39	39	34	5	0
2	D	39	39	34	7	0
3	A	18	14	14	2	0
3	B	18	14	14	6	0
4	A	43	30	30	2	0
4	B	43	30	30	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	14	14	13	0	0
5	B	14	14	12	0	0
6	A	60	84	84	2	0
6	B	20	28	28	1	0
7	A	3	12	0	0	0
7	B	2	8	0	0	0
8	A	285	0	0	1	0
8	B	206	0	0	2	0
All	All	9744	9037	8999	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:ARG:HH11	2:D:2:NAG:H83	1.28	0.97
1:B:216:ARG:HH11	2:D:2:NAG:C8	1.83	0.90
1:B:216:ARG:NH1	2:D:2:NAG:H83	1.91	0.85
1:A:216:ARG:HD3	2:C:2:NAG:H81	1.63	0.80
1:A:216:ARG:HH11	2:C:2:NAG:H83	1.54	0.73
1:B:216:ARG:NH2	8:B:701:HOH:O	2.13	0.70
1:B:216:ARG:HD3	2:D:2:NAG:H81	1.77	0.67
4:B:603:COH:HBB1	4:B:603:COH:HHC	1.78	0.65
4:A:602:COH:HBB1	4:A:602:COH:HHC	1.79	0.64
1:B:185:ARG:NH1	8:B:704:HOH:O	2.34	0.59
1:B:160:PRO:HG2	1:B:165:VAL:HA	1.85	0.59
1:B:160:PRO:CG	1:B:165:VAL:HG12	2.35	0.56
1:B:216:ARG:HD3	2:D:2:NAG:C8	2.38	0.53
1:B:388:HIS:N	1:B:389:PRO:HD2	2.25	0.51
1:B:159:CYS:HB3	1:B:164:GLY:O	2.11	0.49
3:B:602:ID8:H9	3:B:602:ID8:C5	2.43	0.49
1:A:216:ARG:HH11	2:C:2:NAG:C8	2.26	0.48
1:B:523:VAL:HG13	3:B:602:ID8:C4	2.43	0.48
1:B:294:LEU:HD22	1:B:408:ILE:HG23	1.96	0.48
1:B:216:ARG:HH11	2:D:2:NAG:H81	1.76	0.48
1:B:391:LEU:HB2	1:B:441:PRO:HG2	1.95	0.47
1:B:504:TYR:HB3	1:B:505:PRO:HD3	1.96	0.47
1:A:53:ASP:OD1	1:A:53:ASP:N	2.48	0.47
1:A:216:ARG:CD	2:C:2:NAG:H81	2.38	0.47
1:B:160:PRO:HG2	1:B:165:VAL:HG12	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:PRO:HG3	1:B:165:VAL:HG12	1.97	0.45
3:A:601:ID8:H9	3:A:601:ID8:C5	2.46	0.45
1:A:125:ASP:OD2	8:A:701:HOH:O	2.21	0.45
1:B:352:LEU:HD11	1:B:518:PHE:CE2	2.52	0.44
1:B:216:ARG:NH1	2:D:2:NAG:C8	2.63	0.44
1:A:527:ALA:HB3	1:A:528:PRO:HD3	1.99	0.44
1:A:567:LEU:HA	6:A:609:BOG:H8'1	1.98	0.44
1:B:417:HIS:HB3	1:B:421:GLN:HG2	2.01	0.43
1:A:294:LEU:HA	1:A:409:TYR:CE1	2.53	0.43
4:A:602:COH:HMC1	4:A:602:COH:HBC1	2.00	0.43
1:B:506:ALA:HA	1:B:509:VAL:HG22	2.01	0.43
1:B:567:LEU:HA	6:B:601:BOG:H8'1	2.00	0.43
1:A:388:HIS:CE1	1:A:447:VAL:HG11	2.53	0.43
1:B:39:HIS:N	1:B:40:PRO:HD3	2.33	0.43
6:A:609:BOG:H2'1	6:A:609:BOG:H1	1.91	0.43
1:B:523:VAL:HG13	3:B:602:ID8:H4	2.01	0.43
4:B:603:COH:HMC1	4:B:603:COH:HBC1	2.01	0.43
1:A:388:HIS:N	1:A:389:PRO:CD	2.82	0.42
1:A:484:GLU:HG2	1:A:485:LYS:H	1.84	0.42
3:B:602:ID8:C5	3:B:602:ID8:C9	2.96	0.42
1:A:380:GLU:HG2	1:A:466:TYR:CE1	2.55	0.42
3:B:602:ID8:C9	3:B:602:ID8:H5	2.50	0.42
1:B:463:PHE:CD1	1:B:506:ALA:HB3	2.56	0.41
1:A:216:ARG:HD3	2:C:2:NAG:C8	2.41	0.41
1:A:127:PRO:HB2	1:B:538:VAL:HG23	2.03	0.41
1:B:294:LEU:HD13	1:B:295:VAL:HG13	2.02	0.41
1:A:538:VAL:HG23	1:B:127:PRO:HG2	2.02	0.41
1:B:97:LYS:HG3	1:B:356:HIS:CE1	2.56	0.41
1:B:160:PRO:HG2	1:B:164:GLY:O	2.21	0.41
1:B:406:GLN:HE21	1:B:406:GLN:HB3	1.77	0.41
1:A:230:LEU:HG	1:A:337:ILE:HG12	2.03	0.41
1:A:156:PRO:HG2	1:A:159:CYS:SG	2.61	0.40
1:A:172:PRO:HG2	1:A:177:ILE:HD11	2.03	0.40
1:A:233:ILE:HD13	1:A:305:TRP:HB3	2.04	0.40
3:A:601:ID8:C5	3:A:601:ID8:C9	2.96	0.40
1:B:385:TYR:CE1	3:B:602:ID8:H12	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/551 (100%)	533 (97%)	15 (3%)	1 (0%)	43	50
1	B	549/551 (100%)	530 (96%)	19 (4%)	0	100	100
All	All	1098/1102 (100%)	1063 (97%)	34 (3%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	PHE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/490 (100%)	482 (98%)	8 (2%)	55	67
1	B	490/490 (100%)	478 (98%)	12 (2%)	43	55
All	All	980/980 (100%)	960 (98%)	20 (2%)	48	61

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	PHE
1	A	75	LEU
1	A	216	ARG
1	A	289	GLN
1	A	295	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	385	TYR
1	A	428	ARG
1	A	484	GLU
1	B	70	SER
1	B	157	ASP
1	B	216	ARG
1	B	232	HIS
1	B	267	LYS
1	B	289	GLN
1	B	294	LEU
1	B	295	VAL
1	B	385	TYR
1	B	406	GLN
1	B	459	LYS
1	B	484	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	386	HIS
1	A	421	GLN
1	B	42	GLN
1	B	241	GLN
1	B	398	HIS
1	B	421	GLN
1	B	457	GLN
1	B	461	GLN
1	B	571	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.36	0	17,19,21	0.52	0
2	NAG	C	2	2	14,14,15	0.37	0	17,19,21	0.63	0
2	MAN	C	3	2	11,11,12	0.97	1 (9%)	15,15,17	1.52	2 (13%)
2	NAG	D	1	1,2	14,14,15	0.31	0	17,19,21	0.53	0
2	NAG	D	2	2	14,14,15	0.27	0	17,19,21	0.59	0
2	MAN	D	3	2	11,11,12	0.96	2 (18%)	15,15,17	1.38	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsions and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	MAN	C	3	2	-	2/2/19/22	1/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	4/6/23/26	0/1/1/1
2	MAN	D	3	2	-	0/2/19/22	1/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	MAN	C1-C2	2.49	1.58	1.52
2	D	3	MAN	C1-C2	2.21	1.57	1.52
2	D	3	MAN	O5-C5	2.05	1.47	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	MAN	C1-O5-C5	4.26	117.90	112.19
2	D	3	MAN	C1-O5-C5	3.94	117.47	112.19
2	C	3	MAN	O2-C2-C3	-3.14	103.65	110.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	MAN	O2-C2-C3	-2.92	104.10	110.15

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	C	3	MAN	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	C	3	MAN	O5-C5-C6-O6



All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	MAN	C1-C2-C3-C4-C5-O5
2	D	3	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	NAG	7	0
2	C	2	NAG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 15 ligands modelled in this entry, 5 are modelled with single atom - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	BOG	A	607	-	20,20,20	1.17	1 (5%)	25,25,25	0.82	0
6	BOG	A	609	-	20,20,20	1.14	1 (5%)	25,25,25	1.06	1 (4%)
3	ID8	A	601	-	19,19,19	2.04	9 (47%)	26,26,26	1.69	4 (15%)
4	COH	A	602	1	50,50,50	1.62	8 (16%)	66,82,82	1.41	9 (13%)
6	BOG	B	601	-	20,20,20	1.15	1 (5%)	25,25,25	0.81	0
6	BOG	A	608	-	20,20,20	1.18	1 (5%)	25,25,25	0.78	0
5	NAG	A	606	1	14,14,15	0.22	0	17,19,21	0.50	0
3	ID8	B	602	-	19,19,19	1.65	3 (15%)	26,26,26	2.06	5 (19%)
5	NAG	B	607	1	14,14,15	0.21	0	17,19,21	0.46	0
4	COH	B	603	1	50,50,50	1.59	9 (18%)	66,82,82	1.42	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BOG	A	607	-	-	5/11/31/31	0/1/1/1
6	BOG	A	609	-	-	6/11/31/31	0/1/1/1
3	ID8	A	601	-	-	0/8/8/8	0/2/2/2
4	COH	A	602	1	-	1/14/54/54	-
6	BOG	B	601	-	-	5/11/31/31	0/1/1/1
6	BOG	A	608	-	-	3/11/31/31	0/1/1/1
3	ID8	B	602	-	-	0/8/8/8	0/2/2/2
5	NAG	A	606	1	-	0/6/23/26	0/1/1/1
4	COH	B	603	1	1/1/9/9	6/14/54/54	-
5	NAG	B	607	1	-	0/6/23/26	0/1/1/1

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	603	COH	C3D-C2D	6.02	1.54	1.38
4	A	602	COH	C3D-C2D	5.94	1.54	1.38
4	A	602	COH	CO-ND	4.23	2.12	1.96
4	B	603	COH	CO-ND	3.49	2.09	1.96
3	A	601	ID8	O15-C14	-3.27	1.20	1.30
4	B	603	COH	CAB-C3B	3.26	1.56	1.47
4	B	603	COH	CAC-C3C	3.16	1.55	1.47
4	A	602	COH	CAB-C3B	3.15	1.55	1.47
4	A	602	COH	CAC-C3C	3.13	1.55	1.47
4	B	603	COH	CO-NB	3.08	2.08	1.96
4	A	602	COH	CO-NB	3.00	2.07	1.96
3	B	602	ID8	O15-C14	-2.98	1.21	1.30
6	A	608	BOG	O5-C1	2.93	1.49	1.41
6	A	607	BOG	O5-C1	2.88	1.49	1.41
6	B	601	BOG	O5-C1	2.75	1.48	1.41
4	A	602	COH	CO-NA	2.74	2.06	1.96
3	A	601	ID8	C9-C8	-2.68	1.35	1.39
6	A	609	BOG	O5-C1	2.66	1.48	1.41
3	A	601	ID8	C8-N7	-2.48	1.33	1.39
3	A	601	ID8	C6-N7	-2.41	1.33	1.39
3	A	601	ID8	C10-C9	-2.27	1.35	1.38
3	A	601	ID8	C3-C2	-2.26	1.35	1.39
4	B	603	COH	CMC-C2C	2.25	1.55	1.50
4	A	602	COH	CMC-C2C	2.20	1.55	1.50
4	B	603	COH	CO-NA	2.18	2.04	1.96
4	B	603	COH	CMB-C2B	2.17	1.55	1.50
3	B	602	ID8	C8-N7	-2.15	1.34	1.39
4	B	603	COH	CMA-C3A	2.13	1.55	1.50
3	A	601	ID8	C2-C1	-2.13	1.36	1.40
4	A	602	COH	CMB-C2B	2.13	1.55	1.50
3	A	601	ID8	C11-C12	-2.10	1.35	1.38
3	B	602	ID8	C6-N7	-2.06	1.34	1.39
3	A	601	ID8	C4-C3	-2.04	1.35	1.38

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	ID8	C8-C13-C14	5.46	127.80	121.72
3	B	602	ID8	O15-C14-O16	-5.11	112.37	123.35
3	A	601	ID8	O15-C14-O16	-4.78	113.07	123.35
4	B	603	COH	C4D-ND-C1D	4.63	111.94	105.11
4	B	603	COH	C2B-C1B-NB	-4.40	107.46	110.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	602	COH	C2B-C1B-NB	-4.39	107.47	110.88
3	B	602	ID8	O15-C14-C13	4.38	127.73	115.28
4	A	602	COH	C4D-ND-C1D	4.34	111.51	105.11
3	A	601	ID8	O15-C14-C13	4.12	126.99	115.28
3	A	601	ID8	C8-C13-C14	3.73	125.88	121.72
4	A	602	COH	C3A-C4A-NA	-3.23	107.60	110.96
6	A	609	BOG	C1-O5-C5	-3.17	107.52	113.72
4	B	603	COH	C3A-C4A-NA	-2.80	108.04	110.96
4	B	603	COH	CMC-C2C-C1C	-2.62	120.93	125.03
4	A	602	COH	C4B-NB-C1B	2.51	108.81	105.11
4	A	602	COH	CMC-C2C-C1C	-2.51	121.12	125.03
4	B	603	COH	C4B-NB-C1B	2.50	108.80	105.11
4	B	603	COH	C1A-NA-C4A	2.39	108.15	105.12
4	A	602	COH	C1A-NA-C4A	2.29	108.04	105.12
4	A	602	COH	C2C-C1C-NC	-2.27	108.60	110.96
4	B	603	COH	CHC-C4B-C3B	2.18	128.88	125.21
3	B	602	ID8	C5-C6-C1	-2.10	118.24	121.08
4	A	602	COH	C4C-NC-C1C	2.09	107.78	105.12
4	A	602	COH	CHC-C4B-C3B	2.03	128.63	125.21
3	B	602	ID8	C11-C12-C13	2.03	123.46	119.80
3	A	601	ID8	C6-N7-C8	2.02	131.73	125.41
4	B	603	COH	C2C-C1C-NC	-2.02	108.86	110.96
4	B	603	COH	C4C-NC-C1C	2.02	107.68	105.12

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	603	COH	NB

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	603	COH	C1A-C2A-CAA-CBA
6	A	608	BOG	C2'-C1'-O1-C1
6	A	609	BOG	C2'-C1'-O1-C1
6	A	607	BOG	O5-C5-C6-O6
4	B	603	COH	C3A-C2A-CAA-CBA
6	A	607	BOG	C4-C5-C6-O6
6	A	609	BOG	O5-C5-C6-O6
6	B	601	BOG	C4-C5-C6-O6
6	A	609	BOG	C4-C5-C6-O6
6	B	601	BOG	O1-C1'-C2'-C3'

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
6	B	601	BOG	O5-C5-C6-O6
6	A	607	BOG	C4'-C5'-C6'-C7'
4	B	603	COH	C2A-CAA-CBA-CGA
6	A	607	BOG	C3'-C4'-C5'-C6'
6	A	609	BOG	O1-C1'-C2'-C3'
6	A	607	BOG	C2'-C1'-O1-C1
4	A	602	COH	C4B-C3B-CAB-CBB
4	B	603	COH	C4B-C3B-CAB-CBB
6	B	601	BOG	C2'-C1'-O1-C1
6	A	609	BOG	C3'-C4'-C5'-C6'
6	B	601	BOG	C2'-C3'-C4'-C5'
4	B	603	COH	CAA-CBA-CGA-O1A
6	A	609	BOG	C1'-C2'-C3'-C4'
4	B	603	COH	CAA-CBA-CGA-O2A
6	A	608	BOG	C5'-C6'-C7'-C8'
6	A	608	BOG	C3'-C4'-C5'-C6'


There are no ring outliers.



6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	609	BOG	2	0
3	A	601	ID8	2	0
4	A	602	COH	2	0
6	B	601	BOG	1	0
3	B	602	ID8	6	0
4	B	603	COH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/551 (100%)	-0.17	9 (1%) 70 75	14, 36, 64, 104	9 (1%)
1	B	551/551 (100%)	0.20	18 (3%) 49 55	18, 50, 85, 103	25 (4%)
All	All	1102/1102 (100%)	0.02	27 (2%) 58 64	14, 40, 78, 104	34 (3%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	PHE	4.7
1	B	157	ASP	3.7
1	B	409	TYR	3.7
1	B	158	ASP	3.5
1	B	82	LEU	3.5
1	B	107	PHE	3.2
1	B	483	GLY	3.1
1	A	272	GLU	2.7
1	B	85	THR	2.6
1	B	174	SER	2.5
1	A	409	TYR	2.5
1	B	34	ASN	2.5
1	A	107	PHE	2.4
1	A	50	VAL	2.4
1	B	81	PHE	2.4
1	A	157	ASP	2.4
1	A	583	PRO	2.3
1	B	278	GLN	2.3
1	B	96	PHE	2.2
1	A	486	GLU	2.2
1	B	168	LYS	2.1
1	B	187	PHE	2.1
1	B	496	GLY	2.1
1	A	122	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	97	LYS	2.1
1	B	428	ARG	2.0
1	B	185	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MAN	D	3	11/12	0.51	0.19	61,74,88,95	0
2	MAN	C	3	11/12	0.57	0.20	62,76,91,95	0
2	NAG	C	2	14/15	0.84	0.13	31,48,61,74	0
2	NAG	D	2	14/15	0.89	0.10	32,45,56,68	0
2	NAG	D	1	14/15	0.95	0.06	24,31,36,42	0
2	NAG	C	1	14/15	0.96	0.05	18,25,34,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	BOG	A	609	20/20	0.71	0.28	30,134,181,187	0
6	BOG	B	601	20/20	0.76	0.22	32,113,149,154	0
7	NH4	B	609	1/1	0.80	0.30	48,58,58,58	0
5	NAG	B	607	14/15	0.82	0.14	54,71,86,86	0
6	BOG	A	607	20/20	0.83	0.21	38,103,134,204	0
6	BOG	A	608	20/20	0.84	0.19	38,100,124,124	0
7	NH4	A	611	1/1	0.87	0.21	35,43,43,43	0
5	NAG	A	606	14/15	0.88	0.08	33,47,57,59	0
3	ID8	B	602	18/18	0.93	0.08	30,37,45,47	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NH4	A	612	1/1	0.93	0.24	32,39,39,39	0
7	NH4	A	610	1/1	0.93	0.21	24,29,29,29	0
4	COH	A	602	43/43	0.94	0.12	24,50,73,79	0
4	COH	B	603	43/43	0.94	0.14	36,72,88,94	0
3	ID8	A	601	18/18	0.95	0.06	14,22,28,29	0
7	NH4	B	608	1/1	0.96	0.26	27,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around BOG A 609:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG A 607:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG A 608:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COH A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around COH B 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.