



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:06 AM UTC

PDB ID : 2ILN / pdb_00002iln
Title : Crystal structure of the Bowman-Birk inhibitor from snail medic seeds in complex with bovine trypsin
Authors : Capaldi, S.; Perduca, M.; Faggion, B.; Carrizo, M.E.; Tava, A.; Ragona, L.; Monaco, H.L.
Deposited on : 2006-10-03
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	82% 16% .
1	B	223	79% 19% .
2	I	62	61% 19% . . 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cationic trypsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1629	1012	279	324	14			
1	B	223	Total	C	N	O	S	0	0	0
			1629	1012	279	324	14			

- Molecule 2 is a protein called Bowman-Birk type proteinase inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	53	Total	C	N	O	S	0	0	0
			411	249	72	77	13			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	169	Total	O	0	0
			169	169		
3	B	175	Total	O	0	0
			175	175		
3	I	32	Total	O	0	0
			32	32		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cationic trypsin



• Molecule 1: Cationic trypsin



• Molecule 2: Bowman-Birk type proteinase inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	54.47Å 54.47Å 180.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.00 25.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.6 (25.00-2.00) 95.6 (25.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.232 , 0.287 0.230 , 0.281	Depositor DCC
R_{free} test set	1680 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.926	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.437 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4045	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/1660	0.86	3/2250 (0.1%)
1	B	0.60	0/1660	0.85	1/2250 (0.0%)
2	I	0.67	1/422 (0.2%)	1.08	3/572 (0.5%)
All	All	0.60	1/3742 (0.0%)	0.88	7/5072 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	57	CYS	CA-C	5.46	1.56	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	57	CYS	N-CA-C	9.35	125.23	111.02
1	A	151	TYR	CA-C-N	7.81	129.60	119.84
1	A	151	TYR	C-N-CA	7.81	129.60	119.84
1	A	152	PRO	N-CA-C	5.61	124.02	112.47
1	B	211	GLY	N-CA-C	5.26	118.80	110.96
2	I	57	CYS	CA-C-N	5.15	130.97	121.70
2	I	57	CYS	C-N-CA	5.15	130.97	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1588	33	0
1	B	1629	0	1588	35	0
2	I	411	0	370	17	0
3	A	169	0	0	4	0
3	B	175	0	0	2	0
3	I	32	0	0	0	0
All	All	4045	0	3546	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:31:CYS:HA	2:I:32:HIS:HB2	1.31	1.09
2:I:58:TYR:HB3	2:I:59:PRO:HD2	1.46	0.98
1:B:100:ASN:HD21	1:B:179:ASN:HD22	1.19	0.90
2:I:58:TYR:HB3	2:I:59:PRO:CD	2.03	0.89
1:A:100:ASN:HD21	1:A:179:ASN:HD22	1.21	0.88
1:A:100:ASN:ND2	1:A:179:ASN:HD22	1.74	0.86
2:I:28:ARG:HH21	2:I:32:HIS:HE1	1.24	0.82
1:B:48:ASN:HD22	1:B:50:GLN:H	1.30	0.77
1:A:48:ASN:HD22	1:A:50:GLN:H	1.33	0.76
2:I:58:TYR:CB	2:I:59:PRO:HD2	2.16	0.75
1:B:64:GLN:HE22	1:B:66:ARG:HH21	1.36	0.72
1:B:100:ASN:ND2	1:B:179:ASN:HD22	1.88	0.71
1:A:100:ASN:HD21	1:A:179:ASN:ND2	1.88	0.71
2:I:58:TYR:CB	2:I:59:PRO:CD	2.70	0.69
1:B:151:TYR:CE1	2:I:44:PHE:CZ	2.81	0.69
1:A:64:GLN:HE22	1:A:66:ARG:HH21	1.41	0.68
1:B:64:GLN:NE2	1:B:66:ARG:HE	1.92	0.66
2:I:31:CYS:HA	2:I:32:HIS:CB	2.16	0.66
1:B:143:ASN:HD22	1:B:144:THR:N	1.94	0.64
1:B:143:ASN:HD21	1:B:150:SER:H	1.45	0.63
1:B:100:ASN:HD21	1:B:179:ASN:ND2	1.94	0.63
2:I:37:SER:HB3	2:I:51:TYR:CE2	2.34	0.62
1:A:48:ASN:ND2	1:A:50:GLN:H	1.96	0.61
1:B:48:ASN:ND2	1:B:50:GLN:H	1.96	0.60
1:A:149:THR:HG23	1:A:151:TYR:CE2	2.37	0.60
1:B:143:ASN:HD22	1:B:144:THR:H	1.48	0.59
1:B:151:TYR:CE1	2:I:44:PHE:HZ	2.20	0.59
1:A:149:THR:HG23	1:A:151:TYR:HE2	1.69	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ASP:HB3	3:A:398:HOH:O	2.04	0.57
1:B:103:ILE:HD13	1:B:104:MET:N	2.20	0.56
1:B:143:ASN:ND2	1:B:150:SER:H	2.03	0.56
1:A:87:LYS:HD2	3:A:403:HOH:O	2.04	0.56
1:A:115:ASN:HD22	1:A:117:ARG:H	1.54	0.55
1:A:217:SER:HB3	1:A:224:LYS:HD2	1.87	0.55
1:A:143:ASN:HD22	1:A:144:THR:N	2.06	0.54
1:B:48:ASN:HD22	1:B:48:ASN:C	2.14	0.54
2:I:13:PRO:HD2	2:I:23:GLN:O	2.08	0.53
1:A:210:GLN:NE2	3:A:404:HOH:O	2.42	0.53
1:A:101:ASN:HA	1:A:234:TYR:OH	2.09	0.52
1:A:240:GLN:HG3	3:A:304:HOH:O	2.08	0.52
2:I:28:ARG:HH21	2:I:32:HIS:CE1	2.14	0.52
1:A:158:LEU:HD11	1:A:188:LYS:HB3	1.91	0.52
1:B:138:ILE:HG12	1:B:199:VAL:HG22	1.93	0.51
1:B:103:ILE:HD12	1:B:237:TRP:HZ3	1.75	0.50
1:B:158:LEU:HD11	1:B:188:LYS:HB3	1.94	0.49
1:B:64:GLN:HE21	1:B:66:ARG:HE	1.57	0.49
1:A:143:ASN:HD21	1:A:150:SER:H	1.59	0.48
1:B:213:VAL:HA	1:B:228:TYR:CD1	2.47	0.48
1:B:48:ASN:HD21	1:B:51:TRP:HD1	1.60	0.48
1:A:149:THR:CG2	1:A:151:TYR:HE2	2.27	0.48
1:B:69:GLU:HB2	3:B:246:HOH:O	2.14	0.47
1:A:124:PRO:HD3	1:A:209:LEU:O	2.15	0.47
1:B:151:TYR:HE1	2:I:44:PHE:CZ	2.32	0.47
1:B:121:ILE:HD13	1:B:122:SER:N	2.29	0.47
2:I:10:ASP:HA	2:I:60:SER:HB3	1.97	0.47
1:A:211:GLY:HA2	1:A:229:THR:O	2.15	0.47
1:B:184:TYR:CE1	1:B:188:LYS:HE3	2.51	0.45
1:B:91:HIS:CG	1:B:92:PRO:HD2	2.51	0.45
1:B:162:ILE:HD13	1:B:181:PHE:CE1	2.52	0.45
1:A:97:ASN:ND2	1:B:173:PRO:O	2.50	0.45
1:A:91:HIS:CG	1:A:92:PRO:HD2	2.52	0.45
1:A:143:ASN:HD22	1:A:144:THR:H	1.64	0.45
1:A:195:SER:OG	2:I:16:ARG:C	2.60	0.45
1:A:64:GLN:NE2	1:A:66:ARG:HE	2.14	0.45
1:A:27:VAL:HG13	1:A:29:TYR:CZ	2.52	0.44
1:B:100:ASN:ND2	1:B:179:ASN:HB2	2.32	0.44
1:A:146:SER:HB3	1:A:221(A):ALA:HB3	1.99	0.44
2:I:58:TYR:N	2:I:58:TYR:CD1	2.86	0.43
1:A:115:ASN:ND2	1:A:117:ARG:H	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:ASN:ND2	1:A:150:SER:H	2.17	0.43
1:A:213:VAL:HA	1:A:228:TYR:CD1	2.53	0.43
1:B:160:ALA:HB1	1:B:184(A):GLY:HA2	2.01	0.43
1:A:210:GLN:HE21	1:A:210:GLN:HA	1.84	0.43
1:B:115:ASN:ND2	1:B:117:ARG:H	2.17	0.42
1:B:72:ASN:OD1	1:B:153:ASP:HB3	2.19	0.42
1:B:233:ASN:ND2	3:B:387:HOH:O	2.53	0.41
2:I:23:GLN:HB3	2:I:53:ILE:HG13	2.01	0.41
1:A:115:ASN:ND2	1:A:118:VAL:H	2.19	0.41
1:B:210:GLN:HA	1:B:210:GLN:HE21	1.86	0.41
1:B:101:ASN:HA	1:B:234:TYR:OH	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/223 (99%)	215 (97%)	5 (2%)	1 (0%)	24	21
1	B	221/223 (99%)	217 (98%)	4 (2%)	0	100	100
2	I	51/62 (82%)	43 (84%)	6 (12%)	2 (4%)	2	1
All	All	493/508 (97%)	475 (96%)	15 (3%)	3 (1%)	21	17

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	I	32	HIS
2	I	58	TYR
1	A	152	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/184 (100%)	180 (98%)	4 (2%)	45	50
1	B	184/184 (100%)	174 (95%)	10 (5%)	20	17
2	I	52/60 (87%)	50 (96%)	2 (4%)	29	29
All	All	420/428 (98%)	404 (96%)	16 (4%)	29	29

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	109	LYS
1	A	115	ASN
1	A	165	ASP
1	B	31	VAL
1	B	48	ASN
1	B	89	ILE
1	B	103	ILE
1	B	121	ILE
1	B	147	SER
1	B	162	ILE
1	B	186	GLU
1	B	201	CYS
1	B	244	SER
2	I	57	CYS
2	I	58	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	48	ASN
1	A	64	GLN
1	A	74	ASN
1	A	100	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	115	ASN
1	A	143	ASN
1	A	175	GLN
1	A	192	GLN
1	A	210	GLN
1	A	233	ASN
1	A	240	GLN
1	B	48	ASN
1	B	64	GLN
1	B	74	ASN
1	B	100	ASN
1	B	115	ASN
1	B	135	GLN
1	B	143	ASN
1	B	175	GLN
1	B	192	GLN
1	B	210	GLN
1	B	233	ASN
2	I	23	GLN
2	I	32	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/223 (100%)	-0.93	0 100 100	20, 27, 35, 48	0
1	B	223/223 (100%)	-0.92	0 100 100	18, 28, 37, 45	0
2	I	53/62 (85%)	-0.67	0 100 100	21, 36, 54, 58	0
All	All	499/508 (98%)	-0.90	0 100 100	18, 28, 42, 58	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.