



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:47 PM UTC

PDB ID : 3ILF / pdb_00003ilf
Title : Crystal structure of porphyranase A (PorA) in complex with neoporphyrinotetraose
Authors : Hehemann, J.H.; Correc, G.; Barbeyron, T.; Helbert, W.; Michel, G.; Czjzek, M.
Deposited on : 2009-08-07
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	3% 83% 11% ...
2	B	4	75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	L6S	B	4	X	-	-	-
6	ACT	A	282	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 2556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called porphyranase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	8	0
			2159	1386	367	400	6			

- Molecule 2 is an oligosaccharide called 6-O-sulfo-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-6-O-sulfo-alpha-L-galactopyranose-(1-3)-alpha-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	4	Total	C	O	S	0	0	0
			53	24	27	2			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	324	Total	O	0	0
			324	324		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: porphyranase A

Chain A: 



- Molecule 2: 6-O-sulfo-alpha-L-galactopyranose-(1-3)-beta-D-galactopyranose-(1-4)-6-O-sulfo-alpha-L-galactopyranose-(1-3)-alpha-D-galactopyranose

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.09Å 68.38Å 71.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.34 – 1.80 49.34 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.0 (49.34-1.80) 94.0 (49.34-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.93 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.157 , 0.205 (Not available) , 0.196	Depositor DCC
R_{free} test set	1381 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	11.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2556	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, GLA, CL, ACT, L6S, MG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.41	7/2236 (0.3%)	1.22	4/3032 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	GLY	N-CA	7.93	1.50	1.44
1	A	135	GLY	C-O	7.52	1.30	1.23
1	A	69	ALA	CA-CB	6.97	1.62	1.53
1	A	75	PHE	C-O	6.46	1.31	1.23
1	A	58	GLY	N-CA	6.04	1.52	1.45
1	A	209	PHE	C-O	5.42	1.30	1.24
1	A	149	PRO	C-O	5.15	1.29	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	ASP	N-CA-CB	-5.86	103.79	111.23
1	A	148	ASP	CA-C-N	5.43	125.35	119.76
1	A	148	ASP	C-N-CA	5.43	125.35	119.76
1	A	25	ASN	N-CA-C	5.12	118.66	111.74

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2159	0	2063	35	0
2	B	53	0	36	1	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	16	0	12	5	0
7	A	324	0	0	8	2
All	All	2556	0	2111	35	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:ALA:HB2	7:A:449:HOH:O	1.60	1.00
1:A:240[B]:THR:HG23	1:A:241[B]:ALA:N	1.83	0.93
1:A:266:TRP:HE1	6:A:285:ACT:H2	1.33	0.91
1:A:259:LYS:HZ3	6:A:282:ACT:H3	1.39	0.84
1:A:259:LYS:NZ	6:A:282:ACT:H3	1.91	0.84
1:A:240[B]:THR:HG23	1:A:242[B]:GLY:N	1.97	0.79
1:A:240[B]:THR:O	1:A:241[B]:ALA:C	2.28	0.76
1:A:240[B]:THR:HG22	1:A:243:VAL:N	2.01	0.75
1:A:240[B]:THR:HG23	1:A:242[B]:GLY:H	1.55	0.70
1:A:245:ASN:HB2	7:A:567:HOH:O	1.91	0.69
1:A:28:LYS:N	1:A:28:LYS:HE2	2.08	0.69
1:A:240[B]:THR:HG23	1:A:241[B]:ALA:CA	2.22	0.68
1:A:240[B]:THR:CG2	1:A:242[B]:GLY:H	2.06	0.68
1:A:273:VAL:O	1:A:274:ASP:HB2	1.94	0.67
1:A:42:ASN:ND2	7:A:421:HOH:O	2.27	0.67
1:A:240[B]:THR:CG2	1:A:242[B]:GLY:N	2.59	0.64
1:A:56:TRP:CE2	2:B:2:L6S:H4	2.39	0.57
1:A:259:LYS:NZ	6:A:282:ACT:CH3	2.66	0.56
1:A:120:MET:HE2	7:A:5:HOH:O	2.07	0.54
1:A:240[B]:THR:HG22	1:A:242[B]:GLY:C	2.35	0.51
1:A:19:GLN:HE21	1:A:19:GLN:HA	1.78	0.49
1:A:83[B]:ARG:NH1	7:A:376:HOH:O	2.33	0.49
1:A:240[B]:THR:HG23	1:A:241[B]:ALA:C	2.38	0.48
1:A:240[B]:THR:HG23	1:A:241[B]:ALA:HB3	1.95	0.47
1:A:240[B]:THR:HG23	1:A:241[B]:ALA:CB	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83[B]:ARG:HD2	1:A:99:ASP:OD1	2.16	0.46
1:A:83[B]:ARG:CD	1:A:99:ASP:OD1	2.65	0.44
1:A:259:LYS:HZ2	6:A:282:ACT:CH3	2.31	0.44
1:A:50:TYR:O	1:A:103:ALA:HA	2.20	0.42
1:A:133:ARG:HG3	1:A:139:SER:HB3	2.02	0.42
1:A:274:ASP:C	7:A:545:HOH:O	2.62	0.41
1:A:245:ASN:HA	7:A:6:HOH:O	2.19	0.41
1:A:94:LYS:HD3	7:A:439:HOH:O	2.21	0.41
1:A:151:LYS:HD3	1:A:151:LYS:HA	1.96	0.41
1:A:19:GLN:HA	1:A:19:GLN:NE2	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:565:HOH:O	7:A:567:HOH:O[2_554]	1.74	0.46
7:A:558:HOH:O	7:A:561:HOH:O[4_555]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/268 (98%)	247 (94%)	13 (5%)	3 (1%)	11 3

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241[A]	ALA
1	A	241[B]	ALA
1	A	243	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/234 (99%)	224 (97%)	8 (3%)	32	20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	28	LYS
1	A	134	VAL
1	A	165	ASN
1	A	212	LYS
1	A	252[A]	GLU
1	A	252[B]	GLU
1	A	274	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	42	ASN
1	A	165	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLA	B	1	2	12,12,12	1.03	0	17,17,17	3.27	5 (29%)
2	L6S	B	2	2	15,15,16	1.05	2 (13%)	21,22,24	3.08	7 (33%)
2	GAL	B	3	2	11,11,12	1.38	1 (9%)	15,15,17	2.87	4 (26%)
2	L6S	B	4	2	15,15,16	1.88	2 (13%)	21,22,24	4.69	11 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLA	B	1	2	-	0/2/22/22	0/1/1/1
2	L6S	B	2	2	-	0/6/23/26	0/1/1/1
2	GAL	B	3	2	-	0/2/19/22	0/1/1/1
2	L6S	B	4	2	1/1/5/6	3/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	4	L6S	C2-C3	-5.01	1.44	1.52
2	B	4	L6S	O5-C1	4.18	1.50	1.43
2	B	3	GAL	O3-C3	3.45	1.51	1.43
2	B	2	L6S	O5-C1	-2.43	1.39	1.43
2	B	2	L6S	C1-C2	-2.22	1.46	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	L6S	C1-C2-C3	-14.26	88.88	109.64
2	B	1	GLA	C1-C2-C3	-9.02	91.97	110.36
2	B	4	L6S	C1-O5-C5	-8.65	100.60	112.19
2	B	2	L6S	O4-C4-C3	-8.54	90.24	110.38
2	B	3	GAL	O3-C3-C4	8.22	129.74	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	L6S	O2-C2-C3	-6.73	96.22	110.15
2	B	2	L6S	C1-O5-C5	-6.63	103.30	112.19
2	B	4	L6S	O4-C4-C5	-6.48	93.36	109.32
2	B	4	L6S	O4-C4-C3	5.83	124.11	110.38
2	B	1	GLA	O5-C1-C2	5.62	120.19	110.30
2	B	2	L6S	C1-C2-C3	-5.24	102.02	109.64
2	B	1	GLA	O3-C3-C4	-4.83	99.00	110.38
2	B	3	GAL	O3-C3-C2	-4.54	100.80	110.05
2	B	1	GLA	C1-O5-C5	-4.24	105.45	113.65
2	B	2	L6S	C2-C3-C4	4.21	118.26	110.86
2	B	4	L6S	C3-C4-C5	4.19	117.82	110.23
2	B	3	GAL	C1-O5-C5	-3.60	107.36	112.19
2	B	1	GLA	O1-C1-C2	3.44	118.98	108.98
2	B	4	L6S	O5-C5-C4	-3.37	102.62	110.83
2	B	4	L6S	O5-C5-C6	3.37	114.86	107.59
2	B	2	L6S	O2-C2-C1	2.92	115.91	109.22
2	B	3	GAL	C3-C4-C5	-2.75	105.24	110.23
2	B	4	L6S	C2-C3-C4	2.50	115.26	110.86
2	B	4	L6S	O6-S1-O2S	2.45	114.41	106.92
2	B	4	L6S	O6-C6-C5	2.43	111.90	107.57
2	B	2	L6S	O5-C5-C6	2.36	112.69	107.59
2	B	2	L6S	C6-C5-C4	-2.20	107.44	112.07

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	4	L6S	C1

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	L6S	C4-C5-C6-O6
2	B	4	L6S	O5-C5-C6-O6
2	B	4	L6S	C5-C6-O6-S1


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	L6S	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ACT	A	284	-	3,3,3	0.76	0	3,3,3	1.69	1 (33%)
6	ACT	A	285	-	3,3,3	0.96	0	3,3,3	1.53	0
6	ACT	A	282	-	3,3,3	0.97	0	3,3,3	2.38	2 (66%)
6	ACT	A	283	-	3,3,3	1.22	0	3,3,3	1.12	0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	282	ACT	OXT-C-CH3	2.99	127.60	115.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	282	ACT	OXT-C-O	-2.82	111.58	122.03
6	A	284	ACT	OXT-C-CH3	2.20	124.29	115.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	285	ACT	1	0
6	A	282	ACT	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/268 (95%)	-0.56	7 (2%) 56 56	2, 9, 23, 39	10 (3%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	ALA	16.1
1	A	240[A]	THR	7.5
1	A	241[A]	ALA	7.2
1	A	242[A]	GLY	6.8
1	A	18	ALA	4.6
1	A	244	ALA	3.2
1	A	19	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	L6S	B	4	15/16	0.79	0.16	33,39,57,58	0
2	GAL	B	3	11/12	0.93	0.08	10,13,20,26	0
2	L6S	B	2	15/16	0.95	0.06	4,6,9,10	0
2	GLA	B	1	12/12	0.97	0.05	2,5,10,14	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ACT	A	284	4/4	0.63	0.18	62,62,63,63	0
6	ACT	A	285	4/4	0.82	0.12	28,31,31,31	0
6	ACT	A	282	4/4	0.83	0.14	29,29,31,32	0
6	ACT	A	283	4/4	0.94	0.10	23,25,25,26	0
5	CL	A	280	1/1	0.98	0.16	25,25,25,25	0
3	CA	A	278	1/1	0.99	0.02	8,8,8,8	0
5	CL	A	281	1/1	0.99	0.03	8,8,8,8	0
4	MG	A	279	1/1	0.99	0.21	7,7,7,7	0

6.5 Other polymers ⓘ

There are no such residues in this entry.