



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:25 AM UTC

PDB ID : 2IMP / pdb_00002imp
Title : Crystal structure of lactaldehyde dehydrogenase from E. coli: the ternary complex with lactate (occupancy 0.5) and NADH. Crystals soaked with (L)-Lactate.
Authors : Di Costanzo, L.; Gomez, G.A.; Christianson, D.W.
Deposited on : 2006-10-04
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	509	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lactaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	477	3666	2323	631	695	17	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P25553
A	249	OCS	CYS	modified residue	UNP P25553

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (CCD ID: NAI) (formula: $C_{21}H_{29}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	88	42	14	28	4	0	1

- Molecule 4 is LACTIC ACID (CCD ID: LAC) (formula: $C_3H_6O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	221	Total 221	O 221	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lactaldehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	143.03Å 143.03Å 109.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.10) 88.3 (50.00-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.10Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.200 , 0.234 0.204 , 0.238	Depositor DCC
R_{free} test set	2412 reflections (6.67%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3991	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, OCS, NAI, LAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/3726	0.98	11/5048 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	456	GLY	N-CA-C	7.66	121.25	110.38
1	A	124	GLY	N-CA-C	-7.53	100.69	112.58
1	A	366	PRO	N-CA-C	6.51	116.72	110.47
1	A	7	HIS	N-CA-C	6.45	117.93	109.93
1	A	150	LEU	N-CA-C	6.42	118.74	110.39
1	A	329	ILE	N-CA-C	5.93	116.59	110.36
1	A	123	GLU	N-CA-C	5.74	119.07	109.95
1	A	127	ILE	N-CA-C	5.56	116.31	108.36
1	A	27	VAL	N-CA-C	5.33	117.23	108.97
1	A	291	VAL	N-CA-C	5.33	115.77	107.99
1	A	175	ILE	N-CA-C	5.11	115.94	108.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3666	0	3650	99	0
2	A	10	0	0	5	0
3	A	88	0	54	11	0
4	A	6	0	0	0	0
5	A	221	0	0	0	0
All	All	3991	0	3704	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:VAL:HB	1:A:5:VAL:N	1.80	0.96
1:A:305:LEU:HG	1:A:309:MET:HE2	1.47	0.95
1:A:309:MET:HE1	1:A:369:LEU:HD13	1.56	0.86
1:A:277:ARG:HD3	1:A:288:ALA:O	1.77	0.84
1:A:254:GLY:H	3:A:506[B]:NAI:H71N	1.33	0.76
1:A:279:ILE:O	1:A:280:ASN:HB2	1.86	0.74
1:A:254:GLY:N	3:A:506[B]:NAI:H71N	1.87	0.72
1:A:3:VAL:HB	1:A:4:PRO:C	2.15	0.70
1:A:278:VAL:HG21	1:A:324:ALA:HB1	1.73	0.69
1:A:276:SER:O	1:A:279:ILE:HG12	1.94	0.68
1:A:256:ALA:HA	2:A:509:SO4:O3	1.94	0.68
1:A:18:TRP:CG	1:A:39:PRO:HB3	2.29	0.67
1:A:278:VAL:CG2	1:A:324:ALA:HB1	2.24	0.67
1:A:411:LEU:N	2:A:509:SO4:O4	2.27	0.66
1:A:294:GLN:HB3	1:A:394:PHE:CE1	2.30	0.66
1:A:276:SER:OG	1:A:440:GLU:HB2	1.95	0.66
1:A:152:TRP:HH2	1:A:283:GLN:HB3	1.61	0.65
1:A:350:ARG:HH21	1:A:371:LEU:HD12	1.62	0.64
1:A:384:THR:O	1:A:386:GLY:N	2.33	0.61
1:A:281:SER:HB2	1:A:326:GLY:H	1.65	0.61
1:A:287:CYS:HA	1:A:441:ASN:HD22	1.63	0.61
1:A:374:ARG:H	1:A:377:MET:HE3	1.67	0.59
1:A:164:ALA:HB3	1:A:165:PRO:HD3	1.85	0.59
1:A:100:LEU:HD11	1:A:279:ILE:HA	1.85	0.57
1:A:333:ALA:O	1:A:337:VAL:HG23	2.03	0.57
1:A:279:ILE:O	1:A:280:ASN:CB	2.53	0.56
1:A:293:VAL:O	1:A:393:ALA:HA	2.06	0.56
1:A:100:LEU:HD13	1:A:279:ILE:C	2.31	0.56
1:A:262:ASP:OD1	1:A:294:GLN:HG3	2.06	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:SER:HB3	1:A:287:CYS:SG	2.45	0.56
1:A:384:THR:O	1:A:384:THR:HG22	2.07	0.54
1:A:230:SER:HA	3:A:506[B]:NAI:O3D	2.08	0.54
1:A:266:LEU:HD11	1:A:301:PHE:HA	1.88	0.54
1:A:4:PRO:HG2	1:A:37:ARG:NH2	2.23	0.54
1:A:91:VAL:HG13	1:A:317:PRO:HB2	1.89	0.53
1:A:3:VAL:N	1:A:4:PRO:HA	2.23	0.53
1:A:113:ASP:O	1:A:117:GLU:HG3	2.09	0.53
1:A:374:ARG:N	1:A:377:MET:HE3	2.24	0.52
1:A:411:LEU:HB3	2:A:509:SO4:O2	2.10	0.52
1:A:152:TRP:CH2	1:A:283:GLN:HB3	2.44	0.52
1:A:297:ILE:O	1:A:297:ILE:HG12	2.09	0.52
1:A:264:ALA:HB2	1:A:416:TYR:O	2.10	0.52
1:A:266:LEU:HG	1:A:304:ARG:HD2	1.91	0.52
1:A:305:LEU:HG	1:A:309:MET:CE	2.30	0.51
1:A:210:GLU:H	1:A:210:GLU:CD	2.18	0.51
1:A:277:ARG:HG3	1:A:389:LEU:HD21	1.92	0.51
1:A:168:LEU:C	1:A:168:LEU:HD13	2.37	0.49
1:A:254:GLY:C	3:A:506[B]:NAI:N7N	2.71	0.49
1:A:281:SER:OG	1:A:325:MET:HG3	2.13	0.48
1:A:336:ARG:NE	1:A:385:PHE:CZ	2.76	0.48
1:A:254:GLY:N	3:A:506[B]:NAI:N7N	2.57	0.48
1:A:27:VAL:CG1	1:A:32:GLU:HA	2.44	0.47
1:A:284:VAL:O	3:A:506[B]:NAI:H4N	2.14	0.47
1:A:267:GLU:HG2	1:A:304:ARG:NH1	2.30	0.47
1:A:268:LEU:HD23	1:A:268:LEU:C	2.40	0.47
1:A:271:LYS:HG3	1:A:275:ASP:OD2	2.14	0.47
1:A:283:GLN:NE2	1:A:328:LEU:HA	2.29	0.47
1:A:151:PRO:HA	3:A:506[B]:NAI:H52A	1.97	0.47
1:A:325:MET:HE1	1:A:387:PRO:HB3	1.95	0.47
1:A:341:VAL:O	1:A:345:VAL:HG23	2.15	0.47
1:A:294:GLN:HB3	1:A:394:PHE:CZ	2.50	0.46
1:A:82:ARG:NH2	1:A:194:GLU:OE2	2.43	0.46
1:A:255:LYS:N	2:A:509:SO4:O1	2.41	0.46
1:A:87:SER:O	1:A:91:VAL:HG23	2.16	0.46
1:A:263:ASP:O	1:A:418:GLN:HG3	2.16	0.46
1:A:329:ILE:HG23	1:A:330:ASN:N	2.30	0.46
1:A:154:PHE:N	1:A:155:PRO:CD	2.79	0.45
1:A:350:ARG:HD3	1:A:372:ASP:OD2	2.15	0.45
1:A:309:MET:CE	1:A:369:LEU:HD22	2.46	0.45
1:A:270:VAL:O	1:A:274:VAL:HG22	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:PRO:HA	3:A:506[A]:NAI:H52A	1.99	0.45
1:A:276:SER:O	1:A:287:CYS:SG	2.73	0.44
1:A:332:ALA:HB3	3:A:506[A]:NAI:H6N	1.99	0.44
1:A:285:CYS:HA	3:A:506[B]:NAI:C7N	2.47	0.44
1:A:82:ARG:NH2	1:A:194:GLU:OE1	2.51	0.44
1:A:342:ALA:O	1:A:346:GLU:HG3	2.18	0.44
1:A:254:GLY:HA3	2:A:509:SO4:O1	2.18	0.43
1:A:332:ALA:HB1	1:A:336:ARG:HH12	1.83	0.43
1:A:310:GLN:HG2	1:A:353:PHE:CZ	2.54	0.43
1:A:152:TRP:O	1:A:155:PRO:HD3	2.19	0.43
1:A:275:ASP:HA	1:A:278:VAL:HG12	1.99	0.43
1:A:380:MET:HE2	1:A:392:VAL:HG21	2.00	0.42
1:A:280:ASN:ND2	1:A:283:GLN:HB2	2.34	0.42
1:A:57:GLN:HB3	1:A:58:PRO:HD3	2.02	0.42
1:A:231:VAL:O	1:A:235:GLU:HG3	2.20	0.42
1:A:254:GLY:O	3:A:506[B]:NAI:N7N	2.53	0.42
1:A:282:GLY:O	1:A:385:PHE:O	2.38	0.41
1:A:273:ILE:HD12	1:A:301:PHE:HZ	1.85	0.41
1:A:310:GLN:HG2	1:A:353:PHE:CE1	2.55	0.41
1:A:325:MET:CE	1:A:387:PRO:HB3	2.51	0.41
1:A:328:LEU:HD12	1:A:334:LEU:HA	2.02	0.41
1:A:384:THR:C	1:A:385:PHE:CD1	2.99	0.41
1:A:27:VAL:HG12	1:A:28:ASN:N	2.35	0.41
1:A:325:MET:HE1	1:A:387:PRO:CB	2.51	0.41
1:A:124:GLY:C	1:A:125:GLU:HG3	2.46	0.40
1:A:338:GLU:HG2	1:A:357:ALA:HB2	2.03	0.40
1:A:396:THR:OG1	1:A:399:ASP:OD1	2.37	0.40
1:A:158:LEU:HD23	1:A:161:ARG:HD2	2.02	0.40
1:A:280:ASN:HB3	1:A:284:VAL:HG23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	474/479 (99%)	441 (93%)	27 (6%)	6 (1%)	9 6

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	ASN
1	A	385	PHE
1	A	265	ASP
1	A	386	GLY
1	A	41	GLY
1	A	279	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	375/377 (100%)	372 (99%)	3 (1%)	73 81

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	251	GLU
1	A	280	ASN
1	A	385	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	ASN
1	A	280	ASN
1	A	339	GLN
1	A	381	HIS
1	A	441	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	OCS	A	249	1	6,8,9	1.28	1 (16%)	7,11,13	1.78	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	249	1	-	1/4/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	249	OCS	OD3-SG	-2.25	1.38	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	OCS	OD2-SG-OD3	2.68	118.11	111.40
1	A	249	OCS	OD1-SG-CB	-2.37	103.22	106.76

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	249	OCS	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	509	-	4,4,4	0.35	0	6,6,6	0.20	0
3	NAI	A	506[B]	-	47,48,48	1.89	12 (25%)	64,73,73	2.15	18 (28%)
3	NAI	A	506[A]	-	47,48,48	1.78	9 (19%)	64,73,73	1.72	15 (23%)
2	SO4	A	510	-	4,4,4	0.38	0	6,6,6	0.08	0
4	LAC	A	508	-	4,5,5	0.77	0	2,6,6	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LAC	A	508	-	-	0/4/4/4	-
3	NAI	A	506[B]	-	-	10/29/72/72	0/5/5/5
3	NAI	A	506[A]	-	-	6/29/72/72	0/5/5/5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	506[B]	NAI	O7N-C7N	6.37	1.39	1.24
3	A	506[A]	NAI	O7N-C7N	6.36	1.39	1.24
3	A	506[B]	NAI	C2A-N3A	3.99	1.41	1.33
3	A	506[A]	NAI	C2A-N3A	3.90	1.40	1.33
3	A	506[A]	NAI	C2A-N1A	3.78	1.40	1.33
3	A	506[B]	NAI	C2A-N1A	3.70	1.40	1.33
3	A	506[B]	NAI	C4N-C3N	-3.29	1.43	1.50
3	A	506[A]	NAI	C4N-C3N	-3.22	1.43	1.50
3	A	506[A]	NAI	C8A-N7A	2.89	1.37	1.31
3	A	506[A]	NAI	C6N-C5N	2.85	1.41	1.33
3	A	506[B]	NAI	C8A-N7A	2.84	1.37	1.31
3	A	506[B]	NAI	C6N-C5N	2.80	1.41	1.33
3	A	506[B]	NAI	O5D-C5D	-2.73	1.34	1.44
3	A	506[A]	NAI	C2N-C3N	2.68	1.42	1.35
3	A	506[B]	NAI	C2N-C3N	2.63	1.42	1.35
3	A	506[B]	NAI	PN-O3	2.44	1.62	1.59
3	A	506[A]	NAI	C4N-C5N	-2.31	1.43	1.49
3	A	506[B]	NAI	C4N-C5N	-2.27	1.43	1.49
3	A	506[B]	NAI	PN-O5D	-2.26	1.50	1.59
3	A	506[A]	NAI	PN-O3	2.09	1.61	1.59
3	A	506[B]	NAI	PA-O3	2.01	1.61	1.59

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	506[B]	NAI	PN-O5D-C5D	7.18	162.47	121.35
3	A	506[B]	NAI	O5D-C5D-C4D	6.36	130.64	108.99
3	A	506[A]	NAI	N3A-C2A-N1A	-5.75	119.88	128.58
3	A	506[B]	NAI	N3A-C2A-N1A	-5.72	119.93	128.58
3	A	506[B]	NAI	C5A-C4A-N3A	-3.97	121.24	126.72
3	A	506[A]	NAI	C5A-C4A-N3A	-3.97	121.25	126.72
3	A	506[A]	NAI	N9A-C8A-N7A	-3.75	108.62	113.94
3	A	506[B]	NAI	N9A-C8A-N7A	-3.73	108.64	113.94
3	A	506[B]	NAI	C1D-N1N-C2N	-3.55	115.30	121.14
3	A	506[A]	NAI	C5A-N7A-C8A	3.30	108.63	103.45
3	A	506[A]	NAI	C2A-N3A-C4A	3.27	119.83	111.83
3	A	506[B]	NAI	C5A-N7A-C8A	3.27	108.58	103.45
3	A	506[A]	NAI	C3N-C7N-N7N	3.22	123.39	117.67
3	A	506[B]	NAI	C3N-C7N-N7N	3.22	123.39	117.67
3	A	506[B]	NAI	C2A-N3A-C4A	3.18	119.60	111.83
3	A	506[B]	NAI	O4D-C1D-N1N	2.89	113.59	108.08
3	A	506[B]	NAI	C5D-C4D-C3D	2.87	125.53	115.21
3	A	506[A]	NAI	O4D-C1D-N1N	2.84	113.50	108.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	506[A]	NAI	C5D-C4D-C3D	2.64	124.70	115.21
3	A	506[B]	NAI	C6N-N1N-C2N	2.58	122.08	119.32
3	A	506[B]	NAI	O1N-PN-O3	-2.55	100.39	107.27
3	A	506[A]	NAI	C4A-C5A-N7A	-2.50	107.72	110.58
3	A	506[A]	NAI	C6N-N1N-C2N	2.47	121.96	119.32
3	A	506[A]	NAI	C3N-C2N-N1N	-2.45	119.60	123.20
3	A	506[B]	NAI	O7N-C7N-N7N	-2.43	117.44	122.89
3	A	506[B]	NAI	C3N-C2N-N1N	-2.41	119.66	123.20
3	A	506[B]	NAI	C4A-C5A-N7A	-2.41	107.83	110.58
3	A	506[A]	NAI	O7N-C7N-N7N	-2.25	117.86	122.89
3	A	506[A]	NAI	C1D-N1N-C2N	-2.22	117.47	121.14
3	A	506[B]	NAI	N3A-C4A-N9A	2.15	130.82	127.17
3	A	506[A]	NAI	N3A-C4A-N9A	2.09	130.73	127.17
3	A	506[B]	NAI	O5D-PN-O2N	2.03	117.00	108.94
3	A	506[A]	NAI	C5A-C4A-N9A	2.03	108.02	105.81

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	506[A]	NAI	C5D-O5D-PN-O3
3	A	506[A]	NAI	C5D-O5D-PN-O1N
3	A	506[A]	NAI	O4D-C1D-N1N-C6N
3	A	506[B]	NAI	PN-O3-PA-O5B
3	A	506[B]	NAI	C5D-O5D-PN-O1N
3	A	506[B]	NAI	C4D-C5D-O5D-PN
3	A	506[B]	NAI	O4D-C4D-C5D-O5D
3	A	506[B]	NAI	C3D-C4D-C5D-O5D
3	A	506[A]	NAI	PN-O3-PA-O5B
3	A	506[A]	NAI	C5D-O5D-PN-O2N
3	A	506[B]	NAI	C5D-O5D-PN-O3
3	A	506[B]	NAI	C5D-O5D-PN-O2N
3	A	506[A]	NAI	C2N-C3N-C7N-N7N
3	A	506[B]	NAI	O4D-C1D-N1N-C2N
3	A	506[B]	NAI	C2N-C3N-C7N-N7N
3	A	506[B]	NAI	C2D-C1D-N1N-C2N


There are no ring outliers.

3 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	509	SO4	5	0
3	A	506[B]	NAI	9	0
3	A	506[A]	NAI	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/479 (99%)	0.55	68 (14%) 6 6	6, 21, 52, 100	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285	CYS	10.0
1	A	280	ASN	8.9
1	A	281	SER	8.5
1	A	279	ILE	8.0
1	A	3	VAL	7.6
1	A	286	ASN	7.0
1	A	385	PHE	6.8
1	A	282	GLY	6.2
1	A	278	VAL	6.0
1	A	284	VAL	6.0
1	A	283	GLN	5.7
1	A	384	THR	5.2
1	A	336	ARG	4.9
1	A	274	VAL	4.8
1	A	382	GLU	4.8
1	A	276	SER	4.7
1	A	479	SER	4.6
1	A	383	GLU	4.4
1	A	353	PHE	4.3
1	A	287	CYS	4.2
1	A	442	PHE	3.8
1	A	275	ASP	3.7
1	A	374	ARG	3.7
1	A	266	LEU	3.7
1	A	21	ASP	3.7
1	A	346	GLU	3.3
1	A	253	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	345	VAL	3.2
1	A	379	ILE	3.2
1	A	386	GLY	3.2
1	A	340	LYS	3.2
1	A	381	HIS	3.1
1	A	277	ARG	3.0
1	A	264	ALA	3.0
1	A	419	ASN	3.0
1	A	357	ALA	3.0
1	A	273	ILE	3.0
1	A	422	VAL	2.8
1	A	289	GLU	2.8
1	A	351	VAL	2.8
1	A	303	ASN	2.7
1	A	304	ARG	2.7
1	A	395	ASP	2.6
1	A	20	GLY	2.6
1	A	358	VAL	2.5
1	A	421	ASN	2.5
1	A	418	GLN	2.5
1	A	22	ALA	2.5
1	A	295	LYS	2.4
1	A	254	GLY	2.3
1	A	267	GLU	2.3
1	A	288	ALA	2.3
1	A	342	ALA	2.3
1	A	296	GLY	2.3
1	A	399	ASP	2.2
1	A	268	LEU	2.2
1	A	338	GLU	2.1
1	A	376	GLU	2.1
1	A	337	VAL	2.1
1	A	272	ALA	2.1
1	A	360	GLY	2.1
1	A	361	LYS	2.1
1	A	210	GLU	2.1
1	A	343	ARG	2.1
1	A	297	ILE	2.1
1	A	300	GLN	2.0
1	A	387	PRO	2.0
1	A	409	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OCS	A	249	9/10	0.96	0.08	13,14,23,24	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	510	5/5	0.82	0.14	71,71,72,72	0
2	SO4	A	509	5/5	0.86	0.44	46,50,51,51	0
4	LAC	A	508	6/6	0.86	0.26	42,43,43,44	6
3	NAI	A	506[B]	44/44	0.87	0.20	16,35,57,58	44
3	NAI	A	506[A]	44/44	0.87	0.20	11,29,47,48	44

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAI A 506 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.