



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:00 AM UTC

PDB ID : 5IME / pdb_00005ime
Title : Crystal structure of P21-activated kinase 1 (PAK1) in complex with compound 9
Authors : Li, D.; Wang, W.
Deposited on : 2016-03-06
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	 13% 75% 17% • 7%
1	B	300	 4% 84% 9% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase PAK 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2184	1383	366	419	16			
1	B	280	Total	C	N	O	S	0	0	0
			2195	1392	366	422	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	ASN	ASP	conflict	UNP Q13153
A	423	GLU	THR	conflict	UNP Q13153
A	546	GLY	-	expression tag	UNP Q13153
A	547	ASN	-	expression tag	UNP Q13153
A	548	SER	-	expression tag	UNP Q13153
B	389	ASN	ASP	conflict	UNP Q13153
B	423	GLU	THR	conflict	UNP Q13153
B	546	GLY	-	expression tag	UNP Q13153
B	547	ASN	-	expression tag	UNP Q13153
B	548	SER	-	expression tag	UNP Q13153

- Molecule 2 is 8-(3-aminopropyl)-6-[2-chloro-4-(3-methyl-2-oxopyrazin-1(2H)-yl)phenyl]-2-(methylamino)pyrido[2,3-d]pyrimidin-7(8H)-one (CCD ID: 6BZ) (formula: C₂₂H₂₂ClN₇O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 32	C 22	Cl 1	N 7	O 2	0	0
2	B	1	Total 32	C 22	Cl 1	N 7	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total	O	0	0
			71	71		
3	B	111	Total	O	0	0
			111	111		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase PAK 1



- Molecule 1: Serine/threonine-protein kinase PAK 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.48Å 82.82Å 65.89Å 90.00° 106.71° 90.00°	Depositor
Resolution (Å)	37.62 – 2.22 37.62 – 2.22	Depositor EDS
% Data completeness (in resolution range)	96.6 (37.62-2.22) 98.8 (37.62-2.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.224 , 0.263 0.226 , 0.265	Depositor DCC
R_{free} test set	1560 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 70.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4625	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6BZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2217	0.70	2/2995 (0.1%)
1	B	0.33	1/2229 (0.0%)	0.70	0/3010
All	All	0.31	1/4446 (0.0%)	0.70	2/6005 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	278	GLN	CD-OE1	5.24	1.33	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	531	THR	CA-C-N	-5.77	113.07	119.19
1	A	531	THR	C-N-CA	-5.77	113.07	119.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2184	0	2210	33	1
1	B	2195	0	2229	19	0
2	A	32	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	32	0	0	0	0
3	A	71	0	0	0	0
3	B	111	0	0	0	0
All	All	4625	0	4439	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:LYS:N	1:A:286:THR:HG1	1.79	0.80
1:A:357:THR:HG23	1:A:358:GLU:HG2	1.74	0.69
1:B:330:TYR:CD1	1:B:344:MET:HE1	2.30	0.67
1:A:275:LYS:HA	1:A:286:THR:H	1.63	0.62
1:B:342:VAL:HG12	1:B:344:MET:HE3	1.82	0.61
1:A:347:LEU:HD11	1:A:404:LYS:HD2	1.83	0.60
1:A:279:GLY:HA2	1:A:283:THR:HA	1.84	0.59
1:A:278:GLN:HB2	1:A:279:GLY:HA3	1.87	0.57
1:A:388:ARG:NH2	1:A:412:ALA:HB2	2.20	0.56
1:A:388:ARG:HD3	1:A:410:PHE:O	2.08	0.53
1:A:276:ILE:HG12	1:A:286:THR:HG23	1.92	0.51
1:B:298:ILE:HG12	1:B:343:VAL:HG22	1.92	0.51
1:A:302:ASN:HA	1:A:339:GLU:HG2	1.93	0.51
1:A:468:ASN:HB2	1:B:438:ARG:HD2	1.93	0.51
1:A:328:VAL:HG21	1:A:396:LEU:HD12	1.93	0.50
1:A:267:LYS:HD2	1:A:289:ASP:HA	1.93	0.50
1:A:330:TYR:HH	1:A:333:SER:HG	1.60	0.49
1:A:424:MET:HE3	1:A:436:VAL:HA	1.93	0.49
1:A:332:ASP:OD1	1:A:333:SER:N	2.43	0.48
1:B:265:ASP:HB3	1:B:268:LYS:HE2	1.95	0.48
1:A:301:MET:O	1:A:339:GLU:HA	2.14	0.48
1:B:493:ILE:HD11	1:B:519:GLN:HB2	1.95	0.48
1:B:275:LYS:HD2	1:B:278:GLN:NE2	2.29	0.47
1:A:462:PRO:HD2	1:A:465:LEU:HD22	1.96	0.47
1:A:336:VAL:HG22	1:A:341:TRP:CZ3	2.50	0.47
1:A:330:TYR:HA	1:A:344:MET:HG2	1.97	0.46
1:B:280:ALA:HA	1:B:281:SER:HB2	1.97	0.46
1:A:423:GLU:N	1:A:423:GLU:OE1	2.49	0.46
1:B:343:VAL:O	1:B:344:MET:HE2	2.16	0.45
1:A:326:ASN:HB3	1:A:376:ALA:HB2	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:393:ASP:OD1	1:A:393:ASP:N	2.50	0.45
1:B:279:GLY:HA3	1:B:280:ALA:HA	1.54	0.45
1:B:279:GLY:HA3	1:B:281:SER:HB2	1.99	0.44
1:A:454:ALA:HA	1:A:457:MET:HE3	2.00	0.44
1:A:261:VAL:HG11	1:A:335:LEU:HG	2.01	0.43
1:A:375:GLN:HG3	1:A:516:LEU:HD21	2.01	0.43
1:B:261:VAL:HG13	1:B:334:TYR:HA	2.00	0.43
1:B:330:TYR:HD1	1:B:344:MET:HE1	1.79	0.43
1:A:261:VAL:HG21	1:A:335:LEU:HD23	2.01	0.42
1:B:260:ILE:HD13	1:B:317:LEU:HD13	2.00	0.42
1:A:433:PRO:HD3	1:A:448:TRP:CE2	2.55	0.42
1:A:493:ILE:HD11	1:A:519:GLN:C	2.44	0.42
1:B:342:VAL:CG1	1:B:344:MET:HE3	2.50	0.42
1:B:454:ALA:HA	1:B:457:MET:HE3	2.01	0.42
1:B:505:ASP:HB3	1:B:508:LYS:HB3	2.01	0.42
1:A:281:SER:HA	1:A:282:GLY:HA2	1.87	0.42
1:A:336:VAL:HG22	1:A:341:TRP:HZ3	1.84	0.41
1:B:423:GLU:H	1:B:423:GLU:CD	2.28	0.41
1:A:275:LYS:N	1:A:286:THR:O	2.54	0.41
1:B:390:ILE:HB	1:B:449:SER:HB2	2.03	0.40
1:A:479:ASN:OD1	1:A:479:ASN:N	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:GLU:OE2	1:A:474:TYR:OH[2_645]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/300 (91%)	252 (93%)	18 (7%)	2 (1%)	18	18
1	B	274/300 (91%)	263 (96%)	10 (4%)	1 (0%)	30	33
All	All	546/600 (91%)	515 (94%)	28 (5%)	3 (0%)	24	26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	336	VAL
1	B	308	LYS
1	A	337	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/260 (93%)	238 (98%)	4 (2%)	53	68
1	B	243/260 (94%)	241 (99%)	2 (1%)	73	84
All	All	485/520 (93%)	479 (99%)	6 (1%)	63	76

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	253	ILE
1	A	378	GLU
1	A	393	ASP
1	A	479	ASN
1	B	311	LEU
1	B	332	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	314	ASN
1	A	499	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	302	ASN
1	B	326	ASN
1	B	389	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	6BZ	A	601	-	35,35,35	1.30	3 (8%)	39,50,50	2.64	16 (41%)
2	6BZ	B	601	-	35,35,35	1.28	4 (11%)	39,50,50	2.53	16 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6BZ	A	601	-	-	2/14/14/14	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6BZ	B	601	-	-	2/14/14/14	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	6BZ	C21-N31	5.50	1.38	1.34
2	B	601	6BZ	C21-N31	4.93	1.38	1.34
2	B	601	6BZ	C3-N5	-2.95	1.35	1.41
2	A	601	6BZ	C3-N5	-2.59	1.36	1.41
2	B	601	6BZ	C3-C2	-2.19	1.42	1.47
2	A	601	6BZ	C3-C2	-2.13	1.42	1.47
2	B	601	6BZ	C9-N5	-2.07	1.40	1.44

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	6BZ	C3-C2-N8	-8.28	120.00	123.75
2	A	601	6BZ	C3-C2-N8	-7.85	120.20	123.75
2	A	601	6BZ	C9-N5-C3	5.66	122.74	118.74
2	B	601	6BZ	C2-C3-N5	5.34	119.20	115.56
2	A	601	6BZ	C2-C3-N5	5.32	119.18	115.56
2	B	601	6BZ	C19-C18-C23	4.59	118.60	114.55
2	B	601	6BZ	O26-C25-C16	-4.58	119.38	125.37
2	A	601	6BZ	O26-C25-C16	-4.48	119.50	125.37
2	A	601	6BZ	C19-C18-C23	4.46	118.48	114.55
2	A	601	6BZ	N20-C21-N22	-4.33	122.23	126.42
2	B	601	6BZ	N20-C21-N22	-4.11	122.45	126.42
2	A	601	6BZ	C19-N20-C21	3.92	121.18	115.81
2	B	601	6BZ	C19-N20-C21	3.74	120.93	115.81
2	B	601	6BZ	C18-C17-C16	-3.08	119.08	122.47
2	A	601	6BZ	O4-C3-C2	-2.96	119.74	124.36
2	B	601	6BZ	C32-N31-C21	-2.95	120.03	123.59
2	A	601	6BZ	C18-C17-C16	-2.77	119.42	122.47
2	B	601	6BZ	C18-C23-N22	-2.63	118.45	122.95
2	A	601	6BZ	N31-C21-N22	2.61	119.47	116.96
2	A	601	6BZ	C18-C23-N22	-2.58	118.55	122.95
2	B	601	6BZ	C9-N5-C6	2.49	121.05	118.15
2	B	601	6BZ	C1-C2-N8	2.39	119.99	116.83
2	A	601	6BZ	C1-C2-C3	2.32	120.37	117.50
2	B	601	6BZ	O4-C3-C2	-2.29	120.79	124.36
2	A	601	6BZ	C32-N31-C21	-2.16	120.99	123.59
2	A	601	6BZ	C27-N24-C25	2.15	119.60	116.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	6BZ	N31-C21-N22	2.11	118.99	116.96
2	B	601	6BZ	C21-N22-C23	2.10	120.72	113.99
2	A	601	6BZ	C21-N22-C23	2.10	120.72	113.99
2	A	601	6BZ	C7-C6-N5	2.07	120.95	118.44
2	B	601	6BZ	C27-N24-C25	2.06	119.47	116.60
2	B	601	6BZ	C1-C2-C3	2.03	120.01	117.50

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	6BZ	C11-C12-C16-C17
2	A	601	6BZ	C11-C12-C16-C17
2	A	601	6BZ	C13-C12-C16-C17
2	B	601	6BZ	C13-C12-C16-C17

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/300 (93%)	0.95	38 (13%) 7 5	36, 80, 145, 212	0
1	B	280/300 (93%)	0.41	11 (3%) 43 40	30, 55, 126, 179	0
All	All	560/600 (93%)	0.68	49 (8%) 15 13	30, 66, 138, 212	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	GLY	3.9
1	A	334	TYR	3.8
1	A	261	VAL	3.5
1	A	382	SER	3.5
1	A	262	SER	3.4
1	A	283	THR	3.3
1	B	317	LEU	3.1
1	A	266	PRO	3.1
1	A	303	LEU	3.0
1	B	382	SER	2.9
1	B	315	GLU	2.9
1	A	325	PRO	2.8
1	A	330	TYR	2.8
1	A	335	LEU	2.7
1	A	336	VAL	2.6
1	A	298	ILE	2.6
1	B	253	ILE	2.6
1	A	412	ALA	2.5
1	A	302	ASN	2.5
1	B	541	THR	2.4
1	A	275	LYS	2.4
1	A	398	GLY	2.4
1	B	480	GLY	2.4
1	A	276	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	324	ASN	2.3
1	B	314	ASN	2.3
1	A	357	THR	2.3
1	A	422	SER	2.3
1	A	279	GLY	2.3
1	A	267	LYS	2.3
1	A	360	CYS	2.2
1	A	285	TYR	2.2
1	A	259	SER	2.2
1	A	326	ASN	2.2
1	B	319	MET	2.2
1	A	278	GLN	2.2
1	A	543	ASN	2.2
1	B	307	PRO	2.2
1	A	368	ALA	2.2
1	A	312	ILE	2.1
1	A	343	VAL	2.1
1	B	474	TYR	2.1
1	A	287	ALA	2.1
1	A	333	SER	2.1
1	A	313	ILE	2.1
1	A	383	ASN	2.1
1	A	527	LEU	2.0
1	A	296	VAL	2.0
1	B	478	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	6BZ	A	601	32/32	0.90	0.09	56,64,69,103	0
2	6BZ	B	601	32/32	0.93	0.08	45,51,61,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6BZ A 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 6BZ B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.