



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 10:56 AM UTC

PDB ID : 1INV / pdb_00001inv
Title : A SIALIC ACID DERIVED PHOSPHONATE ANALOG INHIBITS DIFFERENT STRAINS OF INFLUENZA VIRUS NEURAMINIDASE WITH DIFFERENT EFFICIENCIES
Authors : White, C.L.; Janakiraman, M.N.; Laver, W.G.; Philippon, C.; Vasella, A.; Air, G.M.; Luo, M.
Deposited on : 1994-09-26
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4157 atoms, of which 963 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INFLUENZA VIRUS B/LEE/40 NEURAMINIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	390	3736	1907	696	531	574	28	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	382	ARG	LYS	conflict	UNP P03474

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	28	8	14	1	5	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ca 2 2	0	0

- Molecule 4 is (1R)-4-acetamido-1,5-anhydro-2,4-dideoxy-1-phosphono-D-glycero-D-galactooctitol (CCD ID: EQP) (formula: $C_{10}H_{20}NO_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	P	0	0
			40	10	19	1	9	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	117	Total	H	O	0	0
			351	234	117		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INFLUENZA VIRUS B/LEE/40 NEURAMINIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	124.73Å 124.73Å 71.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.40 8.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-2.40) 71.8 (8.00-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.39Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.154 , (Not available) 0.157 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4157	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, EQP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	2/3114 (0.1%)	1.11	21/4210 (0.5%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	449	MET	SD-CE	-6.42	1.63	1.79
1	A	104	GLY	C-O	-6.09	1.20	1.24

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	225	GLN	N-CA-C	13.69	126.28	111.36
1	A	455	LEU	N-CA-C	9.17	122.97	111.69
1	A	102	ARG	N-CA-C	-7.64	103.94	113.18
1	A	405	GLU	CA-C-N	7.15	126.91	119.76
1	A	405	GLU	C-N-CA	7.15	126.91	119.76
1	A	441	ALA	N-CA-C	6.98	119.95	108.99
1	A	271	VAL	N-CA-C	6.57	120.58	113.43
1	A	282	ALA	N-CA-C	-6.33	105.38	113.23
1	A	295	SER	N-CA-C	6.28	122.54	114.56
1	A	293	ASP	N-CA-C	-5.82	97.83	108.02
1	A	272	GLU	N-CA-C	5.74	118.28	111.33
1	A	179	GLY	N-CA-C	5.67	119.58	111.12
1	A	211	THR	N-CA-C	5.47	120.44	113.55
1	A	300	ARG	CA-C-N	5.47	125.23	119.76
1	A	300	ARG	C-N-CA	5.47	125.23	119.76
1	A	462	VAL	N-CA-C	5.29	116.20	108.58
1	A	456	TRP	N-CA-C	5.20	117.28	109.23
1	A	171	ILE	N-CA-C	-5.19	101.00	108.48
1	A	290	ALA	N-CA-C	-5.11	100.19	108.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	298	ALA	N-CA-C	-5.02	106.84	113.17
1	A	229	CYS	N-CA-C	-5.00	103.33	110.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3040	696	2934	48	4
2	A	14	14	13	0	0
3	A	2	0	0	0	0
4	A	21	19	18	3	0
5	A	117	234	0	1	2
All	All	3194	963	2965	48	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:PRO:O	1:A:186:ARG:HD3	1.77	0.84
1:A:382:ARG:HH12	1:A:392:ASP:HB2	1.47	0.78
1:A:320:LYS:HG3	1:A:391:SER:O	1.87	0.74
1:A:348:ILE:HD12	1:A:374:ARG:HD2	1.73	0.69
1:A:93:GLN:HE22	1:A:455:LEU:H	1.47	0.62
1:A:270:ARG:HH11	1:A:340:ASN:HD21	1.51	0.58
1:A:200:ALA:HB3	1:A:222:LEU:HB3	1.87	0.56
1:A:321:THR:HB	1:A:394:LEU:HD22	1.87	0.56
1:A:116:ARG:HD2	1:A:428:GLU:OE2	2.06	0.55
1:A:409:TYR:OH	4:A:500:EQP:H2	2.07	0.54
1:A:128:ARG:HH12	1:A:171:ILE:HD13	1.72	0.54
1:A:411:PHE:CZ	1:A:426:GLY:HA3	2.43	0.54
1:A:263:LYS:HB2	1:A:263:LYS:NZ	2.24	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:ASN:HA	5:A:606:HOH:O	2.09	0.52
1:A:268:THR:HG22	1:A:269:GLY:N	2.25	0.52
1:A:103:PHE:O	1:A:110:SER:HB2	2.10	0.51
1:A:227:SER:HB3	1:A:349:LYS:HE2	1.92	0.51
1:A:270:ARG:HD3	1:A:316:LEU:HG	1.93	0.50
1:A:367:ARG:HD3	1:A:380:TYR:CE2	2.47	0.50
1:A:128:ARG:HG3	1:A:188:TRP:CZ2	2.48	0.49
1:A:327:ARG:NH2	1:A:367:ARG:HG3	2.28	0.49
1:A:125:LYS:O	1:A:126:GLU:HB2	2.13	0.48
1:A:116:ARG:HD3	1:A:430:VAL:CG2	2.44	0.48
1:A:116:ARG:HD3	1:A:430:VAL:HG21	1.95	0.47
1:A:374:ARG:NH2	4:A:500:EQP:O3P	2.48	0.47
1:A:78:PRO:HD2	1:A:186:ARG:HD2	1.95	0.46
1:A:315:ARG:HB2	1:A:388:TRP:CD1	2.50	0.46
1:A:367:ARG:HD3	1:A:380:TYR:HE2	1.80	0.45
1:A:116:ARG:NH2	4:A:500:EQP:O3P	2.44	0.44
1:A:326:PRO:HA	1:A:371:LYS:O	2.17	0.44
1:A:234:GLY:O	1:A:257:ARG:HG3	2.18	0.44
1:A:232:ILE:HG23	1:A:232:ILE:O	2.17	0.44
1:A:187:GLU:HB3	1:A:206:TYR:CZ	2.53	0.43
1:A:202:VAL:HB	1:A:214:TYR:HB3	2.01	0.43
1:A:255:LYS:HB3	1:A:263:LYS:HB3	2.01	0.43
1:A:270:ARG:NH1	1:A:340:ASN:HD21	2.15	0.43
1:A:98:ILE:HG22	1:A:100:PRO:HD3	2.00	0.42
1:A:381:VAL:HG22	1:A:382:ARG:N	2.34	0.42
1:A:130:PHE:CD2	1:A:158:SER:HB3	2.54	0.42
1:A:193:VAL:HG22	1:A:202:VAL:HG22	2.02	0.42
1:A:159:VAL:HG11	1:A:165:PRO:HA	2.01	0.41
1:A:361:ILE:HG22	1:A:384:ASP:OD1	2.20	0.41
1:A:202:VAL:O	1:A:213:THR:HA	2.21	0.41
1:A:268:THR:HG22	1:A:269:GLY:H	1.86	0.41
1:A:315:ARG:HB2	1:A:388:TRP:CG	2.56	0.41
1:A:202:VAL:HG23	1:A:222:LEU:HD21	2.02	0.41
1:A:288:GLU:HA	1:A:303:VAL:O	2.21	0.40
1:A:248:ILE:HA	1:A:272:GLU:O	2.22	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:TYR:HH	1:A:463:ASP:H[4_555]	1.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:ALA:H	5:A:508:HOH:H2[4_555]	1.16	0.44
1:A:453:GLN:H	5:A:551:HOH:H2[3_545]	1.23	0.37
1:A:97:LEU:H	1:A:155:HIS:HE2[3_545]	1.35	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/390 (100%)	370 (95%)	17 (4%)	1 (0%)	36	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	126	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/326 (100%)	304 (93%)	22 (7%)	15	26

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	GLU
1	A	84	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	128	ARG
1	A	132	LEU
1	A	147	ARG
1	A	153	LEU
1	A	186	ARG
1	A	229	CYS
1	A	263	LYS
1	A	266	LEU
1	A	270	ARG
1	A	285	LYS
1	A	292	ARG
1	A	367	ARG
1	A	374	ARG
1	A	379	LEU
1	A	394	LEU
1	A	403	ILE
1	A	419	LYS
1	A	453	GLN
1	A	455	LEU
1	A	456	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	144	ASN
1	A	169	ASN
1	A	215	HIS
1	A	230	ASN
1	A	340	ASN
1	A	453	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	467(A)	1	14,14,15	1.71	3 (21%)	17,19,21	1.42	3 (17%)
4	EQP	A	500	-	18,21,21	1.85	3 (16%)	22,31,31	1.60	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	467(A)	1	-	0/6/23/26	0/1/1/1
4	EQP	A	500	-	-	2/14/36/36	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	500	EQP	C4-C5	-4.21	1.49	1.53
4	A	500	EQP	P1-O3P	-3.78	1.48	1.54
2	A	467(A)	NAG	C4-C5	3.52	1.60	1.53
4	A	500	EQP	P1-O2P	-3.46	1.49	1.54
2	A	467(A)	NAG	C4-C3	2.86	1.59	1.52
2	A	467(A)	NAG	C3-C2	2.23	1.57	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	467(A)	NAG	C1-O5-C5	4.44	118.14	112.19
4	A	500	EQP	O3P-P1-O2P	4.20	119.00	107.58
4	A	500	EQP	O3P-P1-O1P	-3.28	105.21	113.45
4	A	500	EQP	O6-C2-C3	-2.90	106.77	110.66
2	A	467(A)	NAG	O7-C7-C8	-2.20	118.13	122.05
2	A	467(A)	NAG	C8-C7-N2	2.06	119.53	116.12

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	500	EQP	C5-C6-C7-O7
4	A	500	EQP	O6-C6-C7-O7

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	500	EQP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/390 (100%)	-0.85	0 100 100	1, 9, 18, 25	45 (11%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	467(A)	14/15	0.88	0.09	0,0,37,56	13
3	CA	A	469	1/1	0.90	0.03	30,30,30,30	1
4	EQP	A	500	21/21	0.96	0.07	0,4,32,68	0
3	CA	A	468	1/1	0.98	0.11	2,2,2,2	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.