



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:36 AM UTC

PDB ID : 2INV / pdb\_00002inv  
Title : Crystal structure of Inulin fructotransferase in the presence of di-fructose  
Authors : Rhee, S.; Jung, W.S.  
Deposited on : 2006-10-09  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	
1	B	410	
1	C	410	
2	D	2	
2	E	2	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	2	 50% 50%

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 9648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inulin fructotransferase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	399	2982	1859	535	578	6	4	0	0	0
1	B	399	2982	1859	535	578	6	4	0	0	0
1	C	399	2982	1859	535	578	6	4	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	D	2	23	12	11	0	0	0
2	E	2	23	12	11	0	0	0
2	F	2	23	12	11	0	0	0

- Molecule 3 is PHOSPHONATE (CCD ID: 2PO) (formula: HO<sub>3</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 4 3 1	0	0
3	B	1	Total O P 4 3 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	203	Total O 203 203	0	0
4	B	210	Total O 210 210	0	0
4	C	212	Total O 212 212	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inulin fructotransferase




- Molecule 1: Inulin fructotransferase



- Molecule 1: Inulin fructotransferase

Chain C:  63% 28% 5% . . .



- Molecule 2: beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain D:  50% 50%

FRU1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain E:  100%

FRU1  
FRU2

- Molecule 2: beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain F:  50% 50%

FRU1  
FRU2

## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	159.17Å 91.90Å 92.92Å 90.00° 124.82° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	82.4 (50.00-1.80) 98.1 (50.00-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.44 (at 1.81Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.207 , 0.227 0.214 , 0.234	Depositor DCC
$R_{free}$ test set	10154 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.2	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 29.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for $1/2^*h+1/2^*k+2^*l, 1/2^*h+1/2^*k, -1/2^*h+1/2^*k-l$ 0.012 for $-1/2^*h-3/2^*k-l, -1/2^*h+1/2^*k-l, 1/2^*h+1/2^*k$ 0.013 for $-1/2^*h+3/2^*k-l, 1/2^*h+1/2^*k+l, 1/2^*h-1/2^*k$ 0.019 for $1/2^*h-1/2^*k+2^*l, -1/2^*h+1/2^*k, -1/2^*h-1/2^*k-l$ 0.019 for $-h+k-l, -l, -k$ 0.005 for $-h-k-l, l, k$ 0.019 for $-1/2^*h+1/2^*k+l, 1/2^*h-1/2^*k+l, 1/2^*h+1/2^*k$ 0.005 for $-1/2^*h-1/2^*k+l, -1/2^*h-1/2^*k-l, 1/2^*h-1/2^*k$ 0.487 for $1/2^*h+3/2^*k, 1/2^*h-1/2^*k, -1/2^*h-1/2^*k-l$ 0.487 for $1/2^*h-3/2^*k, -1/2^*h-1/2^*k, -1/2^*h+1/2^*k-l$ 0.013 for $-h-2^*l, -k, l$	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9648	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, 2PO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.69	9/3042 (0.3%)	1.49	49/4143 (1.2%)
1	B	0.66	5/3042 (0.2%)	1.51	51/4143 (1.2%)
1	C	0.64	5/3042 (0.2%)	1.76	53/4143 (1.3%)
All	All	0.66	19/9126 (0.2%)	1.59	153/12429 (1.2%)

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	291	PRO	C-N	19.16	1.60	1.33
1	B	291	PRO	C-N	18.08	1.58	1.33
1	C	291	PRO	C-N	16.81	1.56	1.33
1	B	285	THR	CB-OG1	8.87	1.57	1.43
1	A	285	THR	CB-OG1	8.78	1.57	1.43

The worst 5 of 153 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	172	ASP	CA-C-N	47.90	179.72	119.84
1	C	172	ASP	C-N-CA	47.90	179.72	119.84
1	B	291	PRO	O-C-N	-34.65	75.86	122.64
1	C	291	PRO	O-C-N	-33.80	77.02	122.64
1	A	291	PRO	O-C-N	-33.23	77.78	122.64

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2982	0	2905	109	0
1	B	2982	0	2905	106	0
1	C	2982	0	2905	102	0
2	D	23	0	21	3	0
2	E	23	0	21	3	0
2	F	23	0	21	3	0
3	A	4	0	1	0	0
3	B	4	0	1	0	0
4	A	203	0	0	1	0
4	B	210	0	0	0	0
4	C	212	0	0	2	0
All	All	9648	0	8780	272	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 272 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:265:MSE:HE3	1:B:273:THR:HG22	1.32	1.10
1:A:319:MSE:HE1	1:A:338:ARG:HB2	1.39	1.01
1:C:291:PRO:HB2	2:F:2:FRU:O6	1.67	0.93
1:A:291:PRO:HB2	2:D:2:FRU:O6	1.70	0.90
1:B:291:PRO:HB2	2:E:2:FRU:O6	1.73	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/410 (97%)	369 (93%)	25 (6%)	3 (1%)	16	6
1	B	397/410 (97%)	369 (93%)	25 (6%)	3 (1%)	16	6
1	C	397/410 (97%)	368 (93%)	26 (6%)	3 (1%)	16	6
All	All	1191/1230 (97%)	1106 (93%)	76 (6%)	9 (1%)	16	6

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	173	PRO
1	A	199	ASN
1	B	199	ASN
1	C	199	ASN
1	B	257	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/325 (99%)	299 (93%)	23 (7%)	13	4
1	B	322/325 (99%)	298 (92%)	24 (8%)	12	4
1	C	322/325 (99%)	297 (92%)	25 (8%)	11	3
All	All	966/975 (99%)	894 (92%)	72 (8%)	12	4

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	256	GLN
1	C	450	PRO
1	C	291	PRO
1	C	349	ARG
1	B	67	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	324	ASN
1	C	89	GLN
1	C	382	ASN
1	B	335	HIS
1	B	416	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FRU	D	1	2	11,12,12	3.93	3 (27%)	10,18,18	2.44	2 (20%)
2	FRU	D	2	2	11,11,12	1.13	1 (9%)	15,15,18	2.51	4 (26%)
2	FRU	E	1	2	11,12,12	3.32	3 (27%)	10,18,18	2.90	3 (30%)
2	FRU	E	2	2	11,11,12	1.13	1 (9%)	15,15,18	2.51	4 (26%)
2	FRU	F	1	2	11,12,12	3.12	3 (27%)	10,18,18	1.98	2 (20%)
2	FRU	F	2	2	11,11,12	1.14	1 (9%)	15,15,18	2.52	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	D	1	2	-	0/5/24/24	0/1/1/1
2	FRU	D	2	2	-	1/4/20/24	0/1/1/1
2	FRU	E	1	2	-	2/5/24/24	0/1/1/1
2	FRU	E	2	2	-	1/4/20/24	0/1/1/1
2	FRU	F	1	2	-	0/5/24/24	0/1/1/1
2	FRU	F	2	2	-	1/4/20/24	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	FRU	O1-C1	-9.18	1.13	1.42
2	D	1	FRU	O1-C1	-9.18	1.13	1.42
2	E	1	FRU	O1-C1	-9.18	1.13	1.42
2	D	1	FRU	O6-C6	-8.34	1.07	1.42
2	E	1	FRU	O6-C6	-4.32	1.24	1.42

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	FRU	O6-C6-C5	7.82	137.96	111.33
2	E	2	FRU	O3-C3-C2	7.54	132.75	111.08
2	D	2	FRU	O3-C3-C2	7.54	132.75	111.08
2	F	2	FRU	O3-C3-C2	7.54	132.73	111.08
2	D	1	FRU	O6-C6-C5	6.21	132.49	111.33

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1	FRU	C4-C5-C6-O6
2	E	1	FRU	O5-C5-C6-O6
2	D	2	FRU	C4-C5-C6-O6
2	E	2	FRU	C4-C5-C6-O6
2	F	2	FRU	C4-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	2	FRU	3	0



*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	FRU	3	0
2	E	2	FRU	3	0
2	E	1	FRU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	2PO	A	1002	-	0,3,3	-	-	0,3,3	-	-
3	2PO	B	1001	-	0,3,3	-	-	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	395/410 (96%)	-1.34	0 100 100	10, 14, 25, 30	0
1	B	395/410 (96%)	-1.34	0 100 100	10, 14, 25, 30	0
1	C	395/410 (96%)	-1.34	0 100 100	10, 14, 25, 30	0
All	All	1185/1230 (96%)	-1.34	0 100 100	10, 14, 25, 30	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FRU	D	1	12/12	0.98	0.04	27,29,30,32	0
2	FRU	F	1	12/12	0.98	0.04	26,29,30,31	0
2	FRU	E	1	12/12	0.99	0.04	28,29,30,32	0
2	FRU	E	2	11/12	0.99	0.04	24,25,26,26	0
2	FRU	D	2	11/12	0.99	0.03	22,23,25,25	0
2	FRU	F	2	11/12	0.99	0.04	22,24,25,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain D:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	2PO	A	1002	4/4	0.99	0.04	45,45,46,47	0
3	2PO	B	1001	4/4	0.99	0.05	33,33,33,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.