



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:19 PM UTC

PDB ID : 4IN7 / pdb_00004in7
Title : (M)L214N mutant of the Rhodobacter sphaeroides Reaction Center
Authors : Saer, R.G.; Hardjasa, A.; Murphy, M.E.P.; Beatty, J.T.
Deposited on : 2013-01-04
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	266	 3% 74% 13% 10%
2	L	282	 87% 12%
3	M	307	 2% 79% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	HTO	L	309	-	X	-	-

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 7359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	240	Total	C	N	O	S	0	5	0
			1849	1183	320	337	9			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-5	HIS	-	expression tag	UNP P0C0Y7
H	-4	HIS	-	expression tag	UNP P0C0Y7
H	-3	HIS	-	expression tag	UNP P0C0Y7
H	-2	HIS	-	expression tag	UNP P0C0Y7
H	-1	HIS	-	expression tag	UNP P0C0Y7
H	0	HIS	-	expression tag	UNP P0C0Y7
H	251	VAL	-	expression tag	UNP P0C0Y7
H	252	VAL	-	expression tag	UNP P0C0Y7
H	253	ALA	-	expression tag	UNP P0C0Y7
H	254	ALA	-	expression tag	UNP P0C0Y7
H	255	MET	-	expression tag	UNP P0C0Y7
H	256	LEU	-	expression tag	UNP P0C0Y7
H	257	ALA	-	expression tag	UNP P0C0Y7
H	258	GLU	-	expression tag	UNP P0C0Y7
H	259	TYR	-	expression tag	UNP P0C0Y7
H	260	ALA	-	expression tag	UNP P0C0Y7

- Molecule 2 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	281	Total	C	N	O	S	0	2	0
			2239	1513	355	363	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	0	MET	-	expression tag	UNP P0C0Y8


- Molecule 3 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	302	Total	C	N	O	S	0	1	0
			2410	1605	396	399	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	0	MET	-	expression tag	UNP P0C0Y9
M	214	ASN	LEU	engineered mutation	UNP P0C0Y9
M	303	MET	-	expression tag	UNP P0C0Y9
M	304	ALA	-	expression tag	UNP P0C0Y9
M	305	PRO	-	expression tag	UNP P0C0Y9
M	306	LEU	-	expression tag	UNP P0C0Y9

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	K	0	0
			1	1		

- Molecule 6 is NONADEC-10-ENOIC ACID 2-[3,4-DIHYDROXY-6-HYDROXYMETHYL-5-(3,4,5-TRIHYDROXY-6-HYDROXYMETHYL-TETRAHYDRO-PYRAN-2-YLOXY)-TETRAHYDRO-PYRAN-2-YLOXY] -1-OCTADEC-9-ENOYLOXYMETHYL-ETHYL ESTER (CCD ID: GGD) (formula: C₅₂H₉₄O₁₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	H	1	Total	C	O	0	0
			57	42	15		

- Molecule 7 is BACTERIOCHLOROPHYLL A (CCD ID: BCL) (formula: C₅₅H₇₄MgN₄O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0

- Molecule 8 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	1	Total	C	N	O	0	0
			16	14	1	1		
8	L	1	Total	C	N	O	0	0
			16	14	1	1		
8	L	1	Total	C	N	O	0	0
			16	14	1	1		
8	M	1	Total	C	N	O	0	0
			16	14	1	1		
8	M	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 9 is UBIQUINONE-10 (CCD ID: U10) (formula: $C_{59}H_{90}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	L	1	Total	C	O	0	1
			46	38	8		
9	M	1	Total	C	O	0	0
			48	44	4		

- Molecule 10 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	L	1	Total	O	P	0	0
			5	4	1		

- Molecule 11 is HEPTANE-1,2,3-TRIOL (CCD ID: HTO) (formula: $C_7H_{16}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	L	1	Total	C	O	0	0
			10	7	3		
11	L	1	Total	C	O	0	0
			10	7	3		

- Molecule 12 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	M	1	Total	Fe		0	0
			1	1			

- Molecule 13 is BACTERIOPHEOPHYTIN A (CCD ID: BPH) (formula: C₅₅H₇₆N₄O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	M	1	Total	C	N	O	0	0
			65	55	4	6		
13	M	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 14 is SPHEROIDENE (CCD ID: SPO) (formula: C₄₁H₆₀O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	M	1	Total	C	O	0	0
			42	41	1		

- Molecule 15 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	M	1	Total	C	O	P	0	0
			81	62	17	2		

- Molecule 16 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
16	M	1	Total	C	N	O	P	0	0
			43	33	1	8	1		


- Molecule 17 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	M	1	Total	Mg	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reaction center protein H chain



- Molecule 2: Reaction center protein L chain



- Molecule 3: Reaction center protein M chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.14Å 139.14Å 185.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.57 – 2.85 38.57 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.57-2.85) 99.8 (38.57-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.86Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.213 0.179 , 0.211	Depositor DCC
R_{free} test set	2474 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7359	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CDL, FE, SPO, BCL, HTO, BPH, PO4, LDA, U10, MG, GOL, GGD, PC1, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	1.20	4/1929 (0.2%)	1.24	6/2619 (0.2%)
2	L	1.10	1/2339 (0.0%)	1.14	10/3203 (0.3%)
3	M	1.12	3/2507 (0.1%)	1.14	12/3422 (0.4%)
All	All	1.14	8/6775 (0.1%)	1.17	28/9244 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	M	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	85[A]	ILE	C-O	-7.12	1.15	1.24
1	H	85[B]	ILE	C-O	-7.12	1.15	1.24
1	H	100	PRO	C-O	-6.79	1.15	1.23
3	M	301	HIS	N-CA	6.79	1.54	1.46
3	M	259	ASN	C-O	-6.34	1.17	1.23

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	28	ASN	N-CA-C	7.43	122.38	113.16
3	M	175	VAL	CA-C-N	7.31	127.89	120.14
3	M	175	VAL	C-N-CA	7.31	127.89	120.14
2	L	220	VAL	CB-CA-C	-7.26	103.51	110.70
3	M	44[A]	ASN	N-CA-C	6.86	120.17	110.23

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	M	300	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1849	0	1873	29	0
2	L	2239	0	2185	24	0
3	M	2410	0	2314	31	0
4	H	30	0	40	0	0
4	L	12	0	16	1	0
5	H	1	0	0	0	0
6	H	57	0	68	3	0
7	L	132	0	148	12	0
7	M	132	0	148	13	0
8	L	48	0	93	0	0
8	M	32	0	62	4	0
9	L	46	0	46	4	0
9	M	48	0	63	2	0
10	L	5	0	0	1	0
11	L	20	0	32	0	0
12	M	1	0	0	0	0
13	M	130	0	150	17	0
14	M	42	0	60	3	0
15	M	81	0	102	1	0
16	M	43	0	60	1	0
17	M	1	0	0	0	0
All	All	7359	0	7460	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:M:412:BPH:HBB3	13:M:412:BPH:HHC	1.50	0.91
3:M:16:ALA:HB1	3:M:32:VAL:HG11	1.63	0.78
13:M:406:BPH:HHC	13:M:406:BPH:HBB3	1.66	0.78
1:H:248:ARG:HA	1:H:249[A]:LYS:HB2	1.68	0.75
1:H:220[B]:LYS:HE2	1:H:221:SER:OG	1.91	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	244/266 (92%)	231 (95%)	11 (4%)	2 (1%)	16	31
2	L	281/282 (100%)	265 (94%)	16 (6%)	0	100	100
3	M	301/307 (98%)	288 (96%)	11 (4%)	2 (1%)	18	35
All	All	826/855 (97%)	784 (95%)	38 (5%)	4 (0%)	30	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	249[A]	LYS
1	H	249[B]	LYS
3	M	30	SER
3	M	301	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	200/214 (94%)	189 (94%)	11 (6%)	19	41
2	L	221/221 (100%)	212 (96%)	9 (4%)	27	52
3	M	237/240 (99%)	225 (95%)	12 (5%)	21	43
All	All	658/675 (98%)	626 (95%)	32 (5%)	24	45

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	M	136	ARG
3	M	170	SER
2	L	158	SER
2	L	21	LEU
3	M	188	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
3	M	193	HIS
3	M	202	HIS
3	M	300	ASN
2	L	159	ASN
2	L	258	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 3 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
11	HTO	L	308	-	9,9,9	1.39	2 (22%)	10,10,10	1.24	1 (10%)
7	BCL	M	401	-	69,74,74	1.76	14 (20%)	79,115,115	2.02	23 (29%)
9	U10	M	407	-	48,48,63	1.38	5 (10%)	60,61,79	2.15	15 (25%)
4	GOL	H	301	-	5,5,5	0.80	0	5,5,5	1.15	0
4	GOL	H	306	-	5,5,5	1.18	0	5,5,5	1.06	1 (20%)
4	GOL	H	302	-	5,5,5	0.51	0	5,5,5	0.69	0
4	GOL	L	311	-	5,5,5	0.37	0	5,5,5	0.37	0
15	CDL	M	409	-	79,79,99	1.54	5 (6%)	84,90,111	1.45	9 (10%)
7	BCL	L	306	-	69,74,74	1.78	14 (20%)	79,115,115	1.93	22 (27%)
8	LDA	M	403	-	13,15,15	2.16	1 (7%)	14,17,17	0.92	1 (7%)
13	BPH	M	406	-	59,70,70	2.35	15 (25%)	59,101,101	2.14	13 (22%)
14	SPO	M	408	-	41,41,41	0.81	1 (2%)	47,50,50	1.87	11 (23%)
4	GOL	L	310	-	5,5,5	0.48	0	5,5,5	0.52	0
8	LDA	L	304	-	13,15,15	2.02	1 (7%)	14,17,17	0.78	0
9	U10	L	305[B]	-	23,23,63	2.12	2 (8%)	30,31,79	1.46	5 (16%)
11	HTO	L	309	-	9,9,9	1.61	3 (33%)	10,10,10	2.00	4 (40%)
16	PC1	M	410	-	42,42,53	1.52	3 (7%)	48,50,61	1.51	6 (12%)
6	GGD	H	307	-	58,58,68	1.15	3 (5%)	72,72,82	1.71	14 (19%)
13	BPH	M	412	-	59,70,70	2.45	16 (27%)	59,101,101	2.70	24 (40%)
7	BCL	L	301	-	69,74,74	1.66	17 (24%)	79,115,115	2.31	30 (37%)
10	PO4	L	307	-	4,4,4	1.22	1 (25%)	6,6,6	0.85	0
8	LDA	L	303	-	13,15,15	2.12	1 (7%)	14,17,17	0.79	1 (7%)
9	U10	L	305[A]	-	23,23,63	1.85	2 (8%)	30,31,79	1.56	6 (20%)
4	GOL	H	304	-	5,5,5	0.82	0	5,5,5	0.79	0
4	GOL	H	303	-	5,5,5	0.58	0	5,5,5	0.68	0
8	LDA	L	302	-	13,15,15	2.08	2 (15%)	14,17,17	0.86	0
8	LDA	M	404	-	13,15,15	1.98	1 (7%)	14,17,17	0.56	0
7	BCL	M	402	-	69,74,74	1.72	17 (24%)	79,115,115	2.03	20 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	HTO	L	308	-	-	6/10/10/10	-
7	BCL	M	401	-	-	11/41/137/137	-
9	U10	M	407	-	-	9/45/69/87	0/1/1/1
4	GOL	H	301	-	-	2/4/4/4	-
4	GOL	H	306	-	-	2/4/4/4	-
4	GOL	H	302	-	-	2/4/4/4	-
4	GOL	L	311	-	-	4/4/4/4	-
15	CDL	M	409	-	-	48/88/88/110	-
7	BCL	L	306	-	-	3/41/137/137	-
8	LDA	M	403	-	-	7/13/13/13	-
13	BPH	M	406	-	-	12/37/105/105	0/5/6/6
14	SPO	M	408	-	-	4/47/47/47	-
4	GOL	L	310	-	-	4/4/4/4	-
8	LDA	L	304	-	-	8/13/13/13	-
9	U10	L	305[B]	-	-	4/15/39/87	0/1/1/1
11	HTO	L	309	-	-	5/10/10/10	-
16	PC1	M	410	-	-	23/46/46/57	-
6	GGD	H	307	-	-	20/47/87/97	0/2/2/2
13	BPH	M	412	-	-	9/37/105/105	0/5/6/6
7	BCL	L	301	-	-	5/41/137/137	-
8	LDA	L	303	-	-	7/13/13/13	-
9	U10	L	305[A]	-	-	1/15/39/87	0/1/1/1
4	GOL	H	304	-	-	2/4/4/4	-
4	GOL	H	303	-	-	2/4/4/4	-
8	LDA	L	302	-	-	5/13/13/13	-
8	LDA	M	404	-	-	7/13/13/13	-
7	BCL	M	402	-	-	2/41/137/137	-

The worst 5 of 126 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	L	305[B]	U10	C6-C1	8.63	1.50	1.35
13	M	412	BPH	C1D-C2D	7.83	1.48	1.39
9	L	305[A]	U10	C6-C1	7.81	1.49	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	M	412	BPH	C1B-C2B	7.30	1.47	1.39
8	L	303	LDA	O1-N1	-7.10	1.24	1.42

The worst 5 of 206 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	M	407	U10	C30-C29-C31	9.78	132.20	115.23
13	M	412	BPH	O1D-CGD-CBD	8.90	138.23	124.72
7	L	301	BCL	O2D-CGD-CBD	7.80	124.87	111.23
14	M	408	SPO	C4-C5-C6	-7.35	114.60	124.91
15	M	409	CDL	OB6-CB5-C51	6.36	125.23	111.48

There are no chirality outliers.

5 of 214 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	306	GOL	O1-C1-C2-O2
4	H	306	GOL	O1-C1-C2-C3
4	H	302	GOL	O1-C1-C2-O2
4	H	302	GOL	O1-C1-C2-C3
4	H	303	GOL	C1-C2-C3-O3

There are no ring outliers.

16 monomers are involved in 52 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	M	401	BCL	6	0
9	M	407	U10	2	0
4	L	311	GOL	1	0
15	M	409	CDL	1	0
7	L	306	BCL	6	0
8	M	403	LDA	2	0
13	M	406	BPH	5	0
14	M	408	SPO	3	0
9	L	305[B]	U10	4	0
16	M	410	PC1	1	0
6	H	307	GGD	3	0
13	M	412	BPH	12	0
7	L	301	BCL	7	0
10	L	307	PO4	1	0
8	M	404	LDA	2	0



Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	M	402	BCL	9	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











Ligand U10 L 305 (B)



Ligand PC1 M 410







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	H	240/266 (90%)	-0.36	8 (3%)	49	40	22, 43, 61, 133	8 (3%)
2	L	281/282 (99%)	-0.41	0	100	100	30, 41, 64, 89	2 (0%)
3	M	302/307 (98%)	-0.36	5 (1%)	69	63	28, 44, 65, 99	7 (2%)
All	All	823/855 (96%)	-0.38	13 (1%)	70	64	22, 43, 64, 133	17 (2%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	1	ALA	18.0
3	M	2	GLU	6.0
3	M	301	HIS	5.5
1	H	247	LYS	3.9
1	H	249[A]	LYS	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
11	HTO	L	309	10/10	0.65	0.31	72,96,120,121	0
8	LDA	L	302	16/16	0.73	0.38	47,125,153,155	0
16	PC1	M	410	43/54	0.73	0.38	61,99,145,170	0
4	GOL	H	306	6/6	0.77	0.21	69,76,89,90	0
8	LDA	L	303	16/16	0.83	0.33	80,90,108,114	0
6	GGD	H	307	57/67	0.85	0.23	51,108,185,203	0
15	CDL	M	409	81/100	0.87	0.26	53,96,139,153	0
8	LDA	L	304	16/16	0.88	0.24	71,98,123,128	0
11	HTO	L	308	10/10	0.90	0.22	66,87,102,105	0
4	GOL	H	303	6/6	0.90	0.13	85,89,92,95	0
4	GOL	L	310	6/6	0.91	0.15	73,81,85,86	0
8	LDA	M	404	16/16	0.91	0.18	59,69,92,96	0
4	GOL	L	311	6/6	0.92	0.12	83,88,91,94	0
4	GOL	H	302	6/6	0.93	0.15	81,82,85,90	0
9	U10	L	305[A]	23/63	0.93	0.13	33,37,65,71	23
9	U10	L	305[B]	23/63	0.93	0.13	35,50,60,62	23
9	U10	M	407	48/63	0.93	0.15	33,46,91,110	0
4	GOL	H	301	6/6	0.94	0.13	59,65,69,80	0
4	GOL	H	304	6/6	0.94	0.13	40,52,63,71	0
7	BCL	M	401	66/66	0.95	0.11	28,38,95,105	0
13	BPH	M	406	65/65	0.95	0.13	35,47,119,134	0
13	BPH	M	412	65/65	0.96	0.09	30,39,48,57	0
14	SPO	M	408	42/42	0.96	0.12	33,47,80,92	0
8	LDA	M	403	16/16	0.96	0.12	52,68,77,78	0
7	BCL	L	301	66/66	0.96	0.10	26,34,56,61	0
10	PO4	L	307	5/5	0.97	0.10	60,62,66,67	0
7	BCL	L	306	66/66	0.97	0.07	28,39,49,65	0
5	K	H	305	1/1	0.98	0.04	50,50,50,50	0
7	BCL	M	402	66/66	0.98	0.08	33,40,51,75	0
17	MG	M	411	1/1	0.99	0.03	39,39,39,39	0
12	FE	M	405	1/1	1.00	0.02	36,36,36,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PC1 M 410:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GGD H 307:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around CDL M 409:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around U10 L 305 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around U10 L 305 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around U10 M 407:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL M 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around BPH M 406:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BPH M 412:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SPO M 408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BCL L 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL L 306:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BCL M 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.