



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:54 PM UTC

PDB ID : 2IOL / pdb_00002iol
Title : Crystal structure of the C-terminal MA3 domain of Pdcd4 (mouse); form 1
Authors : Wlodawer, A.; LaRonde-LeBlanc, N.A.
Deposited on : 2006-10-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	11% 69% 11% 5% 15%
1	B	150	52% 61% 21% • 15%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Programmed Cell Death 4, Pdcd4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	Se	0	0	0
			1030	662	166	195	3	4			
1	B	127	Total	C	N	O	S	Se	0	0	0
			1030	662	166	195	3	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	333	MSE	MET	modified residue	UNP Q61823
A	370	MSE	MET	modified residue	UNP Q61823
A	382	MSE	MET	modified residue	UNP Q61823
A	401	MSE	MET	modified residue	UNP Q61823
B	333	MSE	MET	modified residue	UNP Q61823
B	370	MSE	MET	modified residue	UNP Q61823
B	382	MSE	MET	modified residue	UNP Q61823
B	401	MSE	MET	modified residue	UNP Q61823


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total	O	0	0
			36	36		
2	B	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Programmed Cell Death 4, Pdcd4



• Molecule 1: Programmed Cell Death 4, Pdcd4



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	64.65Å 64.65Å 164.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.00-2.00) 97.9 (30.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.259 , 0.297 0.257 , 0.295	Depositor DCC
R_{free} test set	1381 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2110	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/1047	0.97	0/1409
1	B	0.99	5/1047 (0.5%)	1.05	2/1409 (0.1%)
All	All	0.92	5/2094 (0.2%)	1.01	2/2818 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	342	GLY	C-O	12.04	1.40	1.24
1	B	342	GLY	C-N	9.94	1.50	1.33
1	B	333	MSE	CG-SE	7.26	2.17	1.95
1	B	333	MSE	SE-CE	7.02	2.16	1.95
1	B	401	MSE	SE-CE	-5.47	1.79	1.95

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	322	GLN	CA-C-N	7.02	127.43	119.92
1	B	322	GLN	C-N-CA	7.02	127.43	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1030	0	1028	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1030	0	1028	30	0
2	A	36	0	0	1	0
2	B	14	0	0	5	0
All	All	2110	0	2056	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:333:MSE:SE	1:B:333:MSE:CE	2.16	1.44
1:B:333:MSE:SE	1:B:333:MSE:CG	2.17	1.42
1:B:391:TRP:CD1	1:B:401:MSE:HE3	1.98	0.97
1:B:391:TRP:HD1	1:B:401:MSE:HE3	1.37	0.82
1:A:349:HIS:O	1:A:353:GLU:HG3	1.79	0.81
1:B:348:GLU:O	1:B:352:LYS:HG2	1.88	0.73
1:A:397:THR:HG23	2:A:39:HOH:O	1.90	0.72
1:B:359:PHE:CZ	1:B:362:GLU:HG3	2.25	0.71
1:A:403:ARG:CG	1:A:403:ARG:HH11	2.04	0.70
1:B:344:ILE:HG21	1:B:389:SER:OG	1.97	0.65
1:A:375:THR:N	1:B:410:ASN:HD21	1.95	0.64
1:A:375:THR:H	1:B:410:ASN:HD21	1.47	0.62
1:B:391:TRP:HD1	1:B:401:MSE:CE	2.11	0.62
1:A:396:ILE:CG2	1:A:401:MSE:HG2	2.30	0.62
1:B:326:HIS:O	1:B:330:GLU:HG2	1.99	0.62
1:B:388:LYS:CE	2:B:38:HOH:O	2.47	0.61
1:B:333:MSE:SE	1:B:333:MSE:CB	2.99	0.61
1:B:396:ILE:CG2	1:B:401:MSE:HE2	2.37	0.55
1:A:396:ILE:HG22	1:A:401:MSE:HG2	1.89	0.55
1:B:388:LYS:HE3	2:B:38:HOH:O	2.06	0.55
1:A:356:VAL:O	1:A:356:VAL:HG13	2.07	0.55
1:B:402:LYS:O	1:B:406:GLU:HG3	2.08	0.54
1:A:324:VAL:HG11	1:A:329:LYS:HE2	1.90	0.54
1:A:326:HIS:O	1:A:330:GLU:HG2	2.08	0.53
1:A:403:ARG:CG	1:A:403:ARG:NH1	2.73	0.50
1:A:403:ARG:HH11	1:A:403:ARG:HG2	1.77	0.50
1:A:356:VAL:HG13	1:A:359:PHE:HB3	1.93	0.49
1:A:403:ARG:HH11	1:A:403:ARG:HG3	1.76	0.49
1:B:397:THR:HB	2:B:24:HOH:O	2.13	0.49
1:A:391:TRP:HB2	1:A:401:MSE:HE3	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:356:VAL:HG13	1:B:359:PHE:HB3	1.94	0.48
1:A:396:ILE:HG21	1:A:401:MSE:HG2	1.95	0.48
1:B:388:LYS:HE2	2:B:38:HOH:O	2.13	0.48
1:B:356:VAL:CG1	1:B:356:VAL:O	2.62	0.47
1:B:384:LEU:HG	2:B:34:HOH:O	2.12	0.47
1:A:398:ILE:HD13	1:A:398:ILE:O	2.16	0.46
1:B:351:LEU:HD22	1:B:363:LEU:HD13	1.98	0.45
1:A:334:LEU:HD23	1:A:334:LEU:C	2.42	0.45
1:B:396:ILE:HG22	1:B:401:MSE:HE2	1.99	0.45
1:B:362:GLU:HG2	1:B:407:ARG:HH12	1.84	0.43
1:B:361:HIS:CG	1:B:403:ARG:HG2	2.53	0.43
1:B:388:LYS:HA	1:B:438:ILE:HG21	2.01	0.43
1:B:365:TYR:O	1:B:369:VAL:HG23	2.18	0.43
1:B:388:LYS:HE2	1:B:388:LYS:HB3	1.87	0.42
1:A:388:LYS:HA	1:A:438:ILE:HG21	2.02	0.42
1:A:412:ILE:HG21	1:A:448:PRO:HG2	2.03	0.41
1:B:356:VAL:HG13	1:B:359:PHE:CB	2.51	0.40
1:B:367:ALA:HB1	1:B:383:ILE:HG23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/150 (83%)	125 (100%)	0	0	100	100
1	B	125/150 (83%)	123 (98%)	2 (2%)	0	100	100
All	All	250/300 (83%)	248 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/133 (89%)	107 (91%)	11 (9%)	8	5
1	B	118/133 (89%)	110 (93%)	8 (7%)	14	11
All	All	236/266 (89%)	217 (92%)	19 (8%)	11	7

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	324	VAL
1	A	329	LYS
1	A	334	LEU
1	A	356	VAL
1	A	363	LEU
1	A	397	THR
1	A	398	ILE
1	A	403	ARG
1	A	417	LEU
1	A	443	LEU
1	A	444	ARG
1	B	324	VAL
1	B	356	VAL
1	B	397	THR
1	B	398	ILE
1	B	399	ASP
1	B	403	ARG
1	B	443	LEU
1	B	444	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	416	ASN
1	B	349	HIS
1	B	358	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	410	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	123/150 (82%)	1.18	16 (13%) 7 6	43, 50, 57, 59	0
1	B	123/150 (82%)	2.62	78 (63%) 0 0	41, 52, 59, 61	0
All	All	246/300 (82%)	1.90	94 (38%) 1 1	41, 51, 59, 61	0

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	VAL	7.5
1	B	417	LEU	6.4
1	B	397	THR	5.3
1	B	415	ILE	5.1
1	B	341	SER	4.9
1	B	391	TRP	4.6
1	B	346	GLU	4.6
1	B	424	SER	4.6
1	B	376	GLY	4.6
1	B	323	PRO	4.5
1	B	378	SER	4.4
1	B	377	GLU	4.3
1	B	322	GLN	4.1
1	B	410	ASN	4.0
1	B	328	VAL	4.0
1	B	419	VAL	4.0
1	B	344	ILE	4.0
1	B	440	SER	3.9
1	B	393	SER	3.9
1	B	404	GLY	3.8
1	B	399	ASP	3.7
1	B	338	TYR	3.7
1	B	403	ARG	3.7
1	B	422	SER	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	439	ILE	3.6
1	B	437	GLY	3.6
1	B	340	LEU	3.6
1	B	402	LYS	3.5
1	B	335	LEU	3.5
1	B	400	GLN	3.5
1	A	345	SER	3.5
1	B	385	ASP	3.5
1	B	413	PRO	3.5
1	A	397	THR	3.4
1	B	356	VAL	3.4
1	B	443	LEU	3.4
1	B	425	VAL	3.4
1	B	420	PRO	3.4
1	B	434	PHE	3.4
1	B	421	HIS	3.3
1	B	358	HIS	3.3
1	B	446	LEU	3.3
1	B	369	VAL	3.3
1	B	381	LYS	3.3
1	B	418	ASP	3.3
1	B	325	ASN	3.2
1	B	423	TYR	3.1
1	B	330	GLU	3.1
1	B	349	HIS	3.1
1	B	436	ALA	3.1
1	A	374	SER	3.1
1	A	391	TRP	3.0
1	B	395	THR	3.0
1	B	428	ARG	3.0
1	B	334	LEU	3.0
1	B	398	ILE	2.9
1	A	421	HIS	2.9
1	B	324	VAL	2.9
1	B	447	CYS	2.8
1	B	389	SER	2.8
1	A	322	GLN	2.8
1	B	359	PHE	2.7
1	A	377	GLU	2.7
1	B	388	LYS	2.7
1	B	427	GLU	2.6
1	B	339	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	337	GLU	2.6
1	B	412	ILE	2.6
1	B	448	PRO	2.6
1	B	445	ASP	2.5
1	A	424	SER	2.5
1	B	396	ILE	2.5
1	A	410	ASN	2.4
1	B	405	TYR	2.4
1	B	411	GLU	2.4
1	B	345	SER	2.4
1	B	367	ALA	2.3
1	B	375	THR	2.3
1	A	376	GLY	2.3
1	A	413	PRO	2.2
1	A	418	ASP	2.2
1	B	354	LEU	2.2
1	B	384	LEU	2.2
1	B	365	TYR	2.2
1	B	336	LYS	2.2
1	B	406	GLU	2.1
1	A	405	TYR	2.1
1	B	408	ILE	2.1
1	B	343	ASP	2.1
1	A	355	GLU	2.0
1	B	390	LEU	2.0
1	A	416	ASN	2.0
1	B	326	HIS	2.0
1	B	444	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.