



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 06:10 AM UTC

PDB ID : 4IOA / pdb_00004ioa
Title : Crystal structure of compound 4e bound to large ribosomal subunit (50S) from *Deinococcus radiodurans*
Authors : Han, S.; Marr, E.S.
Deposited on : 2013-01-07
Resolution : 3.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)
RNA backbone	3983	1222 (3.50-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	2880	
2	Y	123	
3	A	274	
4	B	211	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	C	205	
6	D	180	
7	E	185	
8	F	144	
9	G	174	
10	H	134	
11	I	156	
12	J	141	
13	K	116	
14	L	114	
15	M	166	
16	N	118	
17	O	100	
18	P	134	
19	Q	95	
20	R	115	
21	S	237	
22	T	91	
23	U	81	
24	V	67	
25	W	55	
26	Z	60	
27	1	55	
28	2	47	
29	3	66	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
30	4	37	<p>84% 68% 27% 5%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
31	MG	X	2904	-	-	-	X
31	MG	X	2922	-	-	-	X
31	MG	X	2930	-	-	-	X

2 Entry composition [i](#)

There are 32 unique types of molecules in this entry. The entry contains 83879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	X	2686	57651	25718	10642	18606	2685	0	0	0

- Molecule 2 is a RNA chain called 5S ribosomal RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	Y	122	2598	1161	476	840	121	0	0	0

- Molecule 3 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	240	1826	1137	366	321	2	0	0	0

- Molecule 4 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	205	1539	965	295	271	8	0	0	0

- Molecule 5 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	C	197	1506	935	287	282	2	0	0	0

- Molecule 6 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	D	177	1400	892	247	254	7	0	0	0

- Molecule 7 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	E	171	1286	812	237	236	1	0	0	0

- Molecule 8 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	F	71	503	310	91	99	3	0	0	0

- Molecule 9 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	G	142	1114	704	209	198	3	0	0	0

- Molecule 10 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	H	134	997	614	198	180	5	0	0	0

- Molecule 11 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
11	I	141	1067	655	216	196	0	0	0

- Molecule 12 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	J	136	1090	696	202	185	7	0	0	0

- Molecule 13 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	K	113	878	541	178	157	2	0	0	0

- Molecule 14 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
14	L	104	779	476	161	142	0	0	0

- Molecule 15 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
15	M	108	871	543	172	156	0	0	0

- Molecule 16 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
16	N	117	978	608	210	159	1	0	0	0

- Molecule 17 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
17	O	94	741	465	139	137	0	0	0

- Molecule 18 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
18	P	127	1014	639	199	174	2	0	0	0

- Molecule 19 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
19	Q	93	726	458	136	130	2	0	0	0

- Molecule 20 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
20	R	110	825	513	160	151	1	0	0	0

- Molecule 21 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
21	S	175	1345	849	236	254	6	0	0	0

- Molecule 22 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
22	T	84	625	393	122	109	1	0	0	0

- Molecule 23 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
23	U	72	552	341	116	95	0	0	0

- Molecule 24 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
24	V	66	533	327	107	96	3	0	0	0

- Molecule 25 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
25	W	55	424	264	82	76	2	0	0	0

- Molecule 26 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
26	Z	58	457	281	94	77	5	0	0	0

- Molecule 27 is a protein called 50S ribosomal protein L33.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
			Total	C			
27	1	53	53	53	0	0	53

- Molecule 28 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
28	2	46	Total C 46 46	0	0	46

- Molecule 29 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
29	3	63	Total C 63 63	0	0	63


- Molecule 30 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
30	4	37	Total C N O S 297 179 66 47 5	0	0	0


- Molecule 31 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
31	X	30	Total Mg 30 30	0	0
31	Y	5	Total Mg 5 5	0	0


- Molecule 32 is (3aS,4R,7R,8S,9S,10R,11R,13R,15R,15aR)-4-ethyl-11-methoxy-3a,7,9,11,13,15-hexamethyl-2,6,14-trioxo-10-{{3,4,6-trideoxy-3-(dimethylamino)-beta-D-xylo-hexopyranosyl}oxy}tetradecahydro-2H-oxacyclotetradecino[4,3-d][1,3]oxazol-8-yl (2R)-2-[4-(acetylamino)phenyl]-2,3-dihydro-1H-pyrrole-1-carboxylate (CCD ID: 1F3) (formula: C₄₄H₆₆N₄O₁₂).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
32	X	1	60	44	4	12	0	0




• Molecule 4: 50S ribosomal protein L3




• Molecule 5: 50S ribosomal protein L4




• Molecule 6: 50S ribosomal protein L5








• Molecule 7: 50S ribosomal protein L6



• Molecule 8: 50S ribosomal protein L11




• Molecule 9: 50S ribosomal protein L13




• Molecule 10: 50S ribosomal protein L14







• Molecule 11: 50S ribosomal protein L15




• Molecule 12: 50S ribosomal protein L16



• Molecule 13: 50S ribosomal protein L17






• Molecule 14: 50S ribosomal protein L18







- Molecule 15: 50S ribosomal protein L19




- Molecule 16: 50S ribosomal protein L20




- Molecule 17: 50S ribosomal protein L21




- Molecule 18: 50S ribosomal protein L22




- Molecule 19: 50S ribosomal protein L23




• Molecule 20: 50S ribosomal protein L24




• Molecule 21: 50S ribosomal protein L25




• Molecule 22: 50S ribosomal protein L27



• Molecule 23: 50S ribosomal protein L28





- Molecule 24: 50S ribosomal protein L29




- Molecule 25: 50S ribosomal protein L30





- Molecule 26: 50S ribosomal protein L32





- Molecule 27: 50S ribosomal protein L33




- Molecule 28: 50S ribosomal protein L34



- Molecule 29: 50S ribosomal protein L35



- Molecule 30: 50S ribosomal protein L36



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	169.77Å 406.66Å 696.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.20 30.00 – 3.23	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-3.20) 89.7 (30.00-3.23)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.51 (at 3.24Å)	Xtrriage
Refinement program	autoBUSTER	Depositor
R, R_{free}	0.197 , 0.230 0.210 , 0.246	Depositor DCC
R_{free} test set	17364 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å ²)	80.0	Xtrriage
Anisotropy	0.713	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 84.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	83879	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 1F3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	X	0.69	21/64561 (0.0%)	1.38	832/100708 (0.8%)
2	Y	0.69	0/2904	1.24	12/4525 (0.3%)
3	A	1.07	9/1862 (0.5%)	1.62	31/2510 (1.2%)
4	B	1.01	3/1567 (0.2%)	1.55	21/2105 (1.0%)
5	C	1.04	3/1529 (0.2%)	1.70	41/2070 (2.0%)
6	D	0.86	0/1419	1.31	4/1903 (0.2%)
7	E	0.82	0/1308	1.33	4/1771 (0.2%)
8	F	0.92	0/508	1.38	0/683
9	G	1.01	2/1138 (0.2%)	1.71	29/1539 (1.9%)
10	H	0.94	1/1007 (0.1%)	1.48	12/1352 (0.9%)
11	I	1.25	9/1081 (0.8%)	1.83	32/1448 (2.2%)
12	J	1.10	2/1113 (0.2%)	1.61	17/1486 (1.1%)
13	K	1.25	4/886 (0.5%)	1.76	21/1188 (1.8%)
14	L	0.97	0/785	1.75	12/1048 (1.1%)
15	M	1.06	5/884 (0.6%)	1.60	15/1186 (1.3%)
16	N	0.86	0/994	1.55	6/1323 (0.5%)
17	O	0.94	0/750	1.60	14/1000 (1.4%)
18	P	0.93	0/1027	1.47	10/1373 (0.7%)
19	Q	1.05	1/737 (0.1%)	1.70	18/988 (1.8%)
20	R	1.12	1/835 (0.1%)	1.68	23/1121 (2.1%)
21	S	0.92	0/1370	1.40	10/1862 (0.5%)
22	T	0.92	0/633	1.44	11/838 (1.3%)
23	U	1.33	3/556 (0.5%)	1.92	22/741 (3.0%)
24	V	0.83	0/537	1.56	5/714 (0.7%)
25	W	0.79	0/426	1.48	7/568 (1.2%)
26	Z	1.07	1/469 (0.2%)	1.69	11/629 (1.7%)
30	4	0.95	0/298	1.31	0/390
All	All	0.79	65/91184 (0.1%)	1.43	1220/137069 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected

by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	6

The worst 5 of 65 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	I	38	LYS	CA-C	11.43	1.62	1.52
13	K	52	ILE	CG1-CD1	10.65	1.93	1.51
26	Z	53	ASP	CA-C	8.64	1.64	1.52
15	M	29	PRO	CA-C	8.60	1.64	1.52
11	I	29	THR	CA-C	8.14	1.63	1.52

The worst 5 of 1220 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	1288	A	C1'-O4'-C4'	-30.81	78.89	109.70
1	X	1288	A	C4'-C3'-C2'	-19.04	83.56	102.60
1	X	1288	A	C5'-C4'-O4'	16.75	134.22	109.10
11	I	64	GLY	N-CA-C	15.62	126.04	111.67
1	X	1716	G	P-O3'-C3'	14.94	142.60	120.20

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	1288	A	Sidechain
1	X	219	G	Sidechain
1	X	474	G	Sidechain
1	X	683	A	Sidechain
1	X	739	G	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	57651	0	29049	437	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Y	2598	0	1328	17	0
3	A	1826	0	1885	63	0
4	B	1539	0	1600	67	0
5	C	1506	0	1525	58	0
6	D	1400	0	1481	23	0
7	E	1286	0	1336	9	0
8	F	503	0	520	3	0
9	G	1114	0	1144	64	0
10	H	997	0	1046	23	0
11	I	1067	0	1103	48	0
12	J	1090	0	1125	32	0
13	K	878	0	930	29	0
14	L	779	0	820	17	0
15	M	871	0	894	30	0
16	N	978	0	1020	26	0
17	O	741	0	756	31	0
18	P	1014	0	1096	22	0
19	Q	726	0	753	23	0
20	R	825	0	881	29	0
21	S	1345	0	1372	23	0
22	T	625	0	655	11	0
23	U	552	0	604	28	0
24	V	533	0	558	5	0
25	W	424	0	470	9	0
26	Z	457	0	462	16	0
27	1	53	0	0	0	0
28	2	46	0	0	2	0
29	3	63	0	0	2	0
30	4	297	0	330	6	0
31	X	30	0	0	0	0
31	Y	5	0	0	0	0
32	X	60	0	66	2	0
All	All	83879	0	54809	994	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 994 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:K:52:ILE:CD1	13:K:52:ILE:CG1	1.93	1.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:I:62:LYS:NZ	11:I:64:GLY:HA2	1.60	1.15
1:X:1333:G:N2	1:X:1344:C:H41	1.44	1.12
15:M:79:ARG:HG3	15:M:79:ARG:HH11	1.03	1.10
19:Q:29:VAL:HG11	19:Q:38:ILE:HD11	1.35	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	238/274 (87%)	170 (71%)	39 (16%)	29 (12%)	0	1
4	B	203/211 (96%)	172 (85%)	17 (8%)	14 (7%)	1	7
5	C	195/205 (95%)	131 (67%)	37 (19%)	27 (14%)	0	1
6	D	175/180 (97%)	136 (78%)	26 (15%)	13 (7%)	1	6
7	E	169/185 (91%)	132 (78%)	29 (17%)	8 (5%)	2	14
8	F	69/144 (48%)	59 (86%)	8 (12%)	2 (3%)	3	24
9	G	140/174 (80%)	103 (74%)	23 (16%)	14 (10%)	0	2
10	H	132/134 (98%)	117 (89%)	9 (7%)	6 (4%)	2	15
11	I	139/156 (89%)	79 (57%)	34 (24%)	26 (19%)	0	0
12	J	134/141 (95%)	97 (72%)	26 (19%)	11 (8%)	0	4
13	K	111/116 (96%)	96 (86%)	7 (6%)	8 (7%)	1	6
14	L	102/114 (90%)	72 (71%)	20 (20%)	10 (10%)	0	2
15	M	106/166 (64%)	95 (90%)	7 (7%)	4 (4%)	2	18
16	N	115/118 (98%)	101 (88%)	10 (9%)	4 (4%)	3	20
17	O	92/100 (92%)	66 (72%)	13 (14%)	13 (14%)	0	1
18	P	125/134 (93%)	114 (91%)	5 (4%)	6 (5%)	2	14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
19	Q	91/95 (96%)	59 (65%)	19 (21%)	13 (14%)	0	1
20	R	108/115 (94%)	66 (61%)	23 (21%)	19 (18%)	0	0
21	S	173/237 (73%)	145 (84%)	20 (12%)	8 (5%)	2	15
22	T	82/91 (90%)	64 (78%)	9 (11%)	9 (11%)	0	2
23	U	70/81 (86%)	39 (56%)	17 (24%)	14 (20%)	0	0
24	V	64/67 (96%)	57 (89%)	3 (5%)	4 (6%)	1	8
25	W	53/55 (96%)	49 (92%)	4 (8%)	0	100	100
26	Z	56/60 (93%)	47 (84%)	5 (9%)	4 (7%)	1	6
30	4	35/37 (95%)	28 (80%)	5 (14%)	2 (6%)	1	11
All	All	2977/3390 (88%)	2294 (77%)	415 (14%)	268 (9%)	0	3

5 of 268 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	60	ARG
3	A	151	LYS
3	A	170	SER
3	A	187	SER
3	A	199	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	185/215 (86%)	139 (75%)	46 (25%)	0	3
4	B	155/157 (99%)	129 (83%)	26 (17%)	2	11
5	C	157/163 (96%)	123 (78%)	34 (22%)	1	6
6	D	153/156 (98%)	128 (84%)	25 (16%)	2	12
7	E	136/144 (94%)	115 (85%)	21 (15%)	2	14
8	F	51/107 (48%)	46 (90%)	5 (10%)	7	31
9	G	118/146 (81%)	92 (78%)	26 (22%)	1	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	H	103/103 (100%)	76 (74%)	27 (26%)	0	3
11	I	108/121 (89%)	71 (66%)	37 (34%)	0	0
12	J	110/115 (96%)	90 (82%)	20 (18%)	2	9
13	K	90/93 (97%)	73 (81%)	17 (19%)	1	9
14	L	74/82 (90%)	54 (73%)	20 (27%)	0	2
15	M	94/134 (70%)	69 (73%)	25 (27%)	0	3
16	N	96/97 (99%)	79 (82%)	17 (18%)	2	10
17	O	75/79 (95%)	61 (81%)	14 (19%)	1	9
18	P	109/115 (95%)	93 (85%)	16 (15%)	3	16
19	Q	75/76 (99%)	60 (80%)	15 (20%)	1	7
20	R	91/96 (95%)	70 (77%)	21 (23%)	1	5
21	S	149/192 (78%)	123 (83%)	26 (17%)	2	10
22	T	62/67 (92%)	54 (87%)	8 (13%)	4	20
23	U	57/66 (86%)	31 (54%)	26 (46%)	0	0
24	V	54/55 (98%)	46 (85%)	8 (15%)	3	15
25	W	48/48 (100%)	36 (75%)	12 (25%)	0	3
26	Z	51/53 (96%)	36 (71%)	15 (29%)	0	1
30	4	35/35 (100%)	29 (83%)	6 (17%)	2	11
All	All	2436/2715 (90%)	1923 (79%)	513 (21%)	1	6

5 of 513 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
23	U	49	LYS
24	V	25	LEU
23	U	46	LEU
10	H	41	ASN
10	H	26	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 73 such sidechains are listed below:

Mol	Chain	Res	Type
21	S	105	GLN
30	4	13	ASN
21	S	132	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
25	W	15	ASN
10	H	26	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	X	2681/2880 (93%)	664 (24%)	238 (8%)
2	Y	121/123 (98%)	28 (23%)	5 (4%)
All	All	2802/3003 (93%)	692 (24%)	243 (8%)

5 of 692 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	X	2	G
1	X	3	U
1	X	4	C
1	X	13	A
1	X	14	A

5 of 243 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	X	1249	G
1	X	2705	A
1	X	1581	C
1	X	2669	C
1	X	2854	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 35 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
32	1F3	X	2931	-	63,64,64	1.32	7 (11%)	83,96,96	1.87	19 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
32	1F3	X	2931	-	-	9/78/119/119	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
32	X	2931	1F3	C50-C49	4.20	1.50	1.34
32	X	2931	1F3	C48-C47	-4.14	1.51	1.54
32	X	2931	1F3	C50-N45	3.34	1.44	1.38
32	X	2931	1F3	O17-C5	-3.08	1.43	1.47
32	X	2931	1F3	C47-N45	-2.63	1.43	1.47

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
32	X	2931	1F3	C59-C58-N57	-5.48	106.72	114.95
32	X	2931	1F3	C28-O20-C8	-4.96	107.83	116.26
32	X	2931	1F3	C23-C6-C5	-4.94	108.42	115.23
32	X	2931	1F3	O60-C58-N57	3.83	128.31	123.06
32	X	2931	1F3	C47-N45-C50	3.52	113.23	109.61

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
32	X	2931	1F3	C59-C58-N57-C54

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
32	X	2931	1F3	O60-C58-N57-C54
32	X	2931	1F3	C16-C1-C4-N40
32	X	2931	1F3	O20-C8-C9-C7
32	X	2931	1F3	C3-C2-C7-C9

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
32	X	2931	1F3	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	2686/2880 (93%)	0.23	145 (5%) 31 20	43, 87, 194, 279	0
2	Y	122/123 (99%)	0.71	9 (7%) 20 13	82, 129, 165, 187	0
3	A	240/274 (87%)	1.33	56 (23%) 2 2	63, 107, 137, 156	0
4	B	205/211 (97%)	0.30	10 (4%) 35 22	38, 68, 99, 145	0
5	C	197/205 (96%)	1.40	54 (27%) 1 1	55, 107, 150, 178	0
6	D	177/180 (98%)	1.32	46 (25%) 1 2	148, 178, 210, 216	0
7	E	171/185 (92%)	0.96	25 (14%) 6 4	98, 139, 178, 188	0
8	F	71/144 (49%)	2.54	38 (53%) 0 1	221, 234, 251, 259	0
9	G	142/174 (81%)	1.32	37 (26%) 1 2	65, 89, 137, 149	0
10	H	134/134 (100%)	0.04	2 (1%) 72 52	49, 62, 88, 110	0
11	I	141/156 (90%)	2.25	56 (39%) 1 1	54, 120, 171, 195	0
12	J	136/141 (96%)	1.05	28 (20%) 2 2	83, 106, 147, 172	0
13	K	113/116 (97%)	0.15	7 (6%) 26 17	37, 53, 71, 99	0
14	L	104/114 (91%)	1.55	32 (30%) 1 1	91, 122, 149, 166	0
15	M	108/166 (65%)	0.27	6 (5%) 30 19	44, 64, 106, 128	0
16	N	117/118 (99%)	0.68	15 (12%) 7 6	54, 86, 124, 152	0
17	O	94/100 (94%)	0.94	15 (15%) 5 3	67, 106, 146, 160	0
18	P	127/134 (94%)	-0.01	2 (1%) 70 51	48, 64, 103, 143	0
19	Q	93/95 (97%)	1.14	20 (21%) 2 2	69, 101, 156, 193	0
20	R	110/115 (95%)	1.60	33 (30%) 1 1	84, 113, 170, 173	0
21	S	175/237 (73%)	1.08	28 (16%) 5 3	119, 154, 178, 190	0
22	T	84/91 (92%)	1.20	17 (20%) 3 2	72, 103, 176, 195	0
23	U	72/81 (88%)	2.11	32 (44%) 0 1	86, 122, 146, 182	0
24	V	66/67 (98%)	0.85	8 (12%) 8 6	88, 128, 213, 230	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
25	W	55/55 (100%)	0.28	0 100 100	76, 96, 123, 161	0
26	Z	58/60 (96%)	0.61	4 (6%) 23 14	47, 64, 96, 108	0
27	1	53/55 (96%)	5.79	35 (66%) 0 0	6, 28, 62, 73	0
28	2	46/47 (97%)	5.24	39 (84%) 0 0	3, 10, 27, 42	0
29	3	63/66 (95%)	5.68	43 (68%) 0 0	3, 18, 41, 84	0
30	4	37/37 (100%)	5.68	31 (83%) 0 0	191, 239, 247, 252	0
All	All	5997/6561 (91%)	0.82	873 (14%) 6 4	3, 96, 193, 279	0

The worst 5 of 873 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
27	1	7	ARG	50.4
27	1	4	ASP	44.6
29	3	60	LEU	39.0
28	2	46	ASP	21.9
29	3	41	ILE	19.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
31	MG	X	2922	1/1	0.53	0.83	81,81,81,81	0
31	MG	X	2904	1/1	0.58	0.92	90,90,90,90	0
31	MG	X	2930	1/1	0.80	1.20	71,71,71,71	0
31	MG	X	2910	1/1	0.82	0.42	85,85,85,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
31	MG	Y	205	1/1	0.84	0.12	77,77,77,77	0
31	MG	X	2912	1/1	0.85	0.31	60,60,60,60	0
31	MG	X	2906	1/1	0.86	0.85	79,79,79,79	0
31	MG	X	2901	1/1	0.86	0.62	110,110,110,110	0
31	MG	X	2911	1/1	0.87	0.34	35,35,35,35	0
31	MG	X	2929	1/1	0.87	0.35	77,77,77,77	0
31	MG	X	2926	1/1	0.88	0.51	56,56,56,56	0
31	MG	X	2907	1/1	0.88	0.38	53,53,53,53	0
31	MG	Y	201	1/1	0.89	0.63	82,82,82,82	0
31	MG	X	2908	1/1	0.89	0.39	49,49,49,49	0
31	MG	X	2925	1/1	0.90	0.50	39,39,39,39	0
31	MG	X	2913	1/1	0.91	0.45	66,66,66,66	0
31	MG	X	2905	1/1	0.92	0.22	104,104,104,104	0
31	MG	Y	203	1/1	0.92	0.47	87,87,87,87	0
31	MG	Y	204	1/1	0.92	0.16	67,67,67,67	0
31	MG	X	2923	1/1	0.92	0.63	74,74,74,74	0
31	MG	X	2916	1/1	0.93	0.50	53,53,53,53	0
31	MG	X	2903	1/1	0.93	0.17	82,82,82,82	0
31	MG	Y	202	1/1	0.93	0.34	54,54,54,54	0
31	MG	X	2920	1/1	0.94	0.56	38,38,38,38	0
31	MG	X	2927	1/1	0.94	0.20	106,106,106,106	0
31	MG	X	2919	1/1	0.94	0.60	56,56,56,56	0
31	MG	X	2924	1/1	0.95	0.42	39,39,39,39	0
32	1F3	X	2931	60/60	0.95	0.10	38,60,90,99	0
31	MG	X	2917	1/1	0.96	0.47	37,37,37,37	0
31	MG	X	2902	1/1	0.97	0.63	45,45,45,45	0
31	MG	X	2915	1/1	0.97	0.26	24,24,24,24	0
31	MG	X	2914	1/1	0.98	0.54	51,51,51,51	0
31	MG	X	2918	1/1	0.98	0.48	32,32,32,32	0
31	MG	X	2921	1/1	0.98	0.38	18,18,18,18	0
31	MG	X	2928	1/1	0.98	0.58	42,42,42,42	0
31	MG	X	2909	1/1	0.99	0.45	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.