



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 05:00 PM UTC

PDB ID : 5IOV / pdb_00005iov
EMDB ID : EMD-8098
Title : Cryo-EM structure of GluN1/GluN2B NMDA receptor in the glutamate/glycine/Ro25-6981-bound conformation
Authors : Zhu, S.; Stein, A.R.; Yoshioka, C.; Lee, C.H.; Goehring, A.; Mchaourab, S.H.; Gouaux, E.
Deposited on : 2016-03-09
Resolution : 7.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 7.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	436 (7.00 - 8.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	822	15% 90% 7% .
1	C	822	21% 91% 5% ..
2	B	825	24% 87% 9% ..
2	D	825	28% 86% 9% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	QEM	B	901	-	-	X	-
4	QEM	D	901	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-methyl-D-aspartate receptor subunit NR1-8a.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
1	A	803	3963	2358	803	802	0	0
1	C	803	3963	2358	803	802	0	0

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	PHE	LYS	engineered mutation	UNP C0KD18
A	52	PHE	ARG	engineered mutation	UNP C0KD18
A	300	GLN	ASN	engineered mutation	UNP C0KD18
A	350	GLN	ASN	engineered mutation	UNP C0KD18
A	368	ASP	ASN	engineered mutation	UNP C0KD18
A	440	ASP	ASN	engineered mutation	UNP C0KD18
A	469	ASP	ASN	engineered mutation	UNP C0KD18
A	493	ALA	LYS	engineered mutation	UNP C0KD18
A	494	ALA	LYS	engineered mutation	UNP C0KD18
A	495	ALA	GLU	engineered mutation	UNP C0KD18
A	?	-	LYS	deletion	UNP C0KD18
A	?	-	VAL	deletion	UNP C0KD18
A	?	-	ASN	deletion	UNP C0KD18
A	?	-	SER	deletion	UNP C0KD18
A	?	-	GLU	deletion	UNP C0KD18
A	?	-	GLU	deletion	UNP C0KD18
A	?	-	GLU	deletion	UNP C0KD18
A	602	ARG	GLY	engineered mutation	UNP C0KD18
A	609	LEU	ILE	engineered mutation	UNP C0KD18
A	648	ARG	ASP	engineered mutation	UNP C0KD18
A	761	GLU	ASN	engineered mutation	UNP C0KD18
A	829	SER	-	expression tag	UNP C0KD18
A	830	ARG	-	expression tag	UNP C0KD18
A	831	ALA	-	expression tag	UNP C0KD18
A	832	GLU	-	expression tag	UNP C0KD18

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	833	ALA	-	expression tag	UNP C0KD18
A	834	LYS	-	expression tag	UNP C0KD18
A	835	ARG	-	expression tag	UNP C0KD18
A	836	MET	-	expression tag	UNP C0KD18
A	837	LYS	-	expression tag	UNP C0KD18
A	838	GLY	-	expression tag	UNP C0KD18
A	839	LEU	-	expression tag	UNP C0KD18
A	840	GLU	-	expression tag	UNP C0KD18
A	841	VAL	-	expression tag	UNP C0KD18
A	842	LEU	-	expression tag	UNP C0KD18
A	843	PHE	-	expression tag	UNP C0KD18
A	844	GLN	-	expression tag	UNP C0KD18
C	51	PHE	LYS	engineered mutation	UNP C0KD18
C	52	PHE	ARG	engineered mutation	UNP C0KD18
C	300	GLN	ASN	engineered mutation	UNP C0KD18
C	350	GLN	ASN	engineered mutation	UNP C0KD18
C	368	ASP	ASN	engineered mutation	UNP C0KD18
C	440	ASP	ASN	engineered mutation	UNP C0KD18
C	469	ASP	ASN	engineered mutation	UNP C0KD18
C	493	ALA	LYS	engineered mutation	UNP C0KD18
C	494	ALA	LYS	engineered mutation	UNP C0KD18
C	495	ALA	GLU	engineered mutation	UNP C0KD18
C	?	-	LYS	deletion	UNP C0KD18
C	?	-	VAL	deletion	UNP C0KD18
C	?	-	ASN	deletion	UNP C0KD18
C	?	-	SER	deletion	UNP C0KD18
C	?	-	GLU	deletion	UNP C0KD18
C	?	-	GLU	deletion	UNP C0KD18
C	?	-	GLU	deletion	UNP C0KD18
C	?	-	GLU	deletion	UNP C0KD18
C	602	ARG	GLY	engineered mutation	UNP C0KD18
C	609	LEU	ILE	engineered mutation	UNP C0KD18
C	648	ARG	ASP	engineered mutation	UNP C0KD18
C	761	GLU	ASN	engineered mutation	UNP C0KD18
C	829	SER	-	expression tag	UNP C0KD18
C	830	ARG	-	expression tag	UNP C0KD18
C	831	ALA	-	expression tag	UNP C0KD18
C	832	GLU	-	expression tag	UNP C0KD18
C	833	ALA	-	expression tag	UNP C0KD18
C	834	LYS	-	expression tag	UNP C0KD18
C	835	ARG	-	expression tag	UNP C0KD18
C	836	MET	-	expression tag	UNP C0KD18

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	837	LYS	-	expression tag	UNP C0KD18
C	838	GLY	-	expression tag	UNP C0KD18
C	839	LEU	-	expression tag	UNP C0KD18
C	840	GLU	-	expression tag	UNP C0KD18
C	841	VAL	-	expression tag	UNP C0KD18
C	842	LEU	-	expression tag	UNP C0KD18
C	843	PHE	-	expression tag	UNP C0KD18
C	844	GLN	-	expression tag	UNP C0KD18

- Molecule 2 is a protein called Ionotropic glutamate receptor subunit NR2B.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	B	798	3939	2344	798	797	0	0
2	D	797	3934	2341	797	796	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	20	SER	MET	engineered mutation	UNP A7XY94
B	21	ARG	GLY	engineered mutation	UNP A7XY94
B	22	ALA	CYS	engineered mutation	UNP A7XY94
B	64	GLU	ALA	engineered mutation	UNP A7XY94
B	69	GLN	ASN	engineered mutation	UNP A7XY94
B	343	ASP	ASN	engineered mutation	UNP A7XY94
B	?	-	LYS	deletion	UNP A7XY94
B	?	-	TYR	deletion	UNP A7XY94
B	?	-	TYR	deletion	UNP A7XY94
B	?	-	VAL	deletion	UNP A7XY94
B	486	VAL	THR	engineered mutation	UNP A7XY94
B	?	-	ARG	deletion	UNP A7XY94
B	?	-	CYS	deletion	UNP A7XY94
B	?	-	LEU	deletion	UNP A7XY94
B	?	-	ALA	deletion	UNP A7XY94
B	?	-	ASP	deletion	UNP A7XY94
B	?	-	GLY	deletion	UNP A7XY94
B	?	-	ARG	deletion	UNP A7XY94
B	?	-	GLU	deletion	UNP A7XY94
B	?	-	PRO	deletion	UNP A7XY94
B	?	-	GLY	deletion	UNP A7XY94
B	601	LEU	VAL	engineered mutation	UNP A7XY94

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	640	ARG	GLU	engineered mutation	UNP A7XY94
B	641	ARG	GLU	engineered mutation	UNP A7XY94
D	20	SER	MET	engineered mutation	UNP A7XY94
D	21	ARG	GLY	engineered mutation	UNP A7XY94
D	22	ALA	CYS	engineered mutation	UNP A7XY94
D	64	GLU	ALA	engineered mutation	UNP A7XY94
D	69	GLN	ASN	engineered mutation	UNP A7XY94
D	343	ASP	ASN	engineered mutation	UNP A7XY94
D	?	-	LYS	deletion	UNP A7XY94
D	?	-	TYR	deletion	UNP A7XY94
D	?	-	TYR	deletion	UNP A7XY94
D	?	-	VAL	deletion	UNP A7XY94
D	486	VAL	THR	engineered mutation	UNP A7XY94
D	?	-	ARG	deletion	UNP A7XY94
D	?	-	CYS	deletion	UNP A7XY94
D	?	-	LEU	deletion	UNP A7XY94
D	?	-	ALA	deletion	UNP A7XY94
D	?	-	ASP	deletion	UNP A7XY94
D	?	-	GLY	deletion	UNP A7XY94
D	?	-	ARG	deletion	UNP A7XY94
D	?	-	GLU	deletion	UNP A7XY94
D	?	-	PRO	deletion	UNP A7XY94
D	?	-	GLY	deletion	UNP A7XY94
D	601	LEU	VAL	engineered mutation	UNP A7XY94
D	640	ARG	GLU	engineered mutation	UNP A7XY94
D	641	ARG	GLU	engineered mutation	UNP A7XY94

- Molecule 3 is GLYCINE (CCD ID: GLY) (formula: C₂H₅NO₂).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
3	A	1	5	2	1	2	0
3	C	1	5	2	1	2	0

- Molecule 4 is 4-[(1R,2S)-3-(4-benzylpiperidin-1-yl)-1-hydroxy-2-methylpropyl]phenol (CCD ID: QEM) (formula: C₂₂H₂₉NO₂).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	B	1	25	22	1	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	D	1	25	22	1	2	0

- Molecule 5 is GLUTAMIC ACID (CCD ID: GLU) (formula: C₅H₉NO₄).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	B	1	10	5	1	4	0
5	D	1	10	5	1	4	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-methyl-D-aspartate receptor subunit NR1-8a





- Molecule 1: N-methyl-D-aspartate receptor subunit NR1-8a








- Molecule 2: Ionotropic glutamate receptor subunit NR2B



- Molecule 2: Ionotropic glutamate receptor subunit NR2B





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	87851	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	10.2	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.120	Depositor
Minimum map value	-0.042	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.035	Depositor
Map size (Å)	236.47003, 215.07004, 204.37003	wwPDB
Map dimensions	221, 201, 191	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0700002, 1.0700002, 1.0700002	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: QEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.10	0/3961	0.95	27/5512 (0.5%)
1	C	0.13	0/3961	0.80	21/5512 (0.4%)
2	B	0.19	1/3938 (0.0%)	0.72	23/5482 (0.4%)
2	D	0.14	0/3932	0.76	25/5472 (0.5%)
All	All	0.15	1/15792 (0.0%)	0.81	96/21978 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2
2	D	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	393	GLU	C-N	-8.84	1.21	1.33

The worst 5 of 96 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	490	ASN	CB-CA-C	-26.18	72.96	111.70
1	A	478	ALA	N-CA-C	24.20	137.89	111.03
1	C	584	PHE	N-CA-C	23.31	141.02	112.59
1	A	584	PHE	N-CA-C	21.56	140.44	112.41
1	A	584	PHE	CB-CA-C	-20.74	77.42	111.02

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	393	GLU	Mainchain,Peptide
2	D	756	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3963	0	1809	29	0
1	C	3963	0	1808	26	0
2	B	3939	0	1750	43	0
2	D	3934	0	1747	53	0
3	A	5	0	2	0	0
3	C	5	0	2	0	0
4	B	25	0	29	11	0
4	D	25	0	29	11	0
5	B	10	0	5	0	0
5	D	10	0	5	0	0
All	All	15879	0	7186	151	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 151 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:TYR:CB	4:B:901:QEM:C01	2.14	1.25
1:C:649:ARG:O	1:C:651:GLU:N	1.72	1.23
2:B:109:PHE:CB	4:B:901:QEM:H04	1.76	1.14
2:D:102:ALA:HB1	4:D:901:QEM:H09A	1.32	1.10
2:D:462:ALA:HA	2:D:465:VAL:CB	1.81	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	799/822 (97%)	684 (86%)	106 (13%)	9 (1%)	11	46
1	C	799/822 (97%)	681 (85%)	110 (14%)	8 (1%)	12	48
2	B	796/825 (96%)	666 (84%)	110 (14%)	20 (2%)	4	26
2	D	793/825 (96%)	678 (86%)	100 (13%)	15 (2%)	6	32
All	All	3187/3294 (97%)	2709 (85%)	426 (13%)	52 (2%)	10	38

5 of 52 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	651	GLU
1	A	655	THR
1	C	506	GLY
1	C	649	ARG
1	C	650	PRO

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLY	C	1001	-	4,4,4	1.15	1 (25%)	3,4,4	1.64	1 (33%)
3	GLY	A	901	-	4,4,4	1.17	1 (25%)	3,4,4	1.65	1 (33%)
4	QEM	D	901	-	27,27,27	0.85	1 (3%)	35,36,36	1.18	4 (11%)
4	QEM	B	901	-	27,27,27	1.05	2 (7%)	35,36,36	1.27	5 (14%)
5	GLU	D	902	-	8,9,9	1.12	1 (12%)	8,11,11	1.18	1 (12%)
5	GLU	B	902	-	8,9,9	1.11	1 (12%)	8,11,11	1.18	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLY	C	1001	-	-	0/2/2/2	-
3	GLY	A	901	-	-	0/2/2/2	-
4	QEM	D	901	-	-	5/16/26/26	0/3/3/3
4	QEM	B	901	-	-	4/16/26/26	0/3/3/3
5	GLU	D	902	-	-	2/9/9/9	-
5	GLU	B	902	-	-	2/9/9/9	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	901	QEM	C06-C01	-3.09	1.33	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	902	GLU	OXT-C	-2.23	1.23	1.30
5	B	902	GLU	OXT-C	-2.23	1.23	1.30
3	A	901	GLY	OXT-C	-2.22	1.23	1.30
3	C	1001	GLY	OXT-C	-2.18	1.23	1.30

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	901	QEM	C11-N-C10	3.71	116.84	108.84
4	D	901	QEM	C11-N-C10	3.41	116.19	108.84
5	D	902	GLU	OXT-C-O	-2.71	117.94	124.08
5	B	902	GLU	OXT-C-O	-2.69	117.98	124.08
4	D	901	QEM	C14-C13-N	-2.61	109.06	115.44

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	901	QEM	C13-C14-C15-O01
4	B	901	QEM	C01-C02-C07-C08
4	B	901	QEM	C03-C02-C07-C08
4	B	901	QEM	C14-C15-C16-C
4	B	901	QEM	C14-C15-C16-C18


There are no ring outliers.

2 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	901	QEM	11	0
4	B	901	QEM	11	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-8098. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 112



Y Index: 91




Z Index: 93


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.035. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 246 nm³; this corresponds to an approximate mass of 223 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-8098 and PDB model 5IOV. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.










9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.035) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7270	 0.2030
A	 0.7870	 0.2180
B	 0.7010	 0.2020
C	 0.7480	 0.2050
D	 0.6710	 0.1880

