



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:30 AM UTC

PDB ID : 3IPU / pdb_00003ipu
Title : X-ray structure of benzisoxazole urea synthetic agonist bound to the LXR-alpha
Authors : Fradera, X.; Vu, D.; Nimz, O.; Skene, R.; Hosfield, D.; Wijnands, R.; Cooke, A.J.; Haunso, A.; King, A.; Bennet, D.J.; McGuire, R.; Uitdehaag, J.C.M.
Deposited on : 2009-08-18
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	7% 73% 10% • 16%
1	B	283	4% 73% 9% • 17%
2	C	25	44% 16% 40%
2	D	25	4% 60% 40%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxysterols receptor LXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	0	0	0
			1945	1241	344	353	7			
1	B	235	Total	C	N	O	S	12	1	0
			1927	1229	340	351	7			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	MET	-	expression tag	UNP Q13133
A	166	LYS	-	expression tag	UNP Q13133
A	167	HIS	-	expression tag	UNP Q13133
A	168	GLN	-	expression tag	UNP Q13133
A	169	HIS	-	expression tag	UNP Q13133
A	170	GLN	-	expression tag	UNP Q13133
A	171	HIS	-	expression tag	UNP Q13133
A	172	GLN	-	expression tag	UNP Q13133
A	173	HIS	-	expression tag	UNP Q13133
A	174	GLN	-	expression tag	UNP Q13133
A	175	HIS	-	expression tag	UNP Q13133
A	176	GLN	-	expression tag	UNP Q13133
A	177	HIS	-	expression tag	UNP Q13133
A	178	GLN	-	expression tag	UNP Q13133
A	179	GLN	-	expression tag	UNP Q13133
A	180	PRO	-	expression tag	UNP Q13133
A	181	LEU	-	expression tag	UNP Q13133
B	165	MET	-	expression tag	UNP Q13133
B	166	LYS	-	expression tag	UNP Q13133
B	167	HIS	-	expression tag	UNP Q13133
B	168	GLN	-	expression tag	UNP Q13133
B	169	HIS	-	expression tag	UNP Q13133
B	170	GLN	-	expression tag	UNP Q13133
B	171	HIS	-	expression tag	UNP Q13133
B	172	GLN	-	expression tag	UNP Q13133

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	173	HIS	-	expression tag	UNP Q13133
B	174	GLN	-	expression tag	UNP Q13133
B	175	HIS	-	expression tag	UNP Q13133
B	176	GLN	-	expression tag	UNP Q13133
B	177	HIS	-	expression tag	UNP Q13133
B	178	GLN	-	expression tag	UNP Q13133
B	179	GLN	-	expression tag	UNP Q13133
B	180	PRO	-	expression tag	UNP Q13133
B	181	LEU	-	expression tag	UNP Q13133

- Molecule 2 is a protein called Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	15	Total	C	N	O	4	0	0
			131	82	27	22			
2	D	15	Total	C	N	O	4	0	0
			131	82	27	22			

- Molecule 3 is 4-{{methyl(3-{{7-propyl-3-(trifluoromethyl)-1,2-benzisoxazol-6-yl}oxy}propyl)carbamoyl}amino}benzoic acid (CCD ID: O40) (formula: C₂₃H₂₄F₃N₃O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0
			34	23	3	3	5	0
3	B	1	Total	C	F	N	O	0
			34	23	3	3	5	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	35	Total	O	0	0
			35	35		
5	B	42	Total	O	0	0
			42	42		
5	C	7	Total	O	0	0
			7	7		
5	D	1	Total	O	0	0
			1	1		

i

- Molecule 1: Oxysterols receptor LXR-alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	125.18Å 125.18Å 92.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.66 – 2.40 29.66 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.66-2.40) 90.2 (29.66-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.234 , 0.291 0.235 , 0.289	Depositor DCC
R_{free} test set	1375 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 29.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.57$, $\langle L^2 \rangle = 0.41$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4297	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.5903e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: O40, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/1987	0.84	2/2686 (0.1%)
1	B	0.72	1/1969 (0.1%)	0.87	3/2661 (0.1%)
2	C	0.68	0/132	0.89	0/175
2	D	0.70	0/132	0.84	0/175
All	All	0.70	1/4220 (0.0%)	0.86	5/5697 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	226	ARG	CA-CB	7.92	1.65	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	ARG	CB-CA-C	-7.36	99.27	110.90
1	B	226	ARG	N-CA-CB	6.22	119.08	110.07
1	B	317	LYS	N-CA-C	5.55	118.09	111.71
1	A	436	PRO	CA-C-N	5.14	126.26	119.84
1	A	436	PRO	C-N-CA	5.14	126.26	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1945	0	1951	17	0
1	B	1927	0	1936	15	0
2	C	131	0	139	3	0
2	D	131	0	139	0	0
3	A	34	0	23	0	0
3	B	34	0	23	0	0
4	B	5	0	0	0	0
4	C	5	0	0	1	0
5	A	35	0	0	0	0
5	B	42	0	0	0	0
5	C	7	0	0	0	0
5	D	1	0	0	0	0
All	All	4297	0	4211	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:346:MET:HE3	1:B:351:LEU:HD22	1.59	0.82
1:B:440:SER:OG	1:B:444:ASP:OD2	2.22	0.54
2:C:694:GLN:NE2	4:C:802:SO4:O4	2.38	0.53
1:B:232:ARG:O	1:B:235:VAL:HG22	2.08	0.53
1:B:232:ARG:HG2	1:B:263:VAL:HG11	1.90	0.53
1:A:411:LEU:HD12	1:A:414:LEU:HD12	1.91	0.52
1:A:346:MET:HE1	1:A:410:LYS:HB3	1.92	0.52
1:B:230:SER:O	1:B:234:ARG:HD3	2.10	0.51
1:A:236:THR:HG23	1:A:259:GLU:OE1	2.11	0.50
1:A:409:MET:HE1	1:B:411:LEU:HD23	1.94	0.50
1:A:298:MET:HE1	1:A:443:TRP:CZ2	2.47	0.49
1:B:374:ASP:O	1:B:378:VAL:HG23	2.14	0.47
1:B:359:LEU:HD22	1:B:411:LEU:HD11	1.96	0.47
2:C:682:LEU:HD23	2:C:685:ARG:HH22	1.81	0.46
1:A:338:PRO:HB2	1:A:417:LEU:HD21	1.98	0.45
1:B:342:PHE:HB2	1:B:417:LEU:HD11	1.99	0.45
1:B:303:SER:OG	1:B:346:MET:HG2	2.17	0.44
1:A:300:LEU:HG	1:A:414:LEU:HD11	1.99	0.44
1:A:230:SER:O	1:A:234:ARG:HD2	2.18	0.44
1:A:440:SER:OG	1:A:444:ASP:OD2	2.33	0.44
1:B:346:MET:CE	1:B:351:LEU:HD22	2.40	0.44
1:A:299:LEU:HD22	1:A:417:LEU:HD13	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:405:PRO:O	1:A:409:MET:HG2	2.19	0.42
1:B:405:PRO:O	1:B:409:MET:HG2	2.19	0.42
1:A:268:ILE:HG12	1:A:297:VAL:HG11	2.01	0.42
1:A:417:LEU:HD23	1:A:417:LEU:HA	1.92	0.42
1:A:415:ARG:HG2	1:B:412:VAL:HG13	2.02	0.41
1:A:223:GLN:HA	1:A:223:GLN:HE21	1.86	0.41
1:B:228:SER:OG	1:B:267:GLU:OE2	2.35	0.41
1:B:232:ARG:CG	1:B:263:VAL:HG11	2.51	0.41
1:A:287:ILE:HG23	2:C:689:LEU:HD23	2.02	0.40
1:A:298:MET:HE1	1:A:443:TRP:HZ2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	234/283 (83%)	229 (98%)	5 (2%)	0	100	100
1	B	232/283 (82%)	228 (98%)	4 (2%)	0	100	100
2	C	13/25 (52%)	12 (92%)	1 (8%)	0	100	100
2	D	13/25 (52%)	12 (92%)	1 (8%)	0	100	100
All	All	492/616 (80%)	481 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/257 (84%)	206 (95%)	10 (5%)	24	41
1	B	214/257 (83%)	205 (96%)	9 (4%)	26	45
2	C	15/24 (62%)	15 (100%)	0	100	100
2	D	15/24 (62%)	15 (100%)	0	100	100
All	All	460/562 (82%)	441 (96%)	19 (4%)	27	46

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	GLN
1	A	223	GLN
1	A	234	ARG
1	A	257	PHE
1	A	274	GLN
1	A	320	SER
1	A	380	ARG
1	A	411	LEU
1	A	422	SER
1	A	428	LEU
1	B	205	GLN
1	B	234	ARG
1	B	320	SER
1	B	380	ARG
1	B	411	LEU
1	B	417	LEU
1	B	428	LEU
1	B	430	LEU
1	B	432	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN
1	A	223	GLN
1	A	225	ASN
1	A	266	GLN
1	A	307	ASN
1	B	251	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	307	ASN
1	B	332	GLN
1	B	399	HIS
1	B	431	GLN
2	D	694	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	O40	B	801	-	35,36,36	1.32	4 (11%)	45,51,51	1.53	6 (13%)
3	O40	A	801	-	35,36,36	1.27	4 (11%)	45,51,51	2.00	9 (20%)
4	SO4	C	802	-	4,4,4	0.20	0	6,6,6	0.45	0
4	SO4	B	802	-	4,4,4	0.20	0	6,6,6	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	O40	B	801	-	-	1/28/28/28	0/3/3/3
3	O40	A	801	-	-	0/28/28/28	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	801	O40	O30-C32	4.49	1.35	1.22
3	A	801	O40	O30-C32	4.32	1.35	1.22
3	A	801	O40	C23-N24	-2.75	1.32	1.37
3	B	801	O40	C42-N24	-2.44	1.36	1.41
3	B	801	O40	O34-C32	-2.42	1.23	1.30
3	B	801	O40	C23-N24	-2.31	1.32	1.37
3	A	801	O40	O34-C32	-2.25	1.23	1.30
3	A	801	O40	C42-N24	-2.01	1.37	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	O40	C14-C9-C5	-7.19	119.16	129.50
3	A	801	O40	N24-C23-N21	4.49	123.48	115.25
3	B	801	O40	C14-C9-C5	-4.43	123.13	129.50
3	A	801	O40	F39-C14-C9	-4.30	104.26	112.42
3	A	801	O40	C6-C5-C4	4.27	124.91	119.41
3	B	801	O40	N24-C23-N21	4.12	122.80	115.25
3	B	801	O40	F38-C14-C9	-3.87	105.07	112.42
3	B	801	O40	C6-C5-C4	3.38	123.77	119.41
3	A	801	O40	C5-C9-N8	3.29	114.10	111.36
3	A	801	O40	C1-C6-C5	-2.89	116.09	120.86
3	A	801	O40	O7-N8-C9	-2.88	107.40	108.74
3	B	801	O40	C1-C6-C5	-2.80	116.23	120.86
3	A	801	O40	O33-C23-N24	-2.47	118.04	123.46
3	A	801	O40	F38-C14-C9	-2.46	107.76	112.42
3	B	801	O40	O33-C23-N24	-2.09	118.87	123.46

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	801	O40	C3-C12-C17-C20


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	802	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/283 (84%)	0.55	20 (8%) 17 14	16, 28, 58, 68	0
1	B	235/283 (83%)	0.41	11 (4%) 36 32	17, 28, 48, 54	3 (1%)
2	C	15/25 (60%)	0.13	0 100 100	14, 25, 33, 38	1 (6%)
2	D	15/25 (60%)	0.20	1 (6%) 24 20	14, 25, 36, 40	1 (6%)
All	All	503/616 (81%)	0.46	32 (6%) 25 22	14, 28, 53, 68	5 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	PRO	5.7
1	A	432	ASP	4.3
1	A	224	CYS	3.8
1	B	318	ASP	3.8
1	B	233	LEU	3.6
1	A	204	PRO	3.5
1	A	319	PHE	3.4
1	B	222	GLN	3.2
1	A	233	LEU	3.1
1	B	319	PHE	2.9
1	B	247	ARG	2.8
1	A	229	PHE	2.8
1	A	226	ARG	2.8
1	A	245	HIS	2.8
1	A	250	ARG	2.7
1	B	234	ARG	2.6
1	A	318	ASP	2.5
1	B	239	PRO	2.5
1	A	320	SER	2.5
2	D	691	ARG	2.5
1	B	433	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	234	ARG	2.4
1	A	426	PHE	2.3
1	A	317	LYS	2.3
1	A	247	ARG	2.3
1	A	228	SER	2.3
1	B	225	ASN	2.2
1	B	205	GLN	2.2
1	B	430	LEU	2.2
1	A	430	LEU	2.2
1	A	248	GLU	2.0
1	A	239	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	O40	A	801	34/34	0.87	0.14	39,41,45,48	0
3	O40	B	801	34/34	0.92	0.11	34,36,41,44	0
4	SO4	B	802	5/5	0.99	0.05	18,20,20,21	5
4	SO4	C	802	5/5	0.99	0.06	14,14,16,17	5


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around O40 A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around O40 B 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.