



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:00 PM UTC

PDB ID : 6IPU / pdb\_00006ipu  
Title : Human nucleosome core particle containing 145 bp of DNA  
Authors : DeFalco, L.; Davey, C.A.  
Deposited on : 2018-11-04  
Resolution : 1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*






The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	
1	E	98	
2	B	82	
3	C	107	
3	G	107	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	D	95	<p>12% 93% 7%</p>
4	H	95	<p>13% 97% ..</p>
5	F	87	<p>3% 93% 6% .</p>
6	I	145	<p>% 52% 36% 12% .</p>
7	J	145	<p>2% 52% 37% 11%</p>

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 12177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	98	807	508	156	139	4	0	0	0
1	E	98	806	508	156	138	4	0	0	0

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	82	653	412	127	113	1	0	0	0

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	107	828	523	162	143	0	0	0
3	G	107	828	523	162	143	0	0	0

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	95	746	468	136	140	2	0	0	0
4	H	95	746	468	136	140	2	0	0	0

- Molecule 5 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	F	87	703	442	142	118	1	0	0	0

- Molecule 6 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	I	145	2970	1421	538	867	144	0	0	0

- Molecule 7 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
7	J	145	2969	1421	535	869	144	0	0	0

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	1	Total Cl 1 1	0	0
8	G	1	Total Cl 1 1	0	0

- Molecule 9 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	D	1	Total Mn 1 1	0	0
9	I	2	Total Mn 2 2	0	0
9	J	2	Total Mn 2 2	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	16	Total O 16 16	0	0
10	B	14	Total O 14 14	0	0
10	C	13	Total O 13 13	0	0
10	D	8	Total O 8 8	0	0
10	E	27	Total O 27 27	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
10	F	17	Total O 17 17	0	0
10	G	15	Total O 15 15	0	0
10	H	2	Total O 2 2	0	0
10	I	1	Total O 1 1	0	0
10	J	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3.1




- Molecule 1: Histone H3.1




- Molecule 2: Histone H4




- Molecule 3: Histone H2A type 1-B/E



- Molecule 3: Histone H2A type 1-B/E





- Molecule 4: Histone H2B type 1-J




- Molecule 4: Histone H2B type 1-J




- Molecule 5: Histone H4



- Molecule 6: DNA (145-MER)



- Molecule 7: DNA (145-MER)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.56Å 109.71Å 183.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.16 – 1.99 94.16 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.6 (94.16-1.99) 99.6 (94.16-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.31 (at 1.98Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.236 , 0.263 0.242 , 0.266	Depositor DCC
$R_{free}$ test set	3035 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.4	Xtrriage
Anisotropy	0.389	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.017 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12177	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/819	0.90	0/1097
1	E	0.77	0/818	0.93	0/1097
2	B	0.73	0/660	0.84	0/883
3	C	0.75	0/838	0.93	0/1129
3	G	0.69	0/838	0.87	0/1129
4	D	0.75	0/757	0.94	0/1015
4	H	0.69	0/757	0.85	0/1015
5	F	0.81	0/711	0.92	0/948
6	I	0.38	0/3332	1.12	39/5141 (0.8%)
7	J	0.37	0/3330	1.08	32/5138 (0.6%)
All	All	0.58	0/12860	1.02	71/18592 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	I	0	1

There are no bond length outliers.

The worst 5 of 71 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	J	14	DC	C2'-C3'-O3'	-10.50	95.75	111.50
6	I	-50	DT	N1-C1'-C2'	10.25	128.88	113.50
6	I	-28	DT	C2'-C3'-O3'	-9.94	96.59	111.50
7	J	30	DT	C2'-C3'-O3'	-9.49	97.27	111.50
7	J	54	DT	C4'-C3'-O3'	-9.03	96.45	110.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	I	-50	DT	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	807	0	844	6	0
1	E	806	0	844	6	0
2	B	653	0	696	4	0
3	C	828	0	892	3	0
3	G	828	0	892	2	0
4	D	746	0	771	3	0
4	H	746	0	771	1	0
5	F	703	0	755	0	0
6	I	2970	0	1640	69	0
7	J	2969	0	1641	44	0
8	C	1	0	0	0	0
8	G	1	0	0	0	0
9	D	1	0	0	0	0
9	I	2	0	0	0	0
9	J	2	0	0	0	0
10	A	16	0	0	0	0
10	B	14	0	0	0	0
10	C	13	0	0	0	0
10	D	8	0	0	0	0
10	E	27	0	0	1	0
10	F	17	0	0	0	0
10	G	15	0	0	0	0
10	H	2	0	0	0	0
10	I	1	0	0	0	0
10	J	1	0	0	0	0
All	All	12177	0	9746	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:I:-48:DC:H2''	6:I:-47:DC:C6	1.86	1.08
1:A:125:GLN:O	1:A:134:ARG:NH1	1.87	1.07
6:I:-40:DG:H2''	6:I:-39:DT:H5''	1.50	0.93
6:I:51:DG:H2''	6:I:52:DT:H5''	1.50	0.92
6:I:33:DC:H2''	6:I:34:DC:O4'	1.84	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/98 (98%)	96 (100%)	0	0	100	100
1	E	96/98 (98%)	95 (99%)	1 (1%)	0	100	100
2	B	80/82 (98%)	80 (100%)	0	0	100	100
3	C	105/107 (98%)	103 (98%)	2 (2%)	0	100	100
3	G	105/107 (98%)	102 (97%)	3 (3%)	0	100	100
4	D	93/95 (98%)	92 (99%)	0	1 (1%)	11	7
4	H	93/95 (98%)	90 (97%)	2 (2%)	1 (1%)	11	7
5	F	85/87 (98%)	82 (96%)	2 (2%)	1 (1%)	10	6
All	All	753/769 (98%)	740 (98%)	10 (1%)	3 (0%)	30	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	101	GLY
4	H	101	GLY
5	F	18	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	81 (95%)	4 (5%)	23	22
1	E	85/85 (100%)	82 (96%)	3 (4%)	32	32
2	B	67/67 (100%)	64 (96%)	3 (4%)	24	23
3	C	85/85 (100%)	79 (93%)	6 (7%)	13	10
3	G	85/85 (100%)	81 (95%)	4 (5%)	23	22
4	D	81/81 (100%)	78 (96%)	3 (4%)	30	30
4	H	81/81 (100%)	79 (98%)	2 (2%)	42	45
5	F	72/72 (100%)	66 (92%)	6 (8%)	10	7
All	All	641/641 (100%)	610 (95%)	31 (5%)	23	21

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	60	ASN
3	G	118	LYS
1	E	59	GLU
4	H	45	VAL
5	F	93	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	24	GLN
5	F	93	GLN
4	D	44	GLN
3	C	38	ASN
1	E	76	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	98/98 (100%)	0.91	12 (12%) 8 7	36, 51, 81, 126	0
1	E	98/98 (100%)	0.41	7 (7%) 22 20	31, 41, 66, 99	0
2	B	82/82 (100%)	0.70	6 (7%) 21 19	40, 49, 69, 121	0
3	C	107/107 (100%)	0.60	9 (8%) 17 15	31, 47, 79, 118	0
3	G	107/107 (100%)	0.93	13 (12%) 8 8	38, 53, 87, 137	0
4	D	95/95 (100%)	0.80	11 (11%) 9 8	37, 50, 94, 120	0
4	H	95/95 (100%)	0.87	12 (12%) 8 7	40, 54, 94, 143	0
5	F	87/87 (100%)	0.25	3 (3%) 48 47	31, 41, 63, 132	0
6	I	145/145 (100%)	0.79	1 (0%) 84 84	50, 101, 149, 214	0
7	J	145/145 (100%)	0.76	3 (2%) 63 63	54, 102, 149, 202	0
All	All	1059/1059 (100%)	0.71	77 (7%) 21 19	31, 55, 132, 214	0

The worst 5 of 77 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	ALA	7.3
1	E	135	ALA	5.6
1	A	38	PRO	5.2
1	E	39	HIS	4.9
1	E	38	PRO	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	MN	I	102	1/1	0.88	0.18	81,81,81,81	0
9	MN	J	102	1/1	0.88	0.17	80,80,80,80	0
9	MN	J	101	1/1	0.89	0.16	81,81,81,81	0
8	CL	G	201	1/1	0.93	0.10	59,59,59,59	0
9	MN	I	101	1/1	0.94	0.12	84,84,84,84	0
8	CL	C	201	1/1	0.94	0.09	60,60,60,60	0
9	MN	D	201	1/1	0.99	0.16	55,55,55,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.