



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:17 PM UTC

PDB ID : 9IPL / pdb_00009ipl
Title : A tetrapyrrole binding domain variant of CoaR in closed conformation at 2.28 angstrom resolution
Authors : Liu, X.C.
Deposited on : 2024-07-10
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	
1	B	218	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mercuric resistance operon regulatory protein precorrin isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1469	934	260	269	6			
1	B	198	Total	C	N	O	S	0	0	0
			1474	937	262	269	6			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	152	GLY	-	expression tag	UNP Q55938
A	153	PRO	-	expression tag	UNP Q55938
A	177	ASP	ALA	conflict	UNP Q55938
A	256	THR	GLY	conflict	UNP Q55938
A	257	ALA	LEU	conflict	UNP Q55938
A	281	SER	CYS	conflict	UNP Q55938
A	321	ALA	GLY	conflict	UNP Q55938
A	349	ALA	THR	conflict	UNP Q55938
A	359	GLU	ALA	conflict	UNP Q55938
A	363	SER	CYS	conflict	UNP Q55938
A	365	SER	CYS	conflict	UNP Q55938
A	369	SER	CYS	conflict	UNP Q55938
B	152	GLY	-	expression tag	UNP Q55938
B	153	PRO	-	expression tag	UNP Q55938
B	177	ASP	ALA	conflict	UNP Q55938
B	256	THR	GLY	conflict	UNP Q55938
B	257	ALA	LEU	conflict	UNP Q55938
B	281	SER	CYS	conflict	UNP Q55938
B	321	ALA	GLY	conflict	UNP Q55938
B	349	ALA	THR	conflict	UNP Q55938
B	359	GLU	ALA	conflict	UNP Q55938
B	363	SER	CYS	conflict	UNP Q55938
B	365	SER	CYS	conflict	UNP Q55938
B	369	SER	CYS	conflict	UNP Q55938

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		
3	B	57	Total	O	0	0
			57	57		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mercuric resistance operon regulatory protein precorrin isomerase



- Molecule 1: Mercuric resistance operon regulatory protein precorrin isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	96.04Å 96.04Å 148.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.74 – 2.28 45.74 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.74-2.28) 99.8 (45.74-2.28)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.209 , 0.230 0.215 , 0.232	Depositor DCC
R_{free} test set	1733 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	48.6	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.044 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3091	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1498	0.93	1/2050 (0.0%)
1	B	0.53	0/1504	0.94	1/2058 (0.0%)
All	All	0.54	0/3002	0.94	2/4108 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	225	ASP	CA-CB-CG	5.75	118.35	112.60
1	A	225	ASP	CA-CB-CG	5.11	117.71	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1469	0	1499	7	0
1	B	1474	0	1504	7	0
2	A	20	0	0	0	0
2	B	20	0	0	0	0
3	A	51	0	0	0	0
3	B	57	0	0	0	0
All	All	3091	0	3003	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314:ILE:H	1:B:314:ILE:HD12	1.72	0.55
1:A:314:ILE:HD12	1:A:314:ILE:H	1.72	0.54
1:A:194:GLY:HA3	1:B:339:SER:O	2.11	0.50
1:A:293:LEU:HD22	1:A:320:PRO:HB3	1.95	0.47
1:A:339:SER:O	1:B:194:GLY:HA3	2.14	0.47
1:B:312:MET:N	1:B:313:PRO:CD	2.77	0.47
1:A:312:MET:N	1:A:313:PRO:CD	2.78	0.47
1:B:221:PRO:HG2	1:B:278:PRO:HG2	1.97	0.46
1:B:172:LEU:N	1:B:173:PRO:CD	2.79	0.46
1:A:172:LEU:N	1:A:173:PRO:CD	2.79	0.45
1:A:203:ARG:NH1	1:A:337:GLN:OE1	2.51	0.42
1:B:293:LEU:HD22	1:B:320:PRO:HB3	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/218 (90%)	186 (95%)	10 (5%)	0	100	100
1	B	196/218 (90%)	186 (95%)	10 (5%)	0	100	100
All	All	392/436 (90%)	372 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/178 (88%)	155 (99%)	1 (1%)	78	87
1	B	157/178 (88%)	156 (99%)	1 (1%)	78	87
All	All	313/356 (88%)	311 (99%)	2 (1%)	78	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	324	ARG
1	B	170	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	GLN
1	B	272	GLN
1	B	275	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	403	-	4,4,4	0.33	0	6,6,6	0.13	0
2	SO4	A	401	-	4,4,4	0.33	0	6,6,6	0.08	0
2	SO4	B	401	-	4,4,4	0.32	0	6,6,6	0.07	0
2	SO4	B	404	-	4,4,4	0.24	0	6,6,6	0.14	0
2	SO4	B	403	-	4,4,4	0.34	0	6,6,6	0.18	0
2	SO4	B	402	-	4,4,4	0.36	0	6,6,6	0.15	0
2	SO4	A	402	-	4,4,4	0.31	0	6,6,6	0.10	0
2	SO4	A	404	-	4,4,4	0.32	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/218 (90%)	0.73	26 (13%) 7 8	33, 50, 92, 114	0
1	B	198/218 (90%)	0.71	25 (12%) 8 8	34, 49, 88, 114	0
All	All	396/436 (90%)	0.72	51 (12%) 7 8	33, 50, 90, 114	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	164	PHE	8.4
1	A	161	PRO	7.0
1	A	164	PHE	6.1
1	B	161	PRO	6.0
1	A	163	ALA	5.1
1	B	163	ALA	4.8
1	A	319	ALA	4.7
1	A	265	TRP	4.6
1	A	269	HIS	4.5
1	A	165	GLN	4.0
1	A	321	ALA	4.0
1	B	265	TRP	4.0
1	B	358	ILE	3.9
1	B	319	ALA	3.9
1	B	165	GLN	3.7
1	B	168	LEU	3.6
1	A	358	ILE	3.5
1	A	266	HIS	3.5
1	A	162	GLU	3.5
1	B	266	HIS	3.4
1	B	162	GLU	3.2
1	B	199	VAL	3.2
1	B	170	HIS	3.2
1	A	318	HIS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	320	PRO	3.1
1	A	170	HIS	3.1
1	A	167	SER	3.1
1	A	168	LEU	2.9
1	A	236	THR	2.9
1	B	318	HIS	2.8
1	B	166	GLU	2.8
1	A	214	ASP	2.7
1	B	169	LYS	2.7
1	A	317	SER	2.7
1	A	169	LYS	2.6
1	B	198	LEU	2.6
1	A	197	SER	2.5
1	B	321	ALA	2.5
1	B	269	HIS	2.4
1	A	199	VAL	2.4
1	B	240	HIS	2.3
1	B	210	ALA	2.3
1	A	324	ARG	2.2
1	B	197	SER	2.2
1	B	206	GLN	2.2
1	A	262	GLN	2.1
1	A	198	LEU	2.1
1	A	316	PHE	2.1
1	B	167	SER	2.1
1	A	327	MET	2.0
1	B	200	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	402	5/5	0.71	0.12	107,109,122,133	0
2	SO4	B	403	5/5	0.81	0.13	87,92,93,98	0
2	SO4	A	401	5/5	0.85	0.11	98,99,106,111	0
2	SO4	B	401	5/5	0.86	0.11	91,98,102,106	0
2	SO4	A	403	5/5	0.86	0.12	81,89,107,108	0
2	SO4	B	404	5/5	0.86	0.15	59,75,84,96	0
2	SO4	A	404	5/5	0.87	0.15	60,74,94,99	0
2	SO4	B	402	5/5	0.92	0.10	72,75,83,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.