



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:41 AM UTC

PDB ID : 4IQX / pdb_00004iqx
Title : Mutant P44S P169S M296I of Foot-and-mouth disease Virus RNA-dependent RNA polymerase
Authors : Ferrer-Orta, C.; Verdaguer, N.; Agudo, R.; Perez-Luque, R.
Deposited on : 2013-01-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)
RNA backbone	3983	1003 (2.78-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	13% 90% 8% ..
2	B	7	100% 71% 14% 14%
3	C	5	80% 80% 20%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3D polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			3677	2344	641	673	19			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	SER	PRO	engineered mutation	UNP Q9QCE4
A	169	SER	PRO	engineered mutation	UNP Q9QCE4
A	296	ILE	MET	engineered mutation	UNP Q9QCE4
A	471	ALA	-	expression tag	UNP Q9QCE4
A	472	ALA	-	expression tag	UNP Q9QCE4
A	473	LEU	-	expression tag	UNP Q9QCE4
A	474	ALA	-	expression tag	UNP Q9QCE4
A	475	HIS	-	expression tag	UNP Q9QCE4
A	476	HIS	-	expression tag	UNP Q9QCE4
A	477	HIS	-	expression tag	UNP Q9QCE4
A	478	HIS	-	expression tag	UNP Q9QCE4
A	479	HIS	-	expression tag	UNP Q9QCE4
A	480	HIS	-	expression tag	UNP Q9QCE4
A	481	HIS	-	expression tag	UNP Q9QCE4

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*UP*GP*GP*GP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	P	0	0	0
			148	67	28	47	6			

- Molecule 3 is a RNA chain called RNA (5'-R(*GP*GP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	5	Total	C	N	O	P	0	0	0
			103	47	19	33	4			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	35	Total 35	O 35	0	0
4	B	4	Total 4	O 4	0	0
4	C	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3D polymerase



- Molecule 2: RNA (5'-R(*AP*UP*GP*GP*GP*CP*C)-3')



- Molecule 3: RNA (5'-R(*GP*GP*CP*CP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.84Å 93.84Å 99.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.36 – 2.50 29.36 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.36-2.50) 99.2 (29.36-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.83 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.222 , 0.254 0.267 , 0.284	Depositor DCC
R_{free} test set	915 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 28.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3969	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3765	0.75	0/5107
2	B	0.47	0/165	0.84	0/256
3	C	0.53	0/114	0.72	0/176
All	All	0.41	0/4044	0.75	0/5539

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3677	0	3573	30	0
2	B	148	0	78	4	0
3	C	103	0	57	1	0
4	A	35	0	0	10	0
4	B	4	0	0	0	0
4	C	2	0	0	0	0
All	All	3969	0	3708	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:CYS:SG	4:A:513:HOH:O	2.50	0.69
1:A:244:PHE:CZ	4:A:517:HOH:O	2.46	0.68
1:A:427:VAL:HB	4:A:514:HOH:O	1.95	0.66
1:A:333:MET:HE3	1:A:340:ILE:HD13	1.80	0.63
1:A:116:ALA:H	2:B:904:U:H5'	1.67	0.60
1:A:317:TYR:HA	4:A:519:HOH:O	2.02	0.59
1:A:320:ARG:HB2	4:A:519:HOH:O	2.03	0.59
1:A:333:MET:HE3	1:A:340:ILE:CD1	2.33	0.59
1:A:387:LYS:HD3	3:C:919:C:H5''	1.86	0.57
1:A:244:PHE:HZ	4:A:517:HOH:O	1.84	0.57
1:A:132:ILE:HG12	1:A:139:VAL:HG12	1.89	0.54
1:A:396:THR:HB	1:A:398:PHE:HD2	1.75	0.52
1:A:244:PHE:HA	1:A:362:GLN:HE22	1.75	0.52
1:A:321:ARG:HH11	1:A:356:HIS:HD2	1.59	0.50
1:A:396:THR:HB	1:A:398:PHE:CD2	2.48	0.49
1:A:181:VAL:HG11	2:B:904:U:H1'	1.94	0.48
1:A:33:GLU:H	1:A:33:GLU:CD	2.23	0.45
1:A:308:THR:HG23	1:A:335:SER:OG	2.16	0.45
1:A:424:LEU:HA	4:A:514:HOH:O	2.17	0.44
2:B:904:U:H2'	2:B:905:G:C8	2.52	0.44
1:A:433:HIS:HE2	1:A:462:TRP:CD1	2.36	0.44
1:A:203:MET:HE3	1:A:211:ILE:HD12	2.01	0.43
1:A:310:LEU:CB	4:A:517:HOH:O	2.66	0.43
1:A:96:LEU:HD22	1:A:197:GLY:HA3	2.00	0.43
1:A:73:LEU:HG	4:A:513:HOH:O	2.18	0.42
1:A:224:GLN:CD	1:A:398:PHE:HD1	2.28	0.42
1:A:244:PHE:CD2	1:A:338:ASP:HB3	2.54	0.42
1:A:96:LEU:HD13	1:A:198:ARG:CG	2.50	0.41
1:A:369:LYS:HG3	1:A:370:SER:H	1.85	0.41
1:A:321:ARG:HB2	4:A:532:HOH:O	2.20	0.41
1:A:115:THR:HB	2:B:904:U:H5''	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	474/481 (98%)	463 (98%)	10 (2%)	1 (0%)	43 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	379/403 (94%)	371 (98%)	8 (2%)	47 74

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	GLU
1	A	96	LEU
1	A	174	ARG
1	A	221	VAL
1	A	334	ILE
1	A	340	ILE
1	A	368	ASP
1	A	392	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	HIS
1	A	160	GLN
1	A	232	GLN
1	A	280	ASN
1	A	311	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	356	HIS
1	A	362	GLN
1	A	389	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	6/7 (85%)	1 (16%)	0
3	C	4/5 (80%)	0	0
All	All	10/12 (83%)	1 (10%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	904	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/481 (98%)	1.16	62 (13%) 7 6	35, 43, 54, 59	0
2	B	7/7 (100%)	3.61	7 (100%) 0 0	55, 110, 118, 120	1 (14%)
3	C	5/5 (100%)	4.12	4 (80%) 0 0	46, 94, 95, 97	1 (20%)
All	All	488/493 (98%)	1.22	73 (14%) 5 4	35, 43, 55, 120	2 (0%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	919	C	8.5
2	B	906	G	8.5
1	A	473	LEU	4.6
1	A	16	MET	4.5
3	C	916	G	4.5
1	A	287	ASN	4.4
1	A	149	LEU	4.0
1	A	469	ASP	3.7
1	A	447	GLN	3.7
1	A	476	HIS	3.7
1	A	370	SER	3.5
2	B	903	A	3.4
1	A	9	VAL	3.3
1	A	468	GLY	3.3
2	B	907	G	3.3
1	A	145	ALA	3.3
3	C	920	C	3.2
1	A	475	HIS	3.2
1	A	437	ASP	3.1
1	A	120	PRO	3.1
1	A	46	LEU	3.1
1	A	472	ALA	3.0
1	A	329	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	136	ASN	3.0
1	A	146	ALA	3.0
1	A	28	GLY	3.0
1	A	299	GLY	2.9
1	A	139	VAL	2.9
2	B	909	C	2.8
1	A	12	ARG	2.8
1	A	137	GLY	2.7
1	A	181	VAL	2.7
1	A	368	ASP	2.7
1	A	125	GLY	2.7
1	A	10	GLU	2.7
1	A	367	ALA	2.7
2	B	905	G	2.6
1	A	352	ALA	2.6
1	A	14	HIS	2.6
1	A	41	ASN	2.6
1	A	371	ASP	2.6
1	A	467	CYS	2.5
1	A	470	ALA	2.5
1	A	124	GLN	2.5
3	C	917	G	2.5
1	A	53	ASP	2.5
1	A	25	VAL	2.4
2	B	904	U	2.4
1	A	144	GLU	2.4
1	A	47	ASN	2.4
1	A	88	VAL	2.3
1	A	369	LYS	2.3
1	A	263	THR	2.3
1	A	349	ASP	2.3
1	A	147	LEU	2.3
1	A	118	GLY	2.2
1	A	364	ILE	2.2
2	B	908	C	2.2
1	A	471	ALA	2.2
1	A	154	GLU	2.2
1	A	17	ARG	2.2
1	A	350	PHE	2.2
1	A	5	ASP	2.1
1	A	162	PHE	2.1
1	A	64	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	ALA	2.1
1	A	11	GLU	2.1
1	A	372	LYS	2.1
1	A	49	GLY	2.1
1	A	43	ASP	2.1
1	A	474	ALA	2.0
1	A	90	GLY	2.0
1	A	308	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.