



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:02 PM UTC

PDB ID : 8IQ7 / pdb_00008iq7
Title : Ambient Temperature Crystal Structure of Candida boidinii Formate Dehydrogenase
Authors : Gul, M.; DeMirici, H.
Deposited on : 2023-03-16
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	4% 93% 5%
1	B	364	4% 91% 6%
1	C	364	5% 91% 6%
1	D	364	6% 93% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 22547 atoms, of which 11053 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	356	Total	C	H	N	O	S	0	1	0
			5568	1775	2782	477	527	7			
1	B	353	Total	C	H	N	O	S	0	2	0
			5528	1764	2759	473	525	7			
1	C	354	Total	C	H	N	O	S	0	1	0
			5502	1763	2734	474	524	7			
1	D	356	Total	C	H	N	O	S	0	0	0
			5556	1771	2778	476	524	7			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	97	Total	O	0	0
			97	97		
2	B	99	Total	O	0	0
			99	99		
2	C	107	Total	O	0	0
			107	107		
2	D	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Formate dehydrogenase




- Molecule 1: Formate dehydrogenase




- Molecule 1: Formate dehydrogenase



- Molecule 1: Formate dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.67Å 69.51Å 110.80Å 77.86° 89.05° 81.20°	Depositor
Resolution (Å)	30.70 – 2.10 30.70 – 2.10	Depositor EDS
% Data completeness (in resolution range)	73.1 (30.70-2.10) 73.1 (30.70-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.243 , 0.272 0.244 , 0.272	Depositor DCC
R_{free} test set	2000 reflections (2.18%)	wwPDB-VP
Wilson B-factor (Å ²)	12.6	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	22547	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/2843	0.28	0/3857
1	B	0.09	0/2829	0.28	0/3838
1	C	0.11	0/2825	0.30	0/3833
1	D	0.10	0/2835	0.29	0/3846
All	All	0.10	0/11332	0.29	0/15374

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2786	2782	2792	11	1
1	B	2769	2759	2769	9	0
1	C	2768	2734	2765	10	0
1	D	2778	2778	2789	8	1
2	A	97	0	0	5	0
2	B	99	0	0	3	0
2	C	107	0	0	4	0
2	D	90	0	0	5	0
All	All	11494	11053	11115	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:272[A]:GLU:OE2	2:C:401:HOH:O	2.02	0.76
1:C:10:ALA:HB3	1:C:14:ALA:HB2	1.69	0.74
1:A:154:ALA:O	2:A:401:HOH:O	2.07	0.72
1:D:350:ASN:OD1	2:D:401:HOH:O	2.08	0.72
1:D:277:ARG:NH1	2:D:404:HOH:O	2.22	0.70
1:C:208:GLY:O	2:C:402:HOH:O	2.11	0.68
1:B:170:ILE:HD12	1:B:227:ILE:HD13	1.76	0.66
1:B:85:LEU:N	2:B:406:HOH:O	2.30	0.64
1:A:82:ALA:HB1	1:A:85:LEU:HB2	1.84	0.59
1:D:297:ASP:OD2	2:D:402:HOH:O	2.17	0.59
1:B:10:ALA:HB3	1:B:14:ALA:HB2	1.84	0.58
1:A:6:VAL:HG11	1:A:72:ALA:HB2	1.87	0.57
1:B:25:GLU:OE1	1:B:25:GLU:N	2.37	0.56
1:D:45:SER:O	2:D:403:HOH:O	2.18	0.56
1:A:332:GLU:OE1	2:A:403:HOH:O	2.17	0.55
1:A:178:ARG:NH1	2:A:414:HOH:O	2.41	0.54
1:A:149[B]:ASP:OD1	1:A:150:TRP:N	2.39	0.54
1:C:17:GLU:HG3	2:C:420:HOH:O	2.08	0.53
1:C:25:GLU:N	1:C:25:GLU:OE1	2.41	0.53
1:B:82:ALA:HB1	1:B:85:LEU:HB2	1.94	0.49
1:C:158:ASP:OD2	2:C:403:HOH:O	2.19	0.49
1:B:219:VAL:O	2:B:401:HOH:O	2.20	0.49
1:C:181:GLU:OE2	1:C:206:LYS:NZ	2.47	0.48
1:A:51:ASN:OD1	1:A:51:ASN:N	2.47	0.48
1:C:227:ILE:HD11	1:C:239:ILE:HG13	1.98	0.46
1:A:9:ASP:OD1	1:A:24:THR:OG1	2.33	0.46
1:C:317:LEU:HD12	1:D:160:TYR:CZ	2.53	0.44
1:B:314:GLY:O	1:B:323:TYR:OH	2.23	0.43
1:A:195:ASP:OD1	1:A:196:TYR:N	2.51	0.43
1:A:356:LYS:NZ	2:A:415:HOH:O	2.44	0.42
1:B:6:VAL:HG11	1:B:72:ALA:HB2	2.02	0.42
1:D:27:LYS:NZ	2:D:413:HOH:O	2.47	0.42
1:D:118:SER:HG	1:D:323:TYR:HE1	1.64	0.42
1:A:352:GLU:OE2	2:A:404:HOH:O	2.22	0.41
1:C:21:TYR:HB3	1:C:28:LEU:HA	2.02	0.41
1:D:6:VAL:HG11	1:D:72:ALA:HB2	2.02	0.40
1:B:147:ASN:ND2	2:B:420:HOH:O	2.45	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:355:THR:H	1:D:355:THR:OG1[1_645]	1.55	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/364 (98%)	349 (98%)	6 (2%)	0	100	100
1	B	353/364 (97%)	344 (98%)	9 (2%)	0	100	100
1	C	353/364 (97%)	345 (98%)	8 (2%)	0	100	100
1	D	354/364 (97%)	345 (98%)	9 (2%)	0	100	100
All	All	1415/1456 (97%)	1383 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/298 (98%)	290 (99%)	3 (1%)	68	76
1	B	291/298 (98%)	282 (97%)	9 (3%)	35	39
1	C	290/298 (97%)	283 (98%)	7 (2%)	43	49
1	D	292/298 (98%)	287 (98%)	5 (2%)	53	62
All	All	1166/1192 (98%)	1142 (98%)	24 (2%)	47	54

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	ILE
1	A	109	LYS
1	A	356	LYS
1	B	12	LYS
1	B	35	LYS
1	B	37	GLN
1	B	48	GLU
1	B	74	ILE
1	B	128	LEU
1	B	215	ILE
1	B	248	LYS
1	B	292	ASP
1	C	12	LYS
1	C	128	LEU
1	C	143	GLU
1	C	147	ASN
1	C	205	GLU
1	C	213	GLU
1	C	347	ILE
1	D	4	VAL
1	D	18	GLU
1	D	178	ARG
1	D	204	GLU
1	D	356	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	228	ASN
1	B	126	HIS
1	B	214	ASN
1	C	51	ASN
1	C	350	ASN
1	D	350	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	356/364 (97%)	0.59	15 (4%)	40 42	8, 17, 39, 65	1 (0%)
1	B	353/364 (96%)	0.69	16 (4%)	38 40	8, 19, 42, 68	2 (0%)
1	C	354/364 (97%)	0.65	18 (5%)	33 35	9, 18, 38, 64	1 (0%)
1	D	356/364 (97%)	0.67	23 (6%)	25 26	7, 18, 42, 66	0
All	All	1419/1456 (97%)	0.65	72 (5%)	33 35	7, 18, 41, 68	4 (0%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	15	ALA	4.5
1	C	354	ILE	4.4
1	D	354	ILE	4.2
1	C	14	ALA	4.2
1	B	51	ASN	4.2
1	B	50	GLY	4.1
1	B	38	GLY	3.9
1	D	48	GLU	3.8
1	D	337	GLY	3.5
1	A	58	ILE	3.4
1	C	16	ASP	3.2
1	D	128	LEU	3.2
1	B	108	LYS	3.1
1	C	213	GLU	3.0
1	D	50	GLY	3.0
1	A	213	GLU	2.9
1	B	102	TYR	2.9
1	C	18	GLU	2.9
1	A	206	LYS	2.9
1	B	149	ASP	2.9
1	D	51	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	355	THR	2.9
1	C	49	GLY	2.9
1	B	128	LEU	2.8
1	A	356	LYS	2.8
1	A	355	THR	2.8
1	B	19	LYS	2.8
1	A	354	ILE	2.8
1	C	10	ALA	2.7
1	A	149[A]	ASP	2.7
1	D	202	ASP	2.7
1	D	49	GLY	2.7
1	A	215	ILE	2.7
1	D	35	LYS	2.6
1	B	82	ALA	2.5
1	D	38	GLY	2.5
1	C	272[A]	GLU	2.4
1	D	11	GLY	2.4
1	D	215	ILE	2.3
1	C	48	GLU	2.3
1	C	197	GLN	2.3
1	D	334	PHE	2.2
1	D	152	VAL	2.2
1	B	351	GLY	2.2
1	A	295	TRP	2.2
1	D	241	LYS	2.2
1	A	202	ASP	2.2
1	C	50	GLY	2.2
1	C	323	TYR	2.2
1	A	18	GLU	2.2
1	A	49	GLY	2.2
1	B	346	ILE	2.1
1	D	197	GLN	2.1
1	C	13	HIS	2.1
1	C	270	ALA	2.1
1	D	246	LYS	2.1
1	B	48	GLU	2.1
1	C	284	TRP	2.1
1	B	12	LYS	2.1
1	D	216	GLU	2.1
1	B	36	ASP	2.1
1	D	37	GLN	2.1
1	A	263	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	19	LYS	2.1
1	B	21	TYR	2.1
1	A	82	ALA	2.1
1	C	209	ALA	2.1
1	A	55	ASP	2.0
1	B	101	ASP	2.0
1	D	20	LEU	2.0
1	D	356	LYS	2.0
1	C	11	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.