



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:26 PM UTC

PDB ID : 4IRB / pdb_00004irb
Title : Crystal Structure of Vaccinia Virus Uracil DNA Glycosylase Mutant del171-172D4
Authors : Schormann, N.; Zhukovskaya, N.; Sartmatova, D.; Nuth, M.; Ricciardi, R.P.; Chattopadhyay, D.
Deposited on : 2013-01-14
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	
1	B	236	
1	C	236	
1	D	236	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	302	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uracil-DNA glycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1756	1139	291	320	6			
1	B	216	Total	C	N	O	S	0	0	0
			1734	1126	287	315	6			
1	C	215	Total	C	N	O	S	0	0	0
			1731	1126	284	315	6			
1	D	214	Total	C	N	O	S	0	0	0
			1721	1116	284	315	6			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP Q91UM2
A	-18	GLY	-	expression tag	UNP Q91UM2
A	-17	SER	-	expression tag	UNP Q91UM2
A	-16	SER	-	expression tag	UNP Q91UM2
A	-15	HIS	-	expression tag	UNP Q91UM2
A	-14	HIS	-	expression tag	UNP Q91UM2
A	-13	HIS	-	expression tag	UNP Q91UM2
A	-12	HIS	-	expression tag	UNP Q91UM2
A	-11	HIS	-	expression tag	UNP Q91UM2
A	-10	HIS	-	expression tag	UNP Q91UM2
A	-9	SER	-	expression tag	UNP Q91UM2
A	-8	SER	-	expression tag	UNP Q91UM2
A	-7	GLY	-	expression tag	UNP Q91UM2
A	-6	LEU	-	expression tag	UNP Q91UM2
A	-5	VAL	-	expression tag	UNP Q91UM2
A	-4	PRO	-	expression tag	UNP Q91UM2
A	-3	ARG	-	expression tag	UNP Q91UM2
A	-2	GLY	-	expression tag	UNP Q91UM2
A	-1	SER	-	expression tag	UNP Q91UM2
A	0	HIS	-	expression tag	UNP Q91UM2
A	17	ASN	ASP	engineered mutation	UNP Q91UM2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLU	deletion	UNP Q91UM2
A	?	-	SER	deletion	UNP Q91UM2
B	-19	MET	-	initiating methionine	UNP Q91UM2
B	-18	GLY	-	expression tag	UNP Q91UM2
B	-17	SER	-	expression tag	UNP Q91UM2
B	-16	SER	-	expression tag	UNP Q91UM2
B	-15	HIS	-	expression tag	UNP Q91UM2
B	-14	HIS	-	expression tag	UNP Q91UM2
B	-13	HIS	-	expression tag	UNP Q91UM2
B	-12	HIS	-	expression tag	UNP Q91UM2
B	-11	HIS	-	expression tag	UNP Q91UM2
B	-10	HIS	-	expression tag	UNP Q91UM2
B	-9	SER	-	expression tag	UNP Q91UM2
B	-8	SER	-	expression tag	UNP Q91UM2
B	-7	GLY	-	expression tag	UNP Q91UM2
B	-6	LEU	-	expression tag	UNP Q91UM2
B	-5	VAL	-	expression tag	UNP Q91UM2
B	-4	PRO	-	expression tag	UNP Q91UM2
B	-3	ARG	-	expression tag	UNP Q91UM2
B	-2	GLY	-	expression tag	UNP Q91UM2
B	-1	SER	-	expression tag	UNP Q91UM2
B	0	HIS	-	expression tag	UNP Q91UM2
B	17	ASN	ASP	engineered mutation	UNP Q91UM2
B	?	-	GLU	deletion	UNP Q91UM2
B	?	-	SER	deletion	UNP Q91UM2
C	-19	MET	-	initiating methionine	UNP Q91UM2
C	-18	GLY	-	expression tag	UNP Q91UM2
C	-17	SER	-	expression tag	UNP Q91UM2
C	-16	SER	-	expression tag	UNP Q91UM2
C	-15	HIS	-	expression tag	UNP Q91UM2
C	-14	HIS	-	expression tag	UNP Q91UM2
C	-13	HIS	-	expression tag	UNP Q91UM2
C	-12	HIS	-	expression tag	UNP Q91UM2
C	-11	HIS	-	expression tag	UNP Q91UM2
C	-10	HIS	-	expression tag	UNP Q91UM2
C	-9	SER	-	expression tag	UNP Q91UM2
C	-8	SER	-	expression tag	UNP Q91UM2
C	-7	GLY	-	expression tag	UNP Q91UM2
C	-6	LEU	-	expression tag	UNP Q91UM2
C	-5	VAL	-	expression tag	UNP Q91UM2
C	-4	PRO	-	expression tag	UNP Q91UM2
C	-3	ARG	-	expression tag	UNP Q91UM2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP Q91UM2
C	-1	SER	-	expression tag	UNP Q91UM2
C	0	HIS	-	expression tag	UNP Q91UM2
C	17	ASN	ASP	engineered mutation	UNP Q91UM2
C	?	-	GLU	deletion	UNP Q91UM2
C	?	-	SER	deletion	UNP Q91UM2
D	-19	MET	-	initiating methionine	UNP Q91UM2
D	-18	GLY	-	expression tag	UNP Q91UM2
D	-17	SER	-	expression tag	UNP Q91UM2
D	-16	SER	-	expression tag	UNP Q91UM2
D	-15	HIS	-	expression tag	UNP Q91UM2
D	-14	HIS	-	expression tag	UNP Q91UM2
D	-13	HIS	-	expression tag	UNP Q91UM2
D	-12	HIS	-	expression tag	UNP Q91UM2
D	-11	HIS	-	expression tag	UNP Q91UM2
D	-10	HIS	-	expression tag	UNP Q91UM2
D	-9	SER	-	expression tag	UNP Q91UM2
D	-8	SER	-	expression tag	UNP Q91UM2
D	-7	GLY	-	expression tag	UNP Q91UM2
D	-6	LEU	-	expression tag	UNP Q91UM2
D	-5	VAL	-	expression tag	UNP Q91UM2
D	-4	PRO	-	expression tag	UNP Q91UM2
D	-3	ARG	-	expression tag	UNP Q91UM2
D	-2	GLY	-	expression tag	UNP Q91UM2
D	-1	SER	-	expression tag	UNP Q91UM2
D	0	HIS	-	expression tag	UNP Q91UM2
D	17	ASN	ASP	engineered mutation	UNP Q91UM2
D	?	-	GLU	deletion	UNP Q91UM2
D	?	-	SER	deletion	UNP Q91UM2

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	33	Total	O	0	0
			33	33		
5	B	30	Total	O	0	0
			30	30		
5	C	27	Total	O	0	0
			27	27		
5	D	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uracil-DNA glycosylase




- Molecule 1: Uracil-DNA glycosylase



- Molecule 1: Uracil-DNA glycosylase



- Molecule 1: Uracil-DNA glycosylase



P182	
D186	
R193	
S194	
I197	
V200	
L201	
L202	
E203	
Y218	

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.90Å 130.51Å 86.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.64 – 2.30 19.64 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (19.64-2.30) 99.2 (19.64-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.73 (at 2.30Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496, REFMAC 5.7.0029	Depositor
R, R_{free}	0.211 , 0.258 (Not available) , 0.234	Depositor DCC
R_{free} test set	2616 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.323	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7087	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1804	0.85	5/2451 (0.2%)
1	B	0.40	0/1782	0.82	3/2425 (0.1%)
1	C	0.39	0/1778	0.81	0/2417
1	D	0.38	0/1767	0.81	0/2402
All	All	0.39	0/7131	0.82	8/9695 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	204	LEU	N-CA-C	-8.31	102.18	111.82
1	B	202	LEU	CA-C-N	-6.05	112.99	122.73
1	B	202	LEU	C-N-CA	-6.05	112.99	122.73
1	A	77	VAL	CA-C-N	5.58	125.90	119.93
1	A	77	VAL	C-N-CA	5.58	125.90	119.93
1	A	81	SER	CA-C-N	5.16	125.20	119.32
1	A	81	SER	C-N-CA	5.16	125.20	119.32
1	A	187	ARG	N-CA-C	-5.11	104.51	111.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1746	21	0
1	B	1734	0	1727	16	0
1	C	1731	0	1721	18	0
1	D	1721	0	1709	17	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	1	0
3	A	6	0	8	8	0
4	B	1	0	0	0	0
5	A	33	0	0	1	0
5	B	30	0	0	0	0
5	C	27	0	0	0	0
5	D	28	0	0	0	0
All	All	7087	0	6911	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:PHE:H	3:A:302:GOL:H11	1.13	1.10
1:C:197:ILE:HD11	1:D:197:ILE:HD11	1.48	0.95
1:B:149:THR:HB	1:B:174:VAL:HG21	1.65	0.78
1:A:164:SER:HA	1:B:173:PRO:HD3	1.67	0.77
1:A:79:PHE:N	3:A:302:GOL:H11	1.96	0.76
1:C:193:ARG:HH21	1:D:200:VAL:HG13	1.53	0.74
1:A:79:PHE:H	3:A:302:GOL:C1	1.99	0.72
1:A:120:ASN:HD21	3:A:302:GOL:H12	1.58	0.69
1:C:161:THR:O	1:C:164:SER:OG	2.09	0.68
1:A:129:GLU:HG2	1:A:132:SER:HB3	1.76	0.66
1:A:130:THR:HG22	1:A:131:LYS:HG3	1.80	0.64
1:C:130:THR:HG22	1:C:131:LYS:HG3	1.85	0.59
1:A:78:PRO:HB2	1:A:118:PRO:HB2	1.83	0.59
1:B:185:ARG:O	1:B:187:ARG:N	2.36	0.58
1:A:70:TYR:CG	3:A:302:GOL:H31	2.39	0.57
1:D:6:VAL:HG12	1:D:27:VAL:HG13	1.87	0.56
1:C:193:ARG:HH12	1:D:203:GLU:CD	2.13	0.55
1:C:26:LEU:HD12	1:C:144:LEU:HD21	1.90	0.54
1:D:158:LEU:HB3	1:D:182:PRO:HD3	1.90	0.53
1:A:70:TYR:CD1	3:A:302:GOL:H31	2.44	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:204:LEU:HD21	1:D:193:ARG:HB3	1.91	0.52
1:B:81:SER:HB3	1:B:89:ILE:HG21	1.91	0.52
1:B:147:HIS:O	1:B:150:LYS:HG2	2.09	0.52
1:B:36:TRP:HZ2	1:C:132:SER:O	1.92	0.52
1:B:39:ARG:NH2	1:B:136:TYR:OH	2.43	0.52
1:B:203:GLU:HA	1:B:207:LYS:H	1.75	0.52
1:A:205:ASP:OD1	1:B:160:LYS:NZ	2.39	0.51
1:A:70:TYR:CD2	3:A:302:GOL:H31	2.46	0.51
1:A:201:LEU:HD23	1:A:204:LEU:HD12	1.93	0.51
1:A:2:ASN:HB2	1:A:15:TYR:CZ	2.46	0.50
1:D:160:LYS:N	2:D:301:SO4:O4	2.36	0.50
1:D:10:PRO:HB3	1:D:47:ASP:HA	1.93	0.48
1:A:97:SER:OG	1:A:102:VAL:O	2.29	0.48
1:D:177:ILE:HD11	1:D:201:LEU:HD12	1.96	0.48
1:D:2:ASN:HB2	1:D:15:TYR:CZ	2.49	0.48
1:C:169:LYS:O	1:C:174:VAL:HG12	2.14	0.47
1:D:197:ILE:O	1:D:200:VAL:HB	2.15	0.47
1:D:194:SER:O	1:D:197:ILE:HG22	2.14	0.47
1:A:25:GLN:HB3	5:A:414:HOH:O	2.15	0.46
1:C:185:ARG:O	1:C:187:ARG:N	2.49	0.46
1:B:169:LYS:HE3	1:B:169:LYS:HB2	1.83	0.45
1:D:129:GLU:HG2	1:D:132:SER:HB3	1.98	0.45
1:A:78:PRO:HG3	1:A:120:ASN:HD22	1.82	0.45
1:C:1:MET:HE3	1:C:14:THR:HG22	1.99	0.45
1:C:72:LYS:HD3	1:C:127:LEU:HG	1.98	0.44
1:C:194:SER:O	1:C:197:ILE:HG22	2.17	0.44
1:C:10:PRO:HB3	1:C:47:ASP:HA	1.99	0.43
1:D:69:PRO:HD3	1:D:122:TYR:O	2.18	0.43
1:C:158:LEU:HB3	1:C:182:PRO:HD3	2.01	0.43
1:C:23:MET:O	1:C:27:VAL:HG23	2.19	0.43
1:B:163:PHE:HA	1:B:166:ILE:HD13	2.00	0.43
1:B:65:CYS:HA	1:B:119:TRP:O	2.19	0.42
1:C:164:SER:HB2	1:C:165:ASN:H	1.65	0.42
1:A:200:VAL:HG13	1:B:193:ARG:NH2	2.34	0.42
1:B:43:SER:HA	1:B:44:PRO:HA	1.89	0.41
1:B:177:ILE:HD11	1:B:201:LEU:CD1	2.50	0.41
1:A:67:ILE:HB	1:A:162:ASP:HB3	2.02	0.41
1:D:15:TYR:CE2	1:D:23:MET:HG3	2.56	0.41
1:C:142:LYS:HB2	1:C:142:LYS:HE3	1.49	0.41
1:A:48:LYS:HA	1:A:51:ILE:HG23	2.02	0.40
1:B:84:PHE:CD1	1:B:107:GLY:HA2	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:163:PHE:HA	1:D:166:ILE:HD13	2.03	0.40
1:A:70:TYR:CE1	3:A:302:GOL:H31	2.56	0.40
1:D:19:TRP:O	1:D:23:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/236 (90%)	206 (97%)	5 (2%)	2 (1%)	14	17
1	B	214/236 (91%)	206 (96%)	6 (3%)	2 (1%)	14	17
1	C	211/236 (89%)	203 (96%)	5 (2%)	3 (1%)	9	9
1	D	210/236 (89%)	203 (97%)	6 (3%)	1 (0%)	24	31
All	All	848/944 (90%)	818 (96%)	22 (3%)	8 (1%)	14	17

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	186	ASP
1	C	164	SER
1	C	186	ASP
1	A	185	ARG
1	B	166	ILE
1	A	138	ASP
1	C	138	ASP
1	D	186	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/214 (92%)	196 (100%)	0	100	100
1	B	192/214 (90%)	191 (100%)	1 (0%)	81	90
1	C	192/214 (90%)	190 (99%)	2 (1%)	68	82
1	D	191/214 (89%)	191 (100%)	0	100	100
All	All	771/856 (90%)	768 (100%)	3 (0%)	84	92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	169	LYS
1	C	139	LYS
1	C	142	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	120	ASN
1	A	151	HIS
1	A	188	GLN
1	C	146	GLN
1	D	2	ASN
1	D	165	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	D	301	-	4,4,4	0.23	0	6,6,6	0.06	0
2	SO4	C	301	-	4,4,4	0.21	0	6,6,6	0.12	0
3	GOL	A	302	-	5,5,5	0.28	0	5,5,5	0.74	0
2	SO4	B	301	-	4,4,4	0.22	0	6,6,6	0.07	0
2	SO4	A	301	-	4,4,4	0.22	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	302	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	GOL	O1-C1-C2-C3
3	A	302	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	SO4	1	0
3	A	302	GOL	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/236 (91%)	-0.19	4 (1%) 67 69	27, 47, 83, 102	0
1	B	216/236 (91%)	-0.07	2 (0%) 81 82	29, 54, 86, 107	0
1	C	215/236 (91%)	-0.03	3 (1%) 73 75	34, 54, 87, 99	0
1	D	214/236 (90%)	-0.06	0 100 100	37, 56, 89, 106	0
All	All	862/944 (91%)	-0.09	9 (1%) 79 80	27, 53, 87, 107	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	186	ASP	3.0
1	B	173	PRO	2.8
1	C	36	TRP	2.8
1	A	-1	SER	2.6
1	C	193	ARG	2.4
1	A	165	ASN	2.3
1	A	170	LEU	2.1
1	C	186	ASP	2.1
1	B	170	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	302	6/6	0.81	0.21	40,62,67,68	0
2	SO4	C	301	5/5	0.84	0.13	85,86,86,89	0
2	SO4	A	301	5/5	0.88	0.12	89,90,90,91	0
2	SO4	D	301	5/5	0.90	0.11	91,92,93,93	0
2	SO4	B	301	5/5	0.90	0.11	76,77,78,78	0
4	CL	B	302	1/1	0.99	0.07	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.