



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:26 PM UTC

PDB ID : 5IRG / pdb_00005irg
Title : Reverse transcriptase domain of group II intron maturase from *Roseburia intestinalis* in P212121 space group
Authors : Zhao, C.; Pyle, A.M.
Deposited on : 2016-03-13
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	0% 84% 10% 6%
1	B	305	2% 81% 13% 6%
1	C	305	4% 80% 12% 7%
1	D	305	4% 83% 11% 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retron-type reverse transcriptase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	Se	0	2	0
			2317	1463	407	432	2	13			
1	B	288	Total	C	N	O	S	Se	0	0	0
			2309	1457	405	433	2	12			
1	C	284	Total	C	N	O	S	Se	0	0	0
			2283	1440	400	429	2	12			
1	D	286	Total	C	N	O	S	Se	0	1	0
			2299	1451	402	431	2	13			

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	K	0	0
			1	1		
2	B	1	Total	K	0	0
			1	1		
2	C	1	Total	K	0	0
			1	1		
2	D	1	Total	K	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total	O	0	0
			102	102		
3	B	69	Total	O	0	1
			70	70		
3	C	56	Total	O	0	0
			56	56		
3	D	77	Total	O	0	0
			77	77		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retron-type reverse transcriptase




- Molecule 1: Retron-type reverse transcriptase




- Molecule 1: Retron-type reverse transcriptase



- Molecule 1: Retron-type reverse transcriptase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.81Å 145.35Å 200.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.13 – 2.30 49.13 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.13-2.30) 99.9 (49.13-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.29Å)	Xtriage
Refinement program	PHENIX (dev_2254: ???)	Depositor
R, R_{free}	0.191 , 0.231 0.197 , 0.237	Depositor DCC
R_{free} test set	2874 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.323	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9517	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.10	0/2349	0.30	0/3140
1	B	0.09	0/2335	0.29	0/3123
1	C	0.19	0/2307	0.32	0/3084
1	D	0.10	0/2326	0.28	0/3109
All	All	0.13	0/9317	0.30	0/12456

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	0	2347	18	0
1	B	2309	0	2330	23	0
1	C	2283	0	2302	29	0
1	D	2299	0	2323	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	102	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	70	0	0	1	0
3	C	56	0	0	0	0
3	D	77	0	0	2	0
All	All	9517	0	9302	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:270:ASN:OD1	1:C:273:LYS:N	1.88	1.07
1:C:270:ASN:CG	1:C:273:LYS:HB2	2.10	0.77
1:C:153:ASP:CG	1:C:273:LYS:HD2	2.16	0.70
1:D:82:ILE:HD12	1:D:92:LEU:HD12	1.72	0.70
1:A:140:ASP:HB3	1:D:37:ARG:HH12	1.57	0.70
1:A:144:ASP:OD2	1:D:37:ARG:NH2	2.29	0.65
1:D:79:ARG:HH21	1:D:91:ASN:HB3	1.62	0.64
1:D:35:VAL:HG21	1:D:101:PHE:HA	1.81	0.63
1:C:35:VAL:HG21	1:C:101:PHE:HA	1.81	0.63
1:A:35:VAL:HG21	1:A:101:PHE:HA	1.80	0.62
1:A:297:GLN:NE2	3:A:504:HOH:O	2.33	0.62
1:B:229:ARG:HH12	1:B:256:ASN:HB3	1.64	0.61
1:C:65:GLU:OE1	1:C:68:ARG:NH2	2.33	0.61
1:A:90[B]:ARG:NH2	3:A:505:HOH:O	2.34	0.59
1:A:92:LEU:HG	1:A:192[B]:MSE:SE	2.53	0.58
1:D:222:LEU:HG	1:D:226:MSE:HE2	1.85	0.57
1:C:271:MSE:HE3	1:C:271:MSE:HA	1.87	0.56
1:B:150:VAL:HB	1:B:242:ILE:HB	1.88	0.56
1:C:227:GLU:OE1	1:D:70:ARG:NH2	2.38	0.56
1:B:47:MSE:HE3	1:B:52:LEU:HA	1.87	0.55
1:B:35:VAL:HG21	1:B:101:PHE:HA	1.89	0.55
1:A:49:TYR:HA	1:A:97:VAL:HG21	1.89	0.55
1:A:222:LEU:HG	1:A:226:MSE:HE2	1.90	0.54
1:C:93:GLY:H	1:C:192:MSE:SE	2.41	0.54
1:C:153:ASP:OD2	1:C:273:LYS:HD2	2.09	0.53
1:C:222:LEU:HG	1:C:226:MSE:HE2	1.91	0.53
1:C:247:GLU:O	1:C:251:ASN:ND2	2.42	0.52
1:D:90:ARG:NH2	3:D:504:HOH:O	2.42	0.52
1:C:166:LEU:HD13	1:C:218:MSE:HB2	1.92	0.51
1:D:161:VAL:HG23	1:D:218:MSE:HE2	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:103:GLN:HB3	1:C:210:LEU:HB2	1.94	0.50
1:B:229:ARG:NH1	1:B:256:ASN:HB3	2.27	0.50
1:C:270:ASN:ND2	1:C:273:LYS:HB2	2.25	0.50
1:B:222:LEU:HG	1:B:226:MSE:HE2	1.94	0.49
1:D:229:ARG:HH22	1:D:256:ASN:HB3	1.78	0.49
1:C:252:ARG:HH22	1:D:165:LYS:HG2	1.77	0.48
1:D:162:ASN:HB3	1:D:165:LYS:HB2	1.95	0.48
1:C:154:LEU:HD21	1:C:261:ILE:HD13	1.96	0.47
1:D:166:LEU:HD13	1:D:218:MSE:HB2	1.96	0.46
1:D:150:VAL:HB	1:D:242:ILE:HB	1.97	0.46
1:B:38:ASN:ND2	3:B:506:HOH:O	2.44	0.46
1:A:19:MSE:HE2	1:A:23:LEU:HG	1.96	0.46
1:A:229:ARG:HH12	1:A:256:ASN:HB3	1.81	0.45
1:B:158:PHE:HE2	1:B:193:ILE:HG13	1.82	0.45
1:A:83:PRO:HA	1:A:89:VAL:HG12	1.97	0.45
1:B:140:ASP:HB3	1:C:37:ARG:HH21	1.81	0.45
1:A:19:MSE:HE3	1:A:22:ILE:HB	1.98	0.45
1:A:26:ASP:OD2	1:A:30:ARG:NH1	2.50	0.45
1:B:283:LYS:HG3	1:B:303:HIS:NE2	2.32	0.45
1:A:150:VAL:HB	1:A:242:ILE:HB	1.98	0.44
1:C:96:THR:HG22	1:C:99:ASP:H	1.81	0.44
1:B:49:TYR:HA	1:B:97:VAL:HG21	2.00	0.44
1:C:229:ARG:HH22	1:C:256:ASN:HB3	1.81	0.44
1:C:169:ILE:HD13	1:C:221:GLU:HG3	2.00	0.43
1:D:153:ASP:HB2	1:D:273:LYS:HD2	2.00	0.43
1:D:82:ILE:O	1:D:90:ARG:N	2.48	0.43
1:B:32:TYR:OH	1:B:53:LYS:HB2	2.18	0.43
1:C:161:VAL:HG23	1:C:218:MSE:HE3	2.00	0.43
1:B:45:ASP:OD1	1:B:96:THR:OG1	2.37	0.43
1:D:79:ARG:NH2	1:D:91:ASN:HB3	2.32	0.43
1:C:158:PHE:HA	1:C:161:VAL:HG12	2.00	0.43
1:B:157:PHE:CZ	1:B:215:ALA:HA	2.54	0.43
1:B:279:PRO:O	1:B:289:PHE:HB2	2.19	0.42
1:A:150:VAL:HG12	1:A:152:ILE:HG13	2.01	0.42
1:C:269:VAL:O	1:C:271:MSE:N	2.53	0.42
1:B:157:PHE:CZ	1:B:161:VAL:HG11	2.54	0.42
1:C:166:LEU:HD23	1:C:187:LEU:HD11	2.01	0.42
1:D:273:LYS:NZ	3:D:515:HOH:O	2.52	0.42
1:A:32:TYR:OH	1:A:53:LYS:HB2	2.20	0.42
1:B:19:MSE:HE3	1:B:22:ILE:HB	2.01	0.42
1:D:211:SER:OG	1:D:212:PRO:HD3	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:GLU:OE2	3:A:501:HOH:O	2.21	0.41
1:B:158:PHE:HA	1:B:161:VAL:HG22	2.02	0.41
1:C:165:LYS:HE3	1:C:169:ILE:HD11	2.02	0.41
1:A:211:SER:OG	1:A:212:PRO:HD3	2.20	0.41
1:B:141:MSE:SE	1:B:232:ASN:HB3	2.71	0.41
1:B:169:ILE:HD13	1:B:221:GLU:HG3	2.01	0.41
1:B:93:GLY:H	1:B:192:MSE:SE	2.53	0.41
1:B:162:ASN:OD1	1:B:165:LYS:N	2.51	0.41
1:D:64:LYS:O	1:D:68:ARG:HG3	2.21	0.41
1:C:249:SER:OG	1:D:164:ASP:OD1	2.34	0.41
1:C:277:ASP:OD1	1:C:278:ARG:N	2.52	0.41
1:D:93:GLY:H	1:D:192[B]:MSE:SE	2.54	0.41
1:B:211:SER:OG	1:B:212:PRO:HD3	2.19	0.40
1:C:270:ASN:CG	1:C:273:LYS:CB	2.89	0.40
1:C:251:ASN:HA	1:C:254:MSE:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/305 (94%)	284 (99%)	3 (1%)	0	100	100
1	B	286/305 (94%)	284 (99%)	2 (1%)	0	100	100
1	C	280/305 (92%)	275 (98%)	5 (2%)	0	100	100
1	D	283/305 (93%)	278 (98%)	5 (2%)	0	100	100
All	All	1136/1220 (93%)	1121 (99%)	15 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/255 (100%)	252 (99%)	3 (1%)	63	79
1	B	253/255 (99%)	249 (98%)	4 (2%)	55	73
1	C	251/255 (98%)	243 (97%)	8 (3%)	34	51
1	D	253/255 (99%)	247 (98%)	6 (2%)	43	62
All	All	1012/1020 (99%)	991 (98%)	21 (2%)	47	66

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	LEU
1	A	271	MSE
1	A	285	LEU
1	B	92	LEU
1	B	96	THR
1	B	166	LEU
1	B	283	LYS
1	C	15	THR
1	C	25	ASN
1	C	91	ASN
1	C	96	THR
1	C	210	LEU
1	C	269	VAL
1	C	271	MSE
1	C	273	LYS
1	D	18	LEU
1	D	154	LEU
1	D	165	LYS
1	D	227	GLU
1	D	268	LYS
1	D	280	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	29	ASN
1	B	34	GLN
1	B	38	ASN
1	B	209	ASN
1	B	251	ASN
1	C	34	GLN
1	C	59	ASN
1	C	104	GLN
1	C	108	GLN
1	C	146	ASN
1	C	209	ASN
1	C	297	GLN
1	D	34	GLN
1	D	59	ASN
1	D	146	ASN
1	D	220	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/305 (90%)	-0.08	3 (1%) 78 79	21, 41, 63, 93	1 (0%)
1	B	276/305 (90%)	0.14	5 (1%) 67 69	29, 47, 69, 85	0
1	C	272/305 (89%)	0.36	11 (4%) 42 44	35, 52, 86, 110	0
1	D	274/305 (89%)	0.06	11 (4%) 42 44	24, 44, 76, 104	0
All	All	1097/1220 (89%)	0.12	30 (2%) 56 58	21, 45, 76, 110	1 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	208	GLY	3.9
1	D	83	PRO	3.4
1	C	15	THR	3.0
1	D	16	SER	2.9
1	C	272	THR	2.8
1	C	304	ALA	2.7
1	C	152	ILE	2.7
1	B	17	SER	2.6
1	B	161	VAL	2.6
1	C	160	THR	2.5
1	D	15	THR	2.5
1	B	283	LYS	2.5
1	C	274	SER	2.5
1	C	91	ASN	2.4
1	D	90	ARG	2.4
1	C	90	ARG	2.3
1	C	16	SER	2.3
1	B	304	ALA	2.3
1	D	89	VAL	2.3
1	D	209	ASN	2.2
1	D	304	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	159	ASP	2.2
1	D	17	SER	2.2
1	A	17	SER	2.2
1	C	48	LYS	2.1
1	D	273	LYS	2.1
1	D	82	ILE	2.1
1	C	83	PRO	2.1
1	A	71	LYS	2.0
1	B	303	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	K	C	401	1/1	0.86	0.24	114,114,114,114	0
2	K	B	401	1/1	0.87	0.14	96,96,96,96	0
2	K	D	401	1/1	0.95	0.08	84,84,84,84	0
2	K	A	401	1/1	0.96	0.10	64,64,64,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.