



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:02 AM UTC

PDB ID : 1IS9 / pdb_00001is9
Title : Endoglucanase A from Clostridium thermocellum at atomic resolution
Authors : Schmidt, A.; Gonzalez, A.; Morris, R.J.; Costabel, M.; Alzari, P.M.; Lamzin, V.S.
Deposited on : 2001-11-26
Resolution : 1.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1063 (1.06-1.02)
Ramachandran outliers	187476	1034 (1.06-1.02)
Sidechain outliers	187428	1035 (1.06-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	363	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	1010	-	-	X	-
3	CL	A	1014	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called endoglucanase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	7	0
			2827	1790	476	550	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	GLN	LEU	conflict	UNP P04955

- Molecule 2 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	Hg	0	0
			6	6		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	Cl	0	0
			7	7		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	760	Total	O	0	0
			760	760		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: endoglucanase A



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.44Å 62.83Å 103.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.03	Depositor
% Data completeness (in resolution range)	98.0 (25.00-1.03)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.131 , 0.149	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3600	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2940	0.99	6/3994 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	162	HIS	N-CA-C	6.95	121.02	112.54
1	A	3	VAL	CA-C-O	-6.05	111.85	119.95
1	A	335	GLU	N-CA-C	5.93	118.69	111.82
1	A	171	ASP	CA-CB-CG	5.56	118.16	112.60
1	A	120	ASP	CA-CB-CG	5.56	118.16	112.60
1	A	319	ASN	CA-CB-CG	-5.22	107.38	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	3	VAL	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2827	0	2638	71	1
2	A	6	0	0	0	0
3	A	7	0	0	7	0
4	A	760	0	0	54	9
All	All	3600	0	2638	71	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:PHE:HD1	4:A:1722:HOH:O	1.02	1.33
1:A:110:SER:CB	4:A:1725:HOH:O	1.74	1.30
1:A:171:ASP:HA	4:A:1738:HOH:O	1.32	1.27
1:A:283:LYS:HG2	4:A:1717:HOH:O	1.28	1.21
1:A:131:PHE:CD1	4:A:1722:HOH:O	1.79	1.16
1:A:153:ASN:CG	4:A:1773:HOH:O	1.90	1.12
1:A:26[A]:MET:HE2	4:A:1209:HOH:O	1.53	1.05
1:A:153:ASN:ND2	4:A:1773:HOH:O	1.89	1.04
1:A:16:ILE:O	4:A:1471:HOH:O	1.77	1.02
1:A:42:ASN:OD1	4:A:1719:HOH:O	1.77	1.00
1:A:42:ASN:HB3	4:A:1429:HOH:O	1.61	0.99
1:A:5:PHE:CZ	4:A:1742:HOH:O	2.13	0.99
1:A:363:LEU:O	4:A:1765:HOH:O	1.80	0.97
1:A:323:GLU:HG3	4:A:1739:HOH:O	1.67	0.94
1:A:362:ASP:CG	4:A:1770:HOH:O	2.14	0.90
1:A:362:ASP:OD2	4:A:1770:HOH:O	1.89	0.89
1:A:131:PHE:CD2	3:A:1010:CL:CL	2.66	0.85
1:A:323:GLU:OE1	4:A:1690:HOH:O	1.95	0.84
1:A:171:ASP:CA	4:A:1738:HOH:O	2.03	0.81
1:A:153:ASN:CB	4:A:1773:HOH:O	2.26	0.80
1:A:23:VAL:N	4:A:1716:HOH:O	2.14	0.80
1:A:199:THR:HG23	4:A:1684:HOH:O	1.80	0.79
1:A:5:PHE:HZ	4:A:1742:HOH:O	1.56	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LEU:HD23	3:A:1010:CL:CL	2.22	0.76
1:A:110:SER:CA	4:A:1725:HOH:O	2.14	0.75
1:A:170:GLY:HA2	3:A:1014:CL:CL	2.24	0.75
1:A:300:HIS:HE1	1:A:327:GLU:OE2	1.69	0.74
1:A:4:PRO:O	4:A:1549:HOH:O	2.05	0.74
1:A:131:PHE:HB3	4:A:1722:HOH:O	1.87	0.73
1:A:110:SER:HA	4:A:1725:HOH:O	1.81	0.71
1:A:285:ILE:O	1:A:300:HIS:HD2	1.74	0.70
1:A:323:GLU:CD	4:A:1690:HOH:O	2.33	0.69
1:A:322:LYS:HG3	4:A:1473:HOH:O	1.93	0.68
1:A:28:LYS:NZ	4:A:1603:HOH:O	2.19	0.67
1:A:242:ASP:OD2	4:A:1402:HOH:O	2.13	0.66
1:A:111:HIS:CD2	4:A:1713:HOH:O	2.49	0.66
1:A:70:LEU:CD2	3:A:1010:CL:CL	2.81	0.66
1:A:250:TYR:CE2	4:A:1194:HOH:O	2.49	0.65
1:A:158:HIS:HE1	4:A:1082:HOH:O	1.79	0.65
1:A:335:GLU:HB2	4:A:1263:HOH:O	1.98	0.64
1:A:131:PHE:CB	4:A:1722:HOH:O	2.40	0.62
1:A:111:HIS:N	4:A:1723:HOH:O	1.77	0.60
1:A:283:LYS:CG	4:A:1717:HOH:O	2.08	0.60
1:A:153:ASN:HB3	4:A:1773:HOH:O	1.98	0.58
1:A:110:SER:HA	4:A:1723:HOH:O	2.03	0.57
1:A:171:ASP:N	4:A:1738:HOH:O	2.26	0.57
1:A:170:GLY:C	4:A:1738:HOH:O	2.48	0.57
1:A:42:ASN:CB	4:A:1429:HOH:O	2.34	0.56
1:A:26[A]:MET:CE	4:A:1209:HOH:O	2.29	0.56
1:A:117:ALA:C	3:A:1014:CL:CL	2.87	0.54
1:A:322:LYS:HA	4:A:1471:HOH:O	2.12	0.50
1:A:323:GLU:CG	4:A:1739:HOH:O	2.43	0.49
1:A:110:SER:CA	4:A:1723:HOH:O	2.56	0.49
1:A:362:ASP:HA	4:A:1523:HOH:O	2.13	0.48
1:A:111:HIS:HD2	4:A:1713:HOH:O	1.93	0.48
1:A:99:HIS:HD2	1:A:115:ASP:O	1.97	0.47
1:A:7:THR:O	4:A:1721:HOH:O	2.20	0.47
1:A:156:TYR:O	1:A:162:HIS:HE1	1.99	0.46
1:A:35:LYS:NZ	1:A:77:GLU:OE2	2.40	0.45
1:A:153:ASN:ND2	4:A:1325:HOH:O	2.49	0.45
1:A:301:ASN:HB3	4:A:1124:HOH:O	2.17	0.43
1:A:131:PHE:CE2	3:A:1010:CL:CL	3.09	0.43
1:A:23:VAL:CA	4:A:1716:HOH:O	2.64	0.43
1:A:31:TRP:CE2	1:A:35:LYS:HE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:300:HIS:CE1	1:A:327:GLU:OE2	2.60	0.43
1:A:28:LYS:CE	4:A:1603:HOH:O	2.66	0.43
1:A:26[A]:MET:HE3	1:A:329:VAL:HG13	2.03	0.41
1:A:31:TRP:NE1	1:A:35:LYS:HE2	2.35	0.41
1:A:165:TYR:O	3:A:1011:CL:CL	2.76	0.40
1:A:224:LEU:HD23	1:A:273[A]:LEU:CD1	2.51	0.40
1:A:272:MET:CE	4:A:1039:HOH:O	2.69	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:LYS:CD	4:A:1698:HOH:O[1_655]	1.77	0.43
4:A:1417:HOH:O	4:A:1569:HOH:O[4_445]	1.90	0.30
4:A:1643:HOH:O	4:A:1736:HOH:O[4_545]	1.95	0.25
4:A:1672:HOH:O	4:A:1754:HOH:O[3_645]	1.95	0.25
4:A:1094:HOH:O	4:A:1492:HOH:O[4_555]	2.08	0.12
4:A:1741:HOH:O	4:A:1754:HOH:O[3_645]	2.12	0.08
4:A:1404:HOH:O	4:A:1477:HOH:O[3_645]	2.13	0.07
4:A:1518:HOH:O	4:A:1579:HOH:O[4_445]	2.14	0.06
4:A:1652:HOH:O	4:A:1653:HOH:O[3_755]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	361/363 (99%)	352 (98%)	8 (2%)	1 (0%)	36 10

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/292 (100%)	292 (100%)	1 (0%)	86	65

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	HIS
1	A	135	GLN
1	A	153	ASN
1	A	154	ASN
1	A	157	ASN
1	A	158	HIS
1	A	162	HIS
1	A	264	GLN
1	A	299	ASN
1	A	300	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 13 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.