



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 02:54 AM UTC

PDB ID : 4IS3 / pdb_00004is3
Title : Crystal structure of a 3alpha-hydroxysteroid dehydrogenase (BaiA2) associated with secondary bile acid synthesis from *Clostridium scindens* VPI12708 in complex with a putative NAD(+)-OH- adduct at 2.0 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2013-01-16
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	90% 7%
1	B	268	90% 7%
1	C	268	90% 7%
1	D	268	90% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNL	A	301	-	-	X	-
3	UNL	B	301	-	-	X	-
3	UNL	C	301	-	-	X	-
3	UNL	D	301	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile acid 3-alpha hydroxysteroid dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	Se	0	5	0
			1882	1193	316	362	2	9			
1	B	249	Total	C	N	O	S	Se	0	11	0
			1924	1220	322	371	2	9			
1	C	250	Total	C	N	O	S	Se	0	8	0
			1911	1211	324	365	2	9			
1	D	249	Total	C	N	O	S	Se	0	6	0
			1890	1198	316	365	2	9			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	expression tag	UNP P19337
A	-17	GLY	-	expression tag	UNP P19337
A	-16	SER	-	expression tag	UNP P19337
A	-15	ASP	-	expression tag	UNP P19337
A	-14	LYS	-	expression tag	UNP P19337
A	-13	ILE	-	expression tag	UNP P19337
A	-12	HIS	-	expression tag	UNP P19337
A	-11	HIS	-	expression tag	UNP P19337
A	-10	HIS	-	expression tag	UNP P19337
A	-9	HIS	-	expression tag	UNP P19337
A	-8	HIS	-	expression tag	UNP P19337
A	-7	HIS	-	expression tag	UNP P19337
A	-6	GLU	-	expression tag	UNP P19337
A	-5	ASN	-	expression tag	UNP P19337
A	-4	LEU	-	expression tag	UNP P19337
A	-3	TYR	-	expression tag	UNP P19337
A	-2	PHE	-	expression tag	UNP P19337
A	-1	GLN	-	expression tag	UNP P19337
A	0	GLY	-	expression tag	UNP P19337
B	-18	MSE	-	expression tag	UNP P19337
B	-17	GLY	-	expression tag	UNP P19337

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP P19337
B	-15	ASP	-	expression tag	UNP P19337
B	-14	LYS	-	expression tag	UNP P19337
B	-13	ILE	-	expression tag	UNP P19337
B	-12	HIS	-	expression tag	UNP P19337
B	-11	HIS	-	expression tag	UNP P19337
B	-10	HIS	-	expression tag	UNP P19337
B	-9	HIS	-	expression tag	UNP P19337
B	-8	HIS	-	expression tag	UNP P19337
B	-7	HIS	-	expression tag	UNP P19337
B	-6	GLU	-	expression tag	UNP P19337
B	-5	ASN	-	expression tag	UNP P19337
B	-4	LEU	-	expression tag	UNP P19337
B	-3	TYR	-	expression tag	UNP P19337
B	-2	PHE	-	expression tag	UNP P19337
B	-1	GLN	-	expression tag	UNP P19337
B	0	GLY	-	expression tag	UNP P19337
C	-18	MSE	-	expression tag	UNP P19337
C	-17	GLY	-	expression tag	UNP P19337
C	-16	SER	-	expression tag	UNP P19337
C	-15	ASP	-	expression tag	UNP P19337
C	-14	LYS	-	expression tag	UNP P19337
C	-13	ILE	-	expression tag	UNP P19337
C	-12	HIS	-	expression tag	UNP P19337
C	-11	HIS	-	expression tag	UNP P19337
C	-10	HIS	-	expression tag	UNP P19337
C	-9	HIS	-	expression tag	UNP P19337
C	-8	HIS	-	expression tag	UNP P19337
C	-7	HIS	-	expression tag	UNP P19337
C	-6	GLU	-	expression tag	UNP P19337
C	-5	ASN	-	expression tag	UNP P19337
C	-4	LEU	-	expression tag	UNP P19337
C	-3	TYR	-	expression tag	UNP P19337
C	-2	PHE	-	expression tag	UNP P19337
C	-1	GLN	-	expression tag	UNP P19337
C	0	GLY	-	expression tag	UNP P19337
D	-18	MSE	-	expression tag	UNP P19337
D	-17	GLY	-	expression tag	UNP P19337
D	-16	SER	-	expression tag	UNP P19337
D	-15	ASP	-	expression tag	UNP P19337
D	-14	LYS	-	expression tag	UNP P19337
D	-13	ILE	-	expression tag	UNP P19337

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	HIS	-	expression tag	UNP P19337
D	-11	HIS	-	expression tag	UNP P19337
D	-10	HIS	-	expression tag	UNP P19337
D	-9	HIS	-	expression tag	UNP P19337
D	-8	HIS	-	expression tag	UNP P19337
D	-7	HIS	-	expression tag	UNP P19337
D	-6	GLU	-	expression tag	UNP P19337
D	-5	ASN	-	expression tag	UNP P19337
D	-4	LEU	-	expression tag	UNP P19337
D	-3	TYR	-	expression tag	UNP P19337
D	-2	PHE	-	expression tag	UNP P19337
D	-1	GLN	-	expression tag	UNP P19337
D	0	GLY	-	expression tag	UNP P19337

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	B	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	C	1	Total 44	C 21	N 7	O 14	P 2	0	0
2	D	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 3 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	B	1	Total O 1 1	0	0
3	C	1	Total O 1 1	0	0
3	D	1	Total O 1 1	0	0

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	186	Total O 186 186	0	0
5	B	197	Total O 197 197	0	0
5	C	219	Total O 219 219	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	149	Total 149	O 149	0	0


- Molecule 1: Bile acid 3-alpha hydroxysteroid dehydrogenase




- Chain B: 90% • 7%



- Chain C: 90% • 7%



- Chain D: 90% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.51Å 92.89Å 82.92Å 90.00° 90.78° 90.00°	Depositor
Resolution (Å)	29.01 – 2.00 29.01 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.4 (29.01-2.00) 97.3 (29.01-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.00Å)	Xtriage
Refinement program	BUSTER-TNT 2.10.0, BUSTER 2.10.0	Depositor
R, R_{free}	0.152 , 0.182 (Not available) , 0.173	Depositor DCC
R_{free} test set	3444 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.692	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.044 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8546	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, NAD, UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/1917 (0.1%)	1.18	2/2582 (0.1%)
1	B	0.91	1/1977 (0.1%)	1.15	1/2660 (0.0%)
1	C	0.87	0/1954	1.16	1/2628 (0.0%)
1	D	0.87	0/1928	1.17	1/2596 (0.0%)
All	All	0.88	2/7776 (0.0%)	1.16	5/10466 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	73	MSE	SE-CE	-6.70	1.75	1.95
1	A	73	MSE	SE-CE	-5.57	1.78	1.95

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	36	PHE	CA-CB-CG	8.11	121.91	113.80
1	B	36	PHE	CA-CB-CG	7.88	121.68	113.80
1	A	36	PHE	CA-CB-CG	7.69	121.49	113.80
1	D	36	PHE	CA-CB-CG	7.66	121.46	113.80
1	A	196	ASN	CA-CB-CG	5.45	118.05	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1882	0	1920	4	0
1	B	1924	0	1973	5	0
1	C	1911	0	1961	6	0
1	D	1890	0	1925	5	0
2	A	44	0	26	4	0
2	B	44	0	26	2	0
2	C	44	0	26	5	0
2	D	44	0	26	4	0
3	A	1	0	0	4	0
3	B	1	0	0	2	0
3	C	1	0	0	4	0
3	D	1	0	0	4	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	186	0	0	0	0
5	B	197	0	0	0	0
5	C	219	0	0	0	0
5	D	149	0	0	0	0
All	All	8546	0	7889	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:300:NAD:C6N	3:C:301:UNL:O	1.67	1.39
2:D:300:NAD:C6N	3:D:301:UNL:O	1.75	1.35
2:A:300:NAD:C6N	3:A:301:UNL:O	1.84	1.24
2:B:300:NAD:C6N	3:B:301:UNL:O	2.12	0.98
2:C:300:NAD:C5N	3:C:301:UNL:O	2.26	0.83

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/268 (94%)	243 (96%)	9 (4%)	0	100	100
1	B	258/268 (96%)	251 (97%)	7 (3%)	0	100	100
1	C	256/268 (96%)	246 (96%)	10 (4%)	0	100	100
1	D	253/268 (94%)	241 (95%)	11 (4%)	1 (0%)	30	27
All	All	1019/1072 (95%)	981 (96%)	37 (4%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	197	GLY

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are unknown - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	B	401	-	3,3,3	1.42	1 (33%)	3,3,3	0.78	0
2	NAD	D	300	-	46,48,48	1.11	1 (2%)	64,73,73	0.49	0
4	ACT	A	400	-	3,3,3	1.18	0	3,3,3	0.98	0
2	NAD	C	300	-	46,48,48	1.20	1 (2%)	64,73,73	0.57	1 (1%)
2	NAD	B	300	-	46,48,48	1.35	1 (2%)	64,73,73	0.60	1 (1%)
2	NAD	A	300	-	46,48,48	1.31	1 (2%)	64,73,73	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	D	300	-	-	5/30/62/62	0/5/5/5
2	NAD	B	300	-	-	4/30/62/62	0/5/5/5
2	NAD	A	300	-	-	4/30/62/62	0/5/5/5
2	NAD	C	300	-	-	5/30/62/62	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	NAD	C2N-N1N	8.43	1.44	1.35
2	A	300	NAD	C2N-N1N	8.02	1.43	1.35
2	C	300	NAD	C2N-N1N	7.45	1.43	1.35
2	D	300	NAD	C2N-N1N	6.77	1.42	1.35
4	B	401	ACT	CH3-C	2.16	1.57	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	300	NAD	C6N-N1N-C2N	-2.62	119.65	121.88
2	C	300	NAD	C6N-N1N-C2N	-2.54	119.72	121.88

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	300	NAD	C5D-O5D-PN-O3
2	A	300	NAD	C5D-O5D-PN-O1N
2	A	300	NAD	C5D-O5D-PN-O2N
2	B	300	NAD	C5D-O5D-PN-O3
2	B	300	NAD	C5D-O5D-PN-O1N


There are no ring outliers.


4 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	300	NAD	4	0
2	C	300	NAD	5	0
2	B	300	NAD	2	0
2	A	300	NAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/268 (89%)	-0.56	0 100 100	14, 26, 43, 62	5 (2%)
1	B	240/268 (89%)	-0.65	0 100 100	12, 24, 40, 57	11 (4%)
1	C	241/268 (89%)	-0.56	1 (0%) 88 88	14, 23, 49, 85	8 (3%)
1	D	240/268 (89%)	-0.49	1 (0%) 88 88	14, 28, 58, 77	6 (2%)
All	All	961/1072 (89%)	-0.56	2 (0%) 91 90	12, 25, 48, 85	30 (3%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	198	ASN	2.4
1	C	200	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACT	B	401	4/4	0.87	0.14	33,35,35,36	0
3	UNL	B	301	1/-	0.93	0.09	34,34,34,34	0
4	ACT	A	400	4/4	0.93	0.16	37,37,38,38	0
3	UNL	A	301	1/-	0.93	0.13	44,44,44,44	0
3	UNL	D	301	1/-	0.95	0.24	40,40,40,40	0
3	UNL	C	301	1/-	0.97	0.10	30,30,30,30	0
2	NAD	C	300	44/44	0.98	0.04	18,23,27,29	0
2	NAD	D	300	44/44	0.98	0.05	22,31,33,35	0
2	NAD	A	300	44/44	0.98	0.04	20,24,28,29	0
2	NAD	B	300	44/44	0.98	0.04	17,22,27,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAD D 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.