



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:49 PM UTC

PDB ID : 8ITT / pdb\_00008itt  
Title : Crystal structure of lysophosphatidylcholine in complex with human serum albumin and myristate  
Authors : Wang, Y.; Jiang, L.G.; Huang, M.D.  
Deposited on : 2023-03-22  
Resolution : 3.03 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3685 (3.08-3.00)
Clashscore	190562	4007 (3.08-3.00)
Ramachandran outliers	187476	3834 (3.08-3.00)
Sidechain outliers	187428	3836 (3.08-3.00)
RSRZ outliers	180081	3684 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	580	
1	B	580	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MYR	B	601	-	-	X	-
2	MYR	B	602	-	-	X	-
3	LPC	A	604	-	-	X	-
3	LPC	A	605	-	-	X	-
3	LPC	A	606	-	-	X	-
3	LPC	B	603	-	-	X	-

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	579	4443	2808	750	844	41	0	0	0
1	B	579	4442	2807	752	842	41	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	580	GLU	GLN	conflict	UNP P02768
B	580	GLU	GLN	conflict	UNP P02768

- Molecule 2 is MYRISTIC ACID (CCD ID: MYR) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	16	14	2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	14	2		
2	A	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			16	14	2		

- Molecule 3 is [1-MYRISTOYL-GLYCEROL-3-YL]PHOSPHONYLCHOLINE (CCD ID: LPC) (formula: C<sub>22</sub>H<sub>47</sub>NO<sub>7</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	22	1	7	1		
3	A	1	Total	C	N	O	P	0	0
			31	22	1	7	1		
3	A	1	Total	C	N	O	P	0	0
			31	22	1	7	1		
3	B	1	Total	C	N	O	P	0	0
			31	22	1	7	1		
3	B	1	Total	C	N	O	P	0	0
			31	22	1	7	1		
3	B	1	Total	C	N	O	P	0	0
			31	22	1	7	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Albumin



#### • Molecule 1: Albumin





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.26Å 38.47Å 182.46Å 90.00° 104.78° 90.00°	Depositor
Resolution (Å)	47.63 – 3.03 47.63 – 3.03	Depositor EDS
% Data completeness (in resolution range)	86.0 (47.63-3.03) 85.9 (47.63-3.03)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.84 (at 2.73Å)	Xtrriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.261 , 0.291 0.261 , 0.291	Depositor DCC
$R_{free}$ test set	2383 reflections (7.51%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.6	Xtrriage
Anisotropy	0.431	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 43.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9167	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.94 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2227e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LPC, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.00	13/4530 (0.3%)	1.46	78/6141 (1.3%)
1	B	0.97	11/4528 (0.2%)	1.45	72/6134 (1.2%)
All	All	0.99	24/9058 (0.3%)	1.46	150/12275 (1.2%)

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	GLN	CA-C	-8.53	1.43	1.53
1	B	311	GLU	CA-C	-8.06	1.42	1.52
1	B	491	LEU	CA-C	-7.85	1.43	1.52
1	A	93	LYS	CA-C	-7.26	1.43	1.52
1	B	30	TYR	CA-C	-6.82	1.43	1.52

The worst 5 of 150 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	571	GLU	N-CA-C	-20.17	87.77	112.38
1	B	567	CYS	N-CA-C	-19.95	89.59	111.14
1	A	94	GLN	N-CA-C	17.95	131.35	108.19
1	B	299	PRO	N-CA-C	16.22	133.67	114.20
1	A	129	ASP	N-CA-C	15.01	131.02	113.01

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4443	0	4212	542	1
1	B	4442	0	4216	528	0
2	A	48	0	81	6	0
2	B	48	0	81	29	0
3	A	93	0	138	90	0
3	B	93	0	138	80	0
All	All	9167	0	8866	1087	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 60.

The worst 5 of 1087 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:PHE:CE2	3:B:603:LPC:H0A1	1.43	1.54
1:A:485:ARG:HH11	3:A:605:LPC:C0B	1.33	1.40
1:B:149:PHE:CE2	3:B:603:LPC:H0C3	1.58	1.37
1:B:158:ALA:HB1	3:B:603:LPC:CK	1.53	1.36
1:B:149:PHE:CZ	3:B:603:LPC:H0C3	1.61	1.34

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:GLU:O	1:A:497:TYR:OH[1_565]	1.93	0.27

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	577/580 (100%)	432 (75%)	97 (17%)	48 (8%)	0	2
1	B	575/580 (99%)	443 (77%)	95 (16%)	37 (6%)	1	5
All	All	1152/1160 (99%)	875 (76%)	192 (17%)	85 (7%)	1	3

5 of 85 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	GLU
1	A	57	GLU
1	A	95	GLU
1	A	120	VAL
1	A	130	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/508 (91%)	385 (83%)	77 (17%)	2	10
1	B	461/508 (91%)	390 (85%)	71 (15%)	2	12
All	All	923/1016 (91%)	775 (84%)	148 (16%)	2	11

5 of 148 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	324	ASP
1	B	523	ILE
1	B	351	LYS
1	B	464	HIS
1	A	425	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	104	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	128	HIS
1	B	111	ASN
1	B	146	HIS
1	A	390	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	LPC	A	604	-	30,30,30	0.96	4 (13%)	35,37,37	1.24	3 (8%)
2	MYR	B	606	-	15,15,15	0.53	0	15,15,15	1.21	2 (13%)
3	LPC	A	605	-	30,30,30	1.00	1 (3%)	35,37,37	0.93	2 (5%)
2	MYR	B	601	-	15,15,15	0.67	1 (6%)	15,15,15	0.89	0
2	MYR	A	601	-	15,15,15	0.55	0	15,15,15	1.08	1 (6%)
2	MYR	A	603	-	15,15,15	0.55	0	15,15,15	0.97	1 (6%)
2	MYR	A	602	-	15,15,15	0.70	0	15,15,15	2.02	5 (33%)
3	LPC	A	606	-	30,30,30	1.02	1 (3%)	35,37,37	0.97	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	LPC	B	603	-	30,30,30	1.23	1 (3%)	35,37,37	0.95	2 (5%)
3	LPC	B	605	-	30,30,30	1.08	3 (10%)	35,37,37	2.53	13 (37%)
3	LPC	B	604	-	30,30,30	1.13	2 (6%)	35,37,37	0.97	2 (5%)
2	MYR	B	602	-	15,15,15	0.94	1 (6%)	15,15,15	1.61	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LPC	A	604	-	-	16/32/32/32	-
2	MYR	B	606	-	-	6/13/13/13	-
3	LPC	A	605	-	-	22/32/32/32	-
2	MYR	B	601	-	-	6/13/13/13	-
2	MYR	A	601	-	-	7/13/13/13	-
2	MYR	A	603	-	-	8/13/13/13	-
2	MYR	A	602	-	-	2/13/13/13	-
3	LPC	A	606	-	-	21/32/32/32	-
3	LPC	B	603	-	-	17/32/32/32	-
3	LPC	B	605	-	-	18/32/32/32	-
3	LPC	B	604	-	-	21/32/32/32	-
2	MYR	B	602	-	-	10/13/13/13	-

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	603	LPC	OQ2-CA	5.71	1.50	1.33
3	B	604	LPC	OQ2-CA	4.93	1.47	1.33
3	A	605	LPC	OQ2-CA	4.49	1.46	1.33
3	A	606	LPC	OQ2-CA	4.43	1.46	1.33
2	B	602	MYR	O1-C1	3.15	1.32	1.22

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	605	LPC	C0C-N1-C2	-5.26	89.00	109.91
3	B	605	LPC	C0B-N1-C2	-5.09	89.67	109.91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	605	LPC	C0A-N1-C2	-4.96	90.21	109.91
3	B	605	LPC	C0C-N1-C0B	4.84	121.69	108.98
3	B	605	LPC	CG-CF-CE	-4.81	90.08	114.37

There are no chirality outliers.

5 of 154 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	604	LPC	C3-O4-P5-O5B
3	A	604	LPC	C3-O4-P5-O6
3	A	604	LPC	C7-O6-P5-O4
3	A	605	LPC	N1-C2-C3-O4
3	A	605	LPC	C3-O4-P5-O5B


There are no ring outliers.




12 monomers are involved in 205 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	LPC	25	0
2	B	606	MYR	3	0
3	A	605	LPC	44	0
2	B	601	MYR	15	0
2	A	601	MYR	1	0
2	A	603	MYR	1	0
2	A	602	MYR	4	0
3	A	606	LPC	21	0
3	B	603	LPC	42	0
3	B	605	LPC	19	0
3	B	604	LPC	19	0
2	B	602	MYR	11	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	579/580 (99%)	0.22	9 (1%) 70 47	25, 50, 83, 154	0
1	B	579/580 (99%)	0.16	13 (2%) 62 38	21, 49, 83, 124	0
All	All	1158/1160 (99%)	0.19	22 (1%) 66 42	21, 50, 83, 154	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	CYS	4.4
1	B	95	GLU	4.3
1	B	170	GLN	3.8
1	B	568	PHE	3.7
1	B	561	ALA	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LPC	A	606	31/31	0.51	0.22	49,58,66,70	0
3	LPC	B	603	31/31	0.59	0.22	44,53,65,66	0
3	LPC	B	605	31/31	0.63	0.20	48,56,63,65	0
3	LPC	A	604	31/31	0.66	0.20	43,49,57,59	0
3	LPC	A	605	31/31	0.67	0.19	43,48,55,63	0
3	LPC	B	604	31/31	0.68	0.20	42,49,58,60	0
2	MYR	B	601	16/16	0.77	0.22	54,63,72,78	0
2	MYR	A	601	16/16	0.78	0.18	40,48,67,68	0
2	MYR	B	602	16/16	0.78	0.21	40,48,60,61	0
2	MYR	A	603	16/16	0.83	0.15	39,42,61,63	0
2	MYR	B	606	16/16	0.83	0.17	38,43,50,52	0
2	MYR	A	602	16/16	0.84	0.18	47,50,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around LPC B 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around LPC B 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around LPC A 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LPC A 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LPC B 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MYR B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MYR A 601:**

$2mF_o-DF_c$  (at 0.7 rnsd) in gray  
 $mF_o-DF_c$  (at 3 rnsd) in purple (negative)  
and green (positive)



**Electron density around MYR B 602:**

$2mF_o-DF_c$  (at 0.7 rnsd) in gray  
 $mF_o-DF_c$  (at 3 rnsd) in purple (negative)  
and green (positive)






**Electron density around MYR A 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MYR B 606:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around MYR A 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.