



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:29 PM UTC

PDB ID : 3IU7 / pdb_00003iu7
Title : M. tuberculosis methionine aminopeptidase with Mn inhibitor A02
Authors : Ye, Q.Z.; Lu, J.P.
Deposited on : 2009-08-30
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	83% 15% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FCD	A	289	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	0	0	0
			2166	1367	373	417	9			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP P0A5J2
A	-1	ALA	-	expression tag	UNP P0A5J2
A	0	SER	-	expression tag	UNP P0A5J2
A	1	MET	-	expression tag	UNP P0A5J2

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is 5-(2-CHLOROPHENYL)FURAN-2-CARBOXYLIC ACID (CCD ID: FCD) (formula: C₁₁H₇ClO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	O	0	0
			15	11	1	3		
3	A	1	Total	C	Cl	O	0	0
			15	11	1	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Cl 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	249	Total 249	O 249	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionine aminopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	106.41 Å 106.41 Å 50.42 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.14 – 1.40 21.14 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (21.14-1.40) 99.5 (21.14-1.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.90 (at 1.40 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.171 , 0.190 0.168 , 0.187	Depositor DCC
R_{free} test set	3237 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2453	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FCD, SO4, MN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.82	36/2217 (1.6%)	1.50	19/3030 (0.6%)

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	4	ARG	CZ-NH1	8.43	1.44	1.32
1	A	48	GLU	CD-OE2	7.53	1.39	1.25
1	A	28	VAL	CA-C	7.50	1.60	1.52
1	A	262	ARG	N-CA	7.42	1.54	1.46
1	A	267	GLN	C-O	6.90	1.32	1.23
1	A	12	LEU	C-O	6.66	1.32	1.23
1	A	30	LYS	C-O	-6.64	1.17	1.24
1	A	132	VAL	CA-CB	-6.54	1.45	1.54
1	A	149	ALA	N-CA	6.26	1.54	1.46
1	A	250	ILE	C-O	6.07	1.30	1.24
1	A	284	CYS	C-O	-6.05	1.16	1.23
1	A	222	GLN	C-N	5.96	1.39	1.33
1	A	263	LYS	CA-C	5.89	1.60	1.52
1	A	48	GLU	CD-OE1	5.83	1.36	1.25
1	A	262	ARG	CG-CD	-5.82	1.34	1.52
1	A	10	GLY	C-O	5.70	1.29	1.24
1	A	244	GLY	C-O	5.64	1.31	1.23
1	A	164	ARG	N-CA	-5.62	1.39	1.46
1	A	11	VAL	CA-CB	5.60	1.61	1.54
1	A	185	ARG	CG-CD	5.46	1.68	1.52
1	A	92	PRO	C-O	5.44	1.28	1.24
1	A	198	VAL	CA-CB	-5.37	1.47	1.54
1	A	116	ILE	N-CA	-5.28	1.41	1.46
1	A	159	LEU	CA-CB	-5.27	1.45	1.53
1	A	80	ILE	N-CA	5.25	1.52	1.46
1	A	132	VAL	CB-CG1	-5.23	1.35	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	274	VAL	N-CA	5.22	1.52	1.46
1	A	190	TYR	C-O	5.22	1.30	1.24
1	A	33	ALA	N-CA	5.20	1.52	1.45
1	A	185	ARG	CZ-NH2	5.16	1.40	1.33
1	A	168	MET	SD-CE	5.15	1.92	1.79
1	A	178	ARG	CA-CB	-5.15	1.45	1.53
1	A	16	ARG	CZ-NH2	5.10	1.40	1.33
1	A	48	GLU	CG-CD	5.08	1.64	1.52
1	A	123	THR	C-O	5.06	1.30	1.23
1	A	109	ASN	C-O	5.05	1.29	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	16	ARG	NE-CZ-NH2	-9.68	110.49	119.20
1	A	51	ARG	NE-CZ-NH2	-8.29	111.74	119.20
1	A	4	ARG	NE-CZ-NH2	-7.41	112.53	119.20
1	A	162	ARG	NE-CZ-NH2	-7.04	112.86	119.20
1	A	55	ARG	NE-CZ-NH2	6.67	125.21	119.20
1	A	4	ARG	CG-CD-NE	6.59	126.51	112.00
1	A	185	ARG	NH1-CZ-NH2	6.00	127.10	119.30
1	A	4	ARG	CA-C-N	5.80	131.96	121.06
1	A	4	ARG	C-N-CA	5.80	131.96	121.06
1	A	16	ARG	NH1-CZ-NH2	5.67	126.67	119.30
1	A	4	ARG	NE-CZ-NH1	5.52	127.02	121.50
1	A	55	ARG	NE-CZ-NH1	-5.47	116.03	121.50
1	A	144	ASN	CA-CB-CG	5.42	118.02	112.60
1	A	156	HIS	CA-C-N	5.38	127.44	120.44
1	A	156	HIS	C-N-CA	5.38	127.44	120.44
1	A	200	ARG	NE-CZ-NH1	5.24	126.74	121.50
1	A	112	ILE	O-C-N	5.16	127.27	121.90
1	A	28	VAL	N-CA-C	-5.05	102.98	109.30
1	A	152	VAL	O-C-N	5.00	128.63	122.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2166	0	2125	5	1
2	A	2	0	0	0	0
3	A	30	0	12	1	0
4	A	5	0	0	0	0
5	A	1	0	0	0	0
6	A	249	0	0	0	1
All	All	2453	0	2137	5	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:GLU:CG	1:A:260:LYS:HG2	2.27	0.65
1:A:249:GLU:HG3	1:A:260:LYS:HG2	1.81	0.63
1:A:179:ALA:O	1:A:182:VAL:HG22	2.14	0.48
1:A:168:MET:SD	3:A:289:FCD:HG	2.55	0.47
1:A:238:GLU:HA	1:A:268:PHE:O	2.14	0.47

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:ARG:NH2	6:A:436:HOH:O[6_554]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/288 (98%)	276 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/235 (99%)	231 (100%)	1 (0%)	84	67

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FCD	A	288	2	16,16,16	1.84	7 (43%)	21,22,22	1.66	4 (19%)
4	SO4	A	290	-	4,4,4	0.74	0	6,6,6	1.06	1 (16%)
3	FCD	A	289	-	16,16,16	2.72	7 (43%)	21,22,22	3.19	13 (61%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FCD	A	288	2	-	0/8/8/8	0/2/2/2
3	FCD	A	289	-	-	4/8/8/8	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	289	FCD	OA-CD	-7.00	1.25	1.38
3	A	289	FCD	OA-CA	-5.21	1.30	1.38
3	A	289	FCD	CG-CD	3.05	1.41	1.35
3	A	288	FCD	CB-CA	2.99	1.42	1.35
3	A	288	FCD	C2-C1	2.96	1.44	1.39
3	A	288	FCD	C5-C6	2.81	1.44	1.38
3	A	288	FCD	OA-CA	-2.79	1.33	1.38
3	A	289	FCD	C1-CD	2.56	1.53	1.47
3	A	288	FCD	CG-CD	2.50	1.40	1.35
3	A	289	FCD	CB-CA	2.39	1.41	1.35
3	A	289	FCD	C1-C6	2.37	1.42	1.39
3	A	289	FCD	C4-C5	-2.25	1.35	1.38
3	A	288	FCD	OA-CD	-2.21	1.34	1.38
3	A	288	FCD	CA-C	-2.04	1.44	1.48

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	289	FCD	CG-CB-CA	-7.95	99.84	106.94
3	A	289	FCD	C2-C1-C6	-5.80	111.01	117.79
3	A	289	FCD	C3-C2-C1	4.49	127.92	119.80
3	A	289	FCD	OA-CD-C1	4.19	121.95	115.24
3	A	289	FCD	CD-OA-CA	3.77	110.51	106.29
3	A	288	FCD	CG-CB-CA	-3.42	103.89	106.94
3	A	288	FCD	OB-C-OXT	-3.35	115.94	123.90
3	A	289	FCD	CB-CG-CD	3.34	109.19	107.04
3	A	289	FCD	OB-C-CA	3.15	121.19	114.06
3	A	289	FCD	C4-C3-C2	-2.82	116.76	120.24
3	A	289	FCD	OA-CA-CB	2.61	112.70	109.68
3	A	289	FCD	OB-C-OXT	-2.29	118.45	123.90
3	A	288	FCD	OXT-C-CA	2.26	126.11	120.57
3	A	288	FCD	C4-C3-C2	2.23	122.99	120.24
3	A	289	FCD	C1-CD-CG	-2.17	130.26	135.06
3	A	289	FCD	OA-CD-CG	-2.12	107.76	109.39
3	A	289	FCD	C6-C1-CD	2.09	126.97	124.05
4	A	290	SO4	O3-S-O1	2.02	120.11	109.56

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	289	FCD	OB-C-CA-OA
3	A	289	FCD	OXT-C-CA-OA
3	A	289	FCD	OB-C-CA-CB
3	A	289	FCD	OXT-C-CA-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	289	FCD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/288 (98%)	-0.39	1 (0%) 88 91	5, 9, 20, 30	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	PRO	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FCD	A	289	15/15	0.88	0.13	16,31,49,51	0
5	CL	A	291	1/1	0.94	0.22	41,41,41,41	0
4	SO4	A	290	5/5	0.97	0.12	19,19,25,25	0
3	FCD	A	288	15/15	0.98	0.05	7,9,12,12	0
2	MN	A	286	1/1	1.00	0.02	6,6,6,6	0
2	MN	A	287	1/1	1.00	0.01	6,6,6,6	0

6.5 Other polymers [i](#)

There are no such residues in this entry.