



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:31 PM UTC

PDB ID : 9IV7 / pdb_00009iv7
Title : Crystal structure of CcmS-CcmK1-CcmK2 complex from *Synechocystis* sp. PCC 6803
Authors : Li, J.; Deng, J.X.; Jiang, Y.L.; Zhou, C.Z.
Deposited on : 2024-07-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	3% 80% 10% 10%
1	G	142	3% 76% 14% 10%
2	B	111	79% 20% .
2	C	111	86% 12% ..
2	F	111	3% 76% 7% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	111	
3	D	103	
3	E	103	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TAR	C	201	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Slr1911 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	128	Total	C	N	O	S	0	0	0
			1038	678	173	184	3			
1	G	128	Total	C	N	O	S	0	0	0
			1038	678	173	184	3			


- Molecule 2 is a protein called Carboxysome shell protein CcmK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	110	Total	C	N	O	S	0	0	0
			841	528	153	158	2			
2	C	110	Total	C	N	O	S	0	0	0
			841	528	153	158	2			
2	F	92	Total	C	N	O	S	0	0	0
			690	434	124	130	2			
2	H	92	Total	C	N	O	S	0	0	0
			690	434	124	130	2			

- Molecule 3 is a protein called Carboxysome shell protein CcmK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	102	Total	C	N	O	S	0	0	0
			773	484	137	150	2			
3	E	102	Total	C	N	O	S	0	0	0
			773	484	137	150	2			

- Molecule 4 is D(-)-TARTARIC ACID (CCD ID: TAR) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			10	4	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	36	Total	O	0	0
			36	36		
5	B	58	Total	O	0	0
			58	58		
5	C	57	Total	O	0	0
			57	57		
5	D	51	Total	O	0	0
			51	51		
5	E	44	Total	O	0	0
			44	44		
5	F	36	Total	O	0	0
			36	36		
5	G	66	Total	O	0	0
			66	66		
5	H	31	Total	O	0	0
			31	31		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Slr1911 protein




- Molecule 1: Slr1911 protein




- Molecule 2: Carboxysome shell protein CcmK1




- Molecule 2: Carboxysome shell protein CcmK1




- Molecule 2: Carboxysome shell protein CcmK1




- Molecule 2: Carboxysome shell protein CcmK1



• Molecule 3: Carboxysome shell protein CcmK2



• Molecule 3: Carboxysome shell protein CcmK2



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	133.93Å 58.04Å 188.77Å 90.00° 105.48° 90.00°	Depositor
Resolution (Å)	13.24 – 2.50 13.24 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (13.24-2.50) 98.6 (13.24-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 2.50Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.215 , 0.241 0.214 , 0.240	Depositor DCC
R_{free} test set	2415 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	1.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7073	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1068	0.27	0/1452
1	G	0.14	0/1068	0.30	0/1452
2	B	0.11	0/851	0.27	0/1154
2	C	0.11	0/851	0.30	0/1154
2	F	0.11	0/698	0.26	0/947
2	H	0.11	0/698	0.28	0/947
3	D	0.11	0/783	0.26	0/1062
3	E	0.10	0/783	0.26	0/1062
All	All	0.12	0/6800	0.28	0/9230

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1038	0	1026	11	0
1	G	1038	0	1026	18	0
2	B	841	0	867	17	0
2	C	841	0	867	13	0
2	F	690	0	712	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	690	0	712	11	0
3	D	773	0	779	9	0
3	E	773	0	779	12	0
4	C	10	0	4	1	0
5	A	36	0	0	2	0
5	B	58	0	0	4	0
5	C	57	0	0	0	0
5	D	51	0	0	4	0
5	E	44	0	0	2	0
5	F	36	0	0	2	0
5	G	66	0	0	5	0
5	H	31	0	0	2	0
All	All	7073	0	6772	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:71:GLU:OE1	5:E:201:HOH:O	1.92	0.86
1:G:125:THR:OG1	5:G:201:HOH:O	1.98	0.81
3:E:24:VAL:O	5:E:202:HOH:O	2.00	0.79
2:H:19:ALA:HB2	2:H:64:ILE:HD11	1.71	0.70
2:C:101:ARG:NH1	1:G:52:ASP:OD1	2.25	0.68
2:B:99:GLN:N	5:B:202:HOH:O	2.27	0.68
2:C:86:GLU:HG2	1:G:105:LYS:HG3	1.77	0.66
1:G:17:ARG:NH1	5:G:205:HOH:O	2.29	0.64
2:B:19:ALA:HB2	2:B:64:ILE:HD11	1.80	0.64
1:A:79:GLN:NE2	5:A:203:HOH:O	2.30	0.63
2:C:111:ARG:NH1	1:G:127:GLU:OE2	2.31	0.63
1:A:81:MET:HE2	1:A:106:LEU:HG	1.84	0.59
2:F:62:GLU:OE1	2:F:65:ARG:NH2	2.36	0.59
1:A:62:CYS:HB3	2:B:104:VAL:HG12	1.85	0.58
2:H:59:ALA:O	2:H:63:ASN:ND2	2.36	0.58
3:D:86:GLU:OE1	5:D:201:HOH:O	2.17	0.58
3:E:96:GLU:N	3:E:96:GLU:OE1	2.33	0.58
1:A:60:ILE:HG22	2:B:106:PRO:HA	1.87	0.57
3:E:68:ASN:ND2	2:H:75:ASN:OD1	2.37	0.57
2:C:28:ARG:HA	1:G:79:GLN:HG2	1.87	0.57
1:A:54:HIS:HB2	2:B:101:ARG:HH22	1.69	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:59:ALA:O	2:F:63:ASN:ND2	2.34	0.56
2:C:19:ALA:HB2	2:C:64:ILE:HD11	1.87	0.56
2:H:92:ARG:HG3	2:H:93:TYR:H	1.70	0.55
3:E:61:ILE:HD11	3:E:75:THR:HG21	1.88	0.54
2:F:75:ASN:ND2	5:F:204:HOH:O	2.34	0.53
3:D:85:LEU:HG	3:D:89:LEU:HD12	1.90	0.53
2:F:60:GLY:O	2:F:64:ILE:HG12	2.09	0.52
1:G:52:ASP:HB2	1:G:60:ILE:HD13	1.92	0.51
2:F:72:VAL:O	5:F:201:HOH:O	2.19	0.51
2:C:76:HIS:NE2	2:H:18:GLU:OE1	2.36	0.50
1:G:65:GLN:NE2	1:G:69:GLN:OE1	2.35	0.50
1:A:81:MET:HE1	1:A:104:PHE:HB3	1.92	0.50
2:C:84:ASN:HA	1:G:106:LEU:O	2.11	0.50
2:B:20:ALA:HB2	2:B:44:VAL:HG11	1.94	0.50
3:D:2:SER:N	5:D:205:HOH:O	2.45	0.50
2:B:95:GLU:O	5:B:202:HOH:O	2.20	0.50
1:G:22:GLN:HE21	1:G:140:LYS:HD2	1.77	0.49
2:B:36:LYS:HG2	2:B:42:VAL:HG12	1.96	0.48
3:E:16:VAL:HG22	3:E:44:VAL:HG13	1.95	0.48
4:C:201:TAR:O1	3:E:39:SER:N	2.38	0.47
3:E:20:ALA:HB2	3:E:44:VAL:HG11	1.96	0.47
3:E:20:ALA:HB1	3:E:31:LEU:HD22	1.95	0.47
2:H:66:ARG:NH2	5:H:202:HOH:O	2.30	0.47
2:C:20:ALA:HB2	2:C:44:VAL:HG11	1.97	0.47
1:G:33:ALA:O	5:G:202:HOH:O	2.20	0.47
2:B:108:ILE:N	5:B:203:HOH:O	2.27	0.46
2:B:98:GLU:HA	2:B:101:ARG:HD3	1.97	0.46
2:H:27:ALA:HB1	2:H:52:GLU:HB3	1.98	0.46
2:H:62:GLU:O	2:H:65:ARG:HG2	2.15	0.46
2:B:95:GLU:O	2:B:99:GLN:HG2	2.16	0.46
2:B:109:ILE:HG12	5:B:203:HOH:O	2.16	0.46
3:E:17:VAL:HG22	2:H:89:LEU:HD13	1.97	0.45
2:C:103:ILE:HD12	2:C:103:ILE:HA	1.82	0.45
1:A:44:ALA:HB3	1:A:46:GLU:OE2	2.17	0.45
3:D:11:ARG:NH1	5:D:206:HOH:O	2.44	0.45
1:A:82:VAL:HG23	2:B:28:ARG:HD3	1.97	0.45
3:D:2:SER:OG	3:D:3:ILE:N	2.48	0.45
2:C:16:VAL:HG22	2:C:44:VAL:HG13	1.99	0.45
3:E:3:ILE:HD12	3:E:93:TYR:HE2	1.81	0.45
1:G:31:LEU:HD23	1:G:133:LEU:HD13	1.99	0.44
1:A:79:GLN:HG2	2:B:28:ARG:HA	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:GLU:OE1	2:B:2:SER:N	2.51	0.43
3:D:66:ARG:NE	5:D:207:HOH:O	2.48	0.43
2:B:100:PHE:O	2:B:103:ILE:HG22	2.18	0.43
2:C:86:GLU:OE2	1:G:105:LYS:HE2	2.19	0.43
2:C:109:ILE:HD13	1:G:127:GLU:HG3	2.01	0.43
2:C:13:PHE:O	2:C:16:VAL:HG12	2.18	0.43
3:E:40:GLY:HA2	2:H:37:ILE:HG13	2.01	0.42
3:D:84:ASN:HD22	3:D:84:ASN:C	2.28	0.42
1:G:28:GLN:H	1:G:28:GLN:HG2	1.70	0.42
1:G:37:ALA:N	5:G:202:HOH:O	2.25	0.42
2:H:21:ASP:OD2	5:H:201:HOH:O	2.21	0.41
3:D:13:PHE:O	3:D:16:VAL:HG12	2.21	0.41
1:G:126:ILE:HG22	5:G:216:HOH:O	2.19	0.41
1:A:137:LEU:N	5:A:202:HOH:O	2.54	0.41
3:D:20:ALA:HB2	3:D:44:VAL:HG11	2.03	0.40
2:B:7:MET:HG2	2:B:45:ILE:HG13	2.02	0.40
1:G:36:TRP:HB2	1:G:115:PRO:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/142 (89%)	121 (96%)	4 (3%)	1 (1%)	16	31
1	G	126/142 (89%)	121 (96%)	5 (4%)	0	100	100
2	B	108/111 (97%)	107 (99%)	1 (1%)	0	100	100
2	C	108/111 (97%)	106 (98%)	2 (2%)	0	100	100
2	F	90/111 (81%)	89 (99%)	1 (1%)	0	100	100
2	H	90/111 (81%)	88 (98%)	2 (2%)	0	100	100
3	D	100/103 (97%)	97 (97%)	3 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	100/103 (97%)	98 (98%)	2 (2%)	0	100	100
All	All	848/934 (91%)	827 (98%)	20 (2%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/120 (89%)	106 (99%)	1 (1%)	70	87
1	G	107/120 (89%)	107 (100%)	0	100	100
2	B	91/92 (99%)	89 (98%)	2 (2%)	45	73
2	C	91/92 (99%)	88 (97%)	3 (3%)	33	61
2	F	74/92 (80%)	74 (100%)	0	100	100
2	H	74/92 (80%)	74 (100%)	0	100	100
3	D	82/83 (99%)	81 (99%)	1 (1%)	63	83
3	E	82/83 (99%)	80 (98%)	2 (2%)	43	70
All	All	708/774 (92%)	699 (99%)	9 (1%)	61	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	GLN
2	B	5	VAL
2	B	16	VAL
2	C	5	VAL
2	C	16	VAL
2	C	103	ILE
3	D	16	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	3	ILE
3	E	16	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	99	GLN
2	C	54	GLN
3	E	54	GLN
2	F	54	GLN
1	G	22	GLN
1	G	47	GLN
2	H	54	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	TAR	C	201	-	9,9,9	1.37	2 (22%)	12,12,12	1.39	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TAR	C	201	-	-	12/12/12/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	201	TAR	O1-C1	2.93	1.30	1.22
4	C	201	TAR	O11-C1	-2.85	1.21	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	201	TAR	O11-C1-C2	3.40	122.77	113.31
4	C	201	TAR	O1-C1-C2	-3.37	112.65	121.62

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	201	TAR	C1-C2-C3-O3
4	C	201	TAR	O2-C2-C3-O3
4	C	201	TAR	O2-C2-C3-C4
4	C	201	TAR	C1-C2-C3-C4
4	C	201	TAR	O1-C1-C2-C3
4	C	201	TAR	O11-C1-C2-C3
4	C	201	TAR	O1-C1-C2-O2
4	C	201	TAR	O11-C1-C2-O2
4	C	201	TAR	O3-C3-C4-O4
4	C	201	TAR	O3-C3-C4-O41
4	C	201	TAR	C2-C3-C4-O4
4	C	201	TAR	C2-C3-C4-O41

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	201	TAR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/142 (90%)	0.32	4 (3%) 51 47	30, 47, 71, 88	0
1	G	128/142 (90%)	0.07	4 (3%) 51 47	28, 37, 66, 77	0
2	B	110/111 (99%)	-0.04	0 100 100	26, 37, 68, 82	0
2	C	110/111 (99%)	-0.23	0 100 100	26, 36, 54, 69	0
2	F	92/111 (82%)	0.06	3 (3%) 49 45	27, 39, 63, 77	0
2	H	92/111 (82%)	0.13	5 (5%) 31 27	30, 42, 64, 83	0
3	D	102/103 (99%)	-0.01	5 (4%) 35 31	26, 34, 70, 109	0
3	E	102/103 (99%)	-0.07	2 (1%) 65 61	28, 37, 53, 68	0
All	All	864/934 (92%)	0.03	23 (2%) 56 51	26, 38, 66, 109	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	102	THR	5.9
1	A	15	PRO	5.6
1	G	28	GLN	4.9
3	D	103	TYR	4.7
2	F	93	TYR	3.4
3	D	2	SER	3.0
2	F	2	SER	3.0
2	F	66	ARG	2.9
3	E	2	SER	2.9
3	D	87	TYR	2.7
2	H	2	SER	2.7
1	A	69	GLN	2.6
2	H	28	ARG	2.6
3	E	87	TYR	2.6
1	G	15	PRO	2.5
2	H	93	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	62	GLU	2.5
1	G	139	GLU	2.4
2	H	70	GLY	2.2
1	A	142	ALA	2.2
1	A	54	HIS	2.1
3	D	96	GLU	2.1
1	G	101	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	TAR	C	201	10/10	0.63	0.24	20,20,20,20	10

6.5 Other polymers [i](#)

There are no such residues in this entry.