



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:45 AM UTC

PDB ID : 2IW5 / pdb_00002iw5
Title : Structural Basis for CoREST-Dependent Demethylation of Nucleosomes by the Human LSD1 Histone Demethylase
Authors : Yang, M.; Gocke, C.B.; Luo, X.; Borek, D.; Tomchick, D.R.; Machius, M.; Otwinowski, Z.; Yu, H.
Deposited on : 2006-06-26
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	12% 91% 9%
2	B	235	16% 52% 5% 43%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LYSINE-SPECIFIC HISTONE DEMETHYLASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	0	0
			5217	3324	906	967	20			

- Molecule 2 is a protein called REST COREPRESSOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	0
			1076	676	194	203	3			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Cl	0	0
			3	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is AMMONIUM ION (CCD ID: NH4) (formula: H_4N).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total N 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	77	Total O 77 77	0	0
7	B	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LYSINE-SPECIFIC HISTONE DEMETHYLASE 1



• Molecule 2: REST COREPRESSOR 1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	120.35Å 178.22Å 234.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.57 20.00 – 2.57	Depositor EDS
% Data completeness (in resolution range)	93.2 (20.00-2.57) 93.1 (20.00-2.57)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.212 0.194 , 0.203	Depositor DCC
R_{free} test set	1499 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	62.7	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 69.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6440	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, CL, NH₄, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/5331	0.96	12/7232 (0.2%)
2	B	0.50	0/1091	0.95	0/1471
All	All	0.57	0/6422	0.96	12/8703 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	740	VAL	N-CA-C	11.52	118.05	107.56
1	A	287	GLY	N-CA-C	-7.53	101.11	112.41
1	A	697	LEU	N-CA-C	7.02	125.32	114.64
1	A	468	VAL	N-CA-C	-6.75	97.78	107.37
1	A	520	TYR	N-CA-C	-5.85	104.27	111.40
1	A	334	VAL	N-CA-C	-5.75	98.85	107.37
1	A	793	ILE	N-CA-CB	-5.75	103.16	111.21
1	A	304	VAL	N-CA-C	5.73	116.38	108.12
1	A	650	ALA	N-CA-C	-5.70	104.99	111.14
1	A	317	VAL	N-CA-C	-5.69	98.77	106.85
1	A	636	ALA	N-CA-C	-5.28	105.98	112.90
1	A	333	VAL	N-CA-C	5.09	115.92	108.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5217	0	5252	30	0
2	B	1076	0	1091	6	0
3	A	53	0	31	2	0
4	A	3	0	0	0	0
5	A	12	0	16	0	0
6	A	1	0	0	0	0
7	A	77	0	0	0	0
7	B	1	0	0	0	0
All	All	6440	0	6390	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ARG:HB3	2:B:314:MET:HE1	1.47	0.96
1:A:801:GLU:HG2	1:A:809:ALA:H	1.48	0.76
1:A:707:VAL:HG11	1:A:715:MET:HG3	1.75	0.67
1:A:801:GLU:HG3	1:A:809:ALA:HA	1.76	0.66
1:A:384:ARG:HB3	2:B:314:MET:CE	2.27	0.59
1:A:470:PRO:HA	1:A:471:PRO:C	2.28	0.57
1:A:335:THR:HG22	1:A:564:HIS:ND1	2.20	0.57
1:A:761:TYR:CD1	1:A:809:ALA:HB1	2.41	0.55
1:A:695:TRP:CE3	1:A:697:LEU:HD11	2.43	0.54
1:A:711:ALA:O	1:A:715:MET:HG2	2.10	0.52
1:A:624:THR:HG22	1:A:799:ALA:HB3	1.92	0.51
1:A:335:THR:HG22	1:A:564:HIS:CE1	2.47	0.50
1:A:661:LYS:HB3	1:A:704:LEU:HD21	1.96	0.47
1:A:661:LYS:HD3	1:A:704:LEU:HD21	1.96	0.47
1:A:707:VAL:CG1	1:A:715:MET:HG3	2.45	0.47
1:A:449:VAL:O	1:A:453:GLU:HG2	2.16	0.46
1:A:801:GLU:HG2	1:A:809:ALA:N	2.26	0.45
1:A:269:ARG:HG2	1:A:271:LYS:O	2.17	0.45
1:A:724:VAL:O	1:A:728:LEU:HG	2.17	0.45
1:A:627:LEU:HD22	1:A:654:MET:HE2	1.99	0.43
1:A:292:ALA:HB2	1:A:815:LEU:HD22	2.01	0.43
1:A:693:LEU:HD23	1:A:706:LEU:HD22	2.00	0.43
1:A:438:GLN:HG3	2:B:352:ILE:HG22	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:662:VAL:HB	1:A:705:ALA:HB3	2.00	0.43
2:B:418:LYS:O	2:B:421:PHE:HB2	2.18	0.43
1:A:801:GLU:CG	1:A:809:ALA:HA	2.45	0.42
1:A:541:ALA:O	1:A:657:GLY:HA3	2.19	0.42
1:A:666:PHE:O	1:A:701:PRO:HG2	2.20	0.42
1:A:439:GLU:HG3	2:B:352:ILE:HD13	2.02	0.41
1:A:330:GLY:HA2	3:A:900:FAD:HM72	2.01	0.41
2:B:428:PHE:O	2:B:429:ASN:C	2.65	0.40
1:A:331:ALA:HA	3:A:900:FAD:C4X	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/666 (100%)	647 (97%)	17 (3%)	0	100	100
2	B	131/235 (56%)	127 (97%)	4 (3%)	0	100	100
All	All	795/901 (88%)	774 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	566/566 (100%)	551 (97%)	15 (3%)	39	65
2	B	117/203 (58%)	112 (96%)	5 (4%)	26	49
All	All	683/769 (89%)	663 (97%)	20 (3%)	37	63

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	392	LEU
1	A	418	LEU
1	A	429	GLU
1	A	469	LYS
1	A	497	LEU
1	A	517	SER
1	A	538	PHE
1	A	563	SER
1	A	565	LEU
1	A	601	GLU
1	A	704	LEU
1	A	706	LEU
1	A	717	ASN
1	A	737	SER
1	A	801	GLU
2	B	316	LEU
2	B	324	VAL
2	B	337	GLN
2	B	341	GLU
2	B	414	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	358	GLN
1	A	395	GLN
1	A	399	ASN
1	A	532	HIS
1	A	633	GLN
1	A	791	GLN
2	B	318	GLN
2	B	411	ASN
2	B	416	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 3 are monoatomic and 1 is modelled with single atom - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	1840	-	5,5,5	0.36	0	5,5,5	0.33	0
5	GOL	A	1839	-	5,5,5	0.41	0	5,5,5	0.41	0
3	FAD	A	900	-	58,58,58	1.10	5 (8%)	85,89,89	1.78	18 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	1840	-	-	0/4/4/4	-
5	GOL	A	1839	-	-	0/4/4/4	-
3	FAD	A	900	-	-	7/34/50/50	0/6/6/6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	900	FAD	C4X-N5	3.80	1.38	1.30
3	A	900	FAD	C10-N1	2.88	1.39	1.33
3	A	900	FAD	C2A-N1A	2.83	1.39	1.33
3	A	900	FAD	C2A-N3A	2.35	1.38	1.33
3	A	900	FAD	C8A-N7A	2.21	1.35	1.31

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	900	FAD	N3A-C2A-N1A	-6.82	118.27	128.58
3	A	900	FAD	N9A-C8A-N7A	-4.88	107.01	113.94
3	A	900	FAD	C4A-N9A-C8A	3.58	109.50	105.74
3	A	900	FAD	C2A-N3A-C4A	3.51	120.40	111.83
3	A	900	FAD	C5A-C4A-N3A	-3.47	121.93	126.72
3	A	900	FAD	C5A-N7A-C8A	3.46	108.89	103.45
3	A	900	FAD	C5'-C4'-C3'	-3.45	105.71	112.22
3	A	900	FAD	C4X-C4-N3	3.15	121.26	113.25
3	A	900	FAD	C4-N3-C2	-2.99	120.33	125.64
3	A	900	FAD	O4-C4-C4X	-2.87	118.97	126.53
3	A	900	FAD	C4A-C5A-N7A	-2.72	107.48	110.58
3	A	900	FAD	C9A-C5X-N5	-2.66	119.63	122.45
3	A	900	FAD	O3P-PA-O1A	-2.59	102.91	110.70
3	A	900	FAD	C10-N1-C2	2.47	122.20	116.85
3	A	900	FAD	C4X-C10-N1	-2.34	118.85	124.59
3	A	900	FAD	C10-C4X-N5	-2.29	120.14	124.81
3	A	900	FAD	N3A-C4A-N9A	2.21	130.93	127.17
3	A	900	FAD	O2P-P-O3P	-2.14	101.50	107.27

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	900	FAD	C5B-O5B-PA-O1A
3	A	900	FAD	C5'-O5'-P-O3P
3	A	900	FAD	C2'-C3'-C4'-C5'
3	A	900	FAD	C2'-C3'-C4'-O4'
3	A	900	FAD	O3'-C3'-C4'-C5'
3	A	900	FAD	C3'-C4'-C5'-O5'
3	A	900	FAD	C2B-C1B-N9A-C8A

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	900	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	666/666 (100%)	0.95	83 (12%) 8 6	18, 70, 84, 105	0
2	B	133/235 (56%)	1.61	37 (27%) 1 1	59, 73, 96, 101	0
All	All	799/901 (88%)	1.06	120 (15%) 5 4	18, 70, 86, 105	0

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	440	GLU	8.5
2	B	380	ASN	7.2
1	A	836	LEU	6.7
2	B	362	LYS	6.3
1	A	172	SER	5.9
1	A	171	PRO	5.8
2	B	379	CYS	5.7
2	B	308	ARG	5.3
2	B	439	ALA	5.1
1	A	471	PRO	5.0
2	B	378	LYS	4.8
2	B	381	ALA	4.5
1	A	835	THR	4.5
1	A	599	GLY	4.3
1	A	275	THR	4.2
2	B	432	GLU	4.2
2	B	383	TRP	4.0
1	A	558	PHE	4.0
2	B	437	TRP	3.8
1	A	833	MET	3.8
2	B	375	VAL	3.8
1	A	276	LYS	3.7
2	B	359	LEU	3.6
2	B	377	GLN	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	277	LYS	3.5
1	A	771	ASN	3.5
1	A	556	ASP	3.4
1	A	834	TYR	3.4
1	A	721	ASP	3.4
1	A	561	THR	3.4
2	B	434	LEU	3.4
2	B	438	GLU	3.4
1	A	278	THR	3.3
2	B	426	ARG	3.3
1	A	597	ALA	3.3
1	A	620	ALA	3.2
2	B	435	GLN	3.2
1	A	769	SER	3.1
1	A	325	TYR	3.1
2	B	365	GLY	3.1
2	B	382	ARG	3.1
1	A	467	GLU	3.0
1	A	279	GLY	3.0
1	A	773	TYR	2.9
2	B	402	PHE	2.9
1	A	563	SER	2.9
1	A	832	ALA	2.9
1	A	527	GLN	2.9
2	B	376	ILE	2.9
2	B	309	LYS	2.8
1	A	683	SER	2.8
1	A	297	LEU	2.8
1	A	560	PHE	2.7
2	B	397	LYS	2.7
1	A	595	TYR	2.6
1	A	632	GLN	2.6
1	A	190	SER	2.6
1	A	549	LEU	2.6
1	A	699	LYS	2.6
1	A	518	ASP	2.6
1	A	667	ASP	2.5
1	A	774	ASP	2.5
2	B	366	GLY	2.5
2	B	364	ASP	2.5
1	A	550	LYS	2.5
2	B	370	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	735	PHE	2.4
1	A	232	GLU	2.4
2	B	374	GLU	2.4
1	A	652	GLN	2.4
1	A	373	GLU	2.4
1	A	738	SER	2.4
1	A	622	LEU	2.4
2	B	400	ARG	2.4
1	A	289	SER	2.4
1	A	557	ASP	2.4
1	A	717	ASN	2.4
1	A	194	ALA	2.4
1	A	372	LYS	2.4
1	A	552	TRP	2.4
1	A	548	SER	2.3
1	A	611	SER	2.3
1	A	352	GLU	2.3
2	B	415	VAL	2.3
1	A	301	GLY	2.3
1	A	619	ASP	2.3
1	A	616	TYR	2.3
1	A	610	THR	2.2
1	A	791	GLN	2.2
2	B	354	GLN	2.2
2	B	418	LYS	2.2
1	A	555	ASP	2.2
1	A	720	ASP	2.2
2	B	361	GLU	2.2
1	A	368	GLN	2.2
1	A	551	HIS	2.2
1	A	767	GLY	2.2
2	B	414	VAL	2.2
1	A	775	LEU	2.2
1	A	200	ILE	2.2
1	A	739	ALA	2.1
1	A	401	LEU	2.1
1	A	565	LEU	2.1
1	A	281	VAL	2.1
1	A	612	GLN	2.1
1	A	204	GLN	2.1
1	A	288	VAL	2.1
2	B	350	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	350	ASN	2.1
1	A	323	GLY	2.1
1	A	768	SER	2.1
1	A	460	GLN	2.1
1	A	770	GLY	2.1
2	B	430	ILE	2.1
1	A	651	VAL	2.0
2	B	436	GLU	2.0
1	A	223	ASP	2.0
1	A	358	GLN	2.0
1	A	668	ARG	2.0
1	A	772	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	1840	6/6	0.76	0.19	113,114,115,115	0
4	CL	A	1841	1/1	0.85	0.15	92,92,92,92	0
6	NH4	A	1842	1/1	0.88	0.22	70,70,70,70	0
4	CL	A	1843	1/1	0.92	0.15	58,58,58,58	1
5	GOL	A	1839	6/6	0.92	0.21	61,62,64,65	0
4	CL	A	1838	1/1	0.92	0.09	53,53,53,53	1
3	FAD	A	900	53/53	0.92	0.14	38,43,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.