



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:27 PM UTC

PDB ID : 4IXU / pdb_00004ixu
Title : Crystal structure of human Arginase-2 complexed with inhibitor 11d: {(5R)-5-amino-5-carboxy-5-[(3-endo)-8-(3,4-dichlorobenzyl)-8-azabicyclo[3.2.1]oct-3-yl]pentyl}(trihydroxy)borate(1-)
Authors : Cousido-Siah, A.; Mitschler, A.; Ruiz, F.X.; Whitehouse, D.; Beckett, P.; Van Zandt, M.C.; Ji, M.K.; Ryder, T.; Jagdmann, R.; Andreoli, M.; Olczak, J.; Mazur, M.; Czystkowski, W.; Piotrowska, W.; Schroeter, H.; Golebiowski, A.; Podjarny, A.
Deposited on : 2013-01-28
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
1	B	306	
1	C	306	

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BME	A	405[B]	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Arginase-2, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	5	13	0
			2383	1505	410	459	9			
1	B	306	Total	C	N	O	S	0	13	0
			2382	1505	409	457	11			
1	C	306	Total	C	N	O	S	0	14	0
			2391	1515	411	454	11			

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		
2	B	2	Total	Mn	0	0
			2	2		
2	C	2	Total	Mn	0	0
			2	2		

- Molecule 3 is BENZAMIDINE (CCD ID: BEN) (formula: C₇H₈N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			9	7	2		
3	B	1	Total	C	N	0	0
			9	7	2		
3	C	1	Total	C	N	0	0
			9	7	2		

- Molecule 4 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	1
			4	2	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	1
			4	2	1	1		
4	B	1	Total	C	O	S	0	1
			4	2	1	1		
4	B	1	Total	C	O	S	0	1
			4	2	1	1		
4	C	1	Total	C	O	S	0	1
			4	2	1	1		
4	C	1	Total	C	O	S	0	1
			4	2	1	1		

- Molecule 5 is {(5R)-5-amino-5-carboxy-5-[(3-endo)-8-(3,4-dichlorobenzyl)-8-azabicyclo[3.2.1]oct-3-yl]pentyl}(trihydroxy)borate(1-) (CCD ID: 38I) (formula: C₂₀H₃₀BCl₂N₂O₅).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	B	C	Cl	N	O	0	0
			30	1	20	2	2	5		
5	B	1	Total	B	C	Cl	N	O	0	0
			30	1	20	2	2	5		
5	C	1	Total	B	C	Cl	N	O	0	0
			30	1	20	2	2	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	283	Total	O	0	9
			293	293		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	294	Total 315	O 315	1	16
6	C	221	Total 231	O 231	0	9



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Arginase-2, mitochondrial



- Molecule 1: Arginase-2, mitochondrial



- Molecule 1: Arginase-2, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.40Å 127.40Å 159.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.34 – 1.90 46.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.34-1.90) 100.0 (46.34-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.167 , 0.193 0.168 , 0.195	Depositor DCC
R_{free} test set	5167 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8142	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 38I, MN, BEN, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	3/2464 (0.1%)	0.79	0/3351
1	B	0.56	1/2459 (0.0%)	0.81	0/3344
1	C	0.57	2/2478 (0.1%)	0.81	2/3370 (0.1%)
All	All	0.58	6/7401 (0.1%)	0.80	2/10065 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	LEU	C-N	-6.41	1.25	1.33
1	C	197	GLY	C-O	-6.21	1.16	1.24
1	B	169	ARG	C-O	-6.17	1.17	1.24
1	C	56	LYS	C-O	-5.49	1.17	1.24
1	A	56	LYS	C-O	-5.26	1.18	1.24
1	A	160	HIS	C-N	-5.19	1.26	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	185	LYS	CA-C-N	5.93	126.27	119.93
1	C	185	LYS	C-N-CA	5.93	126.27	119.93

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	133	HIS	Mainchain
1	C	171[A]	LEU	Mainchain
1	C	171[B]	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2383	0	2374	26	0
1	B	2382	0	2379	25	0
1	C	2391	0	2400	23	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	9	0	7	0	0
3	B	9	0	7	0	0
3	C	9	0	7	0	0
4	A	8	0	12	0	0
4	B	8	0	12	0	0
4	C	8	0	12	2	0
5	A	30	0	27	2	0
5	B	30	0	27	3	0
5	C	30	0	27	2	0
6	A	293	0	0	9	0
6	B	315	0	0	6	0
6	C	231	0	0	1	0
All	All	8142	0	7291	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:172[B]:GLN:HG3	1:C:186:PRO:HG3	1.50	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172[B]:GLN:HG3	1:B:186:PRO:HG3	1.52	0.90
5:A:406:38I:CL2	6:A:659:HOH:O	2.30	0.87
1:A:172[B]:GLN:HG3	1:A:186:PRO:HG3	1.56	0.86
1:C:130:HIS:O	1:C:134[A]:CYS:HB2	1.84	0.78
1:A:187:CYS:O	1:A:188:ILE:HD13	1.84	0.77
1:B:211:ASN:HB2	6:B:558[B]:HOH:O	1.86	0.75
1:A:173:ASP:HB3	6:A:668:HOH:O	1.88	0.74
1:A:87:VAL:HG11	6:A:777:HOH:O	1.88	0.74
1:A:206[B]:HIS:HD2	6:A:768[B]:HOH:O	1.70	0.73
1:C:33[B]:PHE:HE2	1:C:92:VAL:HG22	1.56	0.71
1:A:206[B]:HIS:CD2	6:A:768[B]:HOH:O	2.48	0.64
1:C:40:LYS:HG3	6:C:698[B]:HOH:O	1.97	0.63
1:A:328:GLU:HG2	6:A:506[B]:HOH:O	2.00	0.61
1:A:187:CYS:C	1:A:188:ILE:HD13	2.27	0.60
1:A:87:VAL:CG1	6:A:777:HOH:O	2.47	0.59
1:C:107:VAL:HG23	1:C:113[C]:CYS:SG	2.43	0.59
1:C:33[B]:PHE:CE2	1:C:92:VAL:HG22	2.37	0.58
1:B:107:VAL:HG23	1:B:113[C]:CYS:SG	2.44	0.58
1:B:185:LYS:NZ	6:B:740:HOH:O	2.37	0.58
1:B:202:ASP:OD1	5:B:406:38I:H3	2.05	0.56
1:A:81:LEU:N	6:A:724:HOH:O	2.39	0.55
1:A:306:GLU:O	1:A:310[B]:THR:HG23	2.07	0.55
1:B:137:LEU:HD11	1:B:188:ILE:HD11	1.89	0.55
1:C:202:ASP:OD1	5:C:406:38I:H4	2.07	0.55
1:B:327:ARG:O	1:C:199:ARG:HA	2.07	0.54
1:A:120:HIS:CG	1:A:295:VAL:HG21	2.43	0.53
1:C:147:ASP:HB3	1:C:163:PRO:HD2	1.91	0.53
1:A:327:ARG:O	1:B:199:ARG:HA	2.08	0.53
1:B:306:GLU:O	1:B:310[B]:THR:HG23	2.09	0.53
1:C:133:HIS:C	4:C:405[B]:BME:H12	2.34	0.52
1:B:199:ARG:HD3	1:B:219:MET:HG3	1.92	0.52
1:B:120:HIS:CG	1:B:295:VAL:HG21	2.44	0.52
1:C:199:ARG:HD3	1:C:219:MET:HG3	1.92	0.52
1:B:25:SER:HB3	1:B:111:TYR:CD1	2.46	0.51
1:C:306:GLU:O	1:C:310[B]:THR:HG23	2.11	0.50
1:C:33[B]:PHE:CE2	1:C:92:VAL:HA	2.46	0.50
1:C:55:MET:HE3	1:C:65:LEU:HD13	1.92	0.50
1:B:50:ARG:NH1	1:B:67:ASP:OD2	2.44	0.49
1:B:147:ASP:HB3	1:B:163:PRO:HD2	1.95	0.49
1:C:120:HIS:CG	1:C:295:VAL:HG21	2.48	0.48
5:B:406:38I:H23	5:B:406:38I:H16	1.69	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ARG:HD3	1:A:219:MET:HG3	1.94	0.48
1:A:199:ARG:HA	1:C:327:ARG:O	2.14	0.48
1:A:202:ASP:OD1	5:A:406:38I:H3	2.14	0.48
1:C:185:LYS:HA	1:C:185:LYS:HD3	1.77	0.47
1:B:106:ALA:HB3	1:B:113[C]:CYS:SG	2.55	0.47
1:B:66:LYS:NZ	6:B:731:HOH:O	2.29	0.47
1:B:134[A]:CYS:HB3	1:B:245:PRO:HG3	1.97	0.47
1:C:137:LEU:HD11	1:C:188:ILE:HD11	1.97	0.46
1:A:55:MET:HE3	1:A:65:LEU:HD13	1.96	0.46
1:A:105:ARG:HB3	6:A:581:HOH:O	2.16	0.45
1:B:105:ARG:HB3	6:B:664:HOH:O	2.15	0.45
1:A:147:ASP:HB3	1:A:163:PRO:HD2	1.98	0.45
1:C:130:HIS:CD2	4:C:405[B]:BME:H21	2.53	0.44
1:B:81:LEU:HD21	1:B:87:VAL:HG12	2.00	0.44
1:A:172[B]:GLN:CG	1:A:186:PRO:HG3	2.38	0.43
1:B:202:ASP:OD1	5:B:406:38I:H4	2.18	0.43
1:A:82:TYR:HB3	1:A:86:ILE:HB	1.99	0.43
1:C:82:TYR:HB3	1:C:86:ILE:HB	1.99	0.43
1:C:284:HIS:CD2	1:C:324:GLY:HA2	2.54	0.43
1:C:137:LEU:HD12	1:C:137:LEU:C	2.44	0.42
1:A:172[A]:GLN:HE21	1:A:172[A]:GLN:HB2	1.60	0.42
1:B:105:ARG:NE	6:B:625:HOH:O	2.52	0.42
1:A:87:VAL:HG22	1:A:88:ASN:CG	2.45	0.41
1:A:150:THR:HG22	1:A:165:SER:HB2	2.02	0.41
1:A:87:VAL:HG22	1:A:88:ASN:N	2.35	0.41
1:B:137:LEU:C	1:B:137:LEU:HD12	2.45	0.41
1:B:307:GLU:HG2	6:B:655:HOH:O	2.19	0.41
1:A:185:LYS:HE2	1:A:185:LYS:HB3	1.78	0.41
1:B:105:ARG:HA	1:B:105:ARG:HD2	1.71	0.41
5:C:406:38I:H23	5:C:406:38I:H16	1.83	0.41
1:B:294:LEU:C	1:B:294:LEU:HD13	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/306 (103%)	307 (98%)	6 (2%)	1 (0%)	36	29
1	B	314/306 (103%)	308 (98%)	5 (2%)	1 (0%)	36	29
1	C	316/306 (103%)	310 (98%)	5 (2%)	1 (0%)	36	29
All	All	944/918 (103%)	925 (98%)	16 (2%)	3 (0%)	36	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	GLN
1	C	162	GLN
1	B	162	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/257 (104%)	264 (99%)	3 (1%)	65	67
1	B	267/257 (104%)	265 (99%)	2 (1%)	76	78
1	C	269/257 (105%)	267 (99%)	2 (1%)	76	78
All	All	803/771 (104%)	796 (99%)	7 (1%)	73	73

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ARG
1	A	172[A]	GLN
1	A	172[B]	GLN
1	B	87	VAL
1	B	294	LEU
1	C	294	LEU
1	C	303	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	35	GLN
1	A	64	HIS
1	A	149	ASN
1	A	177	GLN
1	A	228	GLN
1	B	88	ASN
1	B	149	ASN
1	B	228	GLN
1	C	149	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 6 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BME	A	405[B]	-	3,3,3	0.32	0	2,2,2	0.32	0
4	BME	A	404[B]	-	3,3,3	0.27	0	2,2,2	0.27	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BME	C	405[B]	-	3,3,3	0.28	0	2,2,2	0.22	0
3	BEN	B	403	-	9,9,9	0.73	0	7,11,11	0.88	0
3	BEN	C	403	-	9,9,9	0.79	0	7,11,11	0.82	0
4	BME	B	404[B]	-	3,3,3	0.25	0	2,2,2	0.15	0
5	38I	B	406	2	28,32,32	1.16	2 (7%)	35,48,48	2.57	13 (37%)
4	BME	C	404[B]	-	3,3,3	0.25	0	2,2,2	0.41	0
3	BEN	A	403	-	9,9,9	0.65	0	7,11,11	1.47	1 (14%)
5	38I	A	406	2	28,32,32	1.14	3 (10%)	35,48,48	2.15	11 (31%)
5	38I	C	406	2	28,32,32	1.18	3 (10%)	35,48,48	2.01	13 (37%)
4	BME	B	405[B]	-	3,3,3	0.36	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	A	405[B]	-	-	1/1/1/1	-
4	BME	A	404[B]	-	-	1/1/1/1	-
4	BME	C	405[B]	-	-	0/1/1/1	-
3	BEN	B	403	-	-	0/4/4/4	0/1/1/1
3	BEN	C	403	-	-	0/4/4/4	0/1/1/1
4	BME	B	404[B]	-	-	0/1/1/1	-
5	38I	B	406	2	-	5/19/46/46	0/4/3/3
4	BME	C	404[B]	-	-	0/1/1/1	-
3	BEN	A	403	-	-	0/4/4/4	0/1/1/1
5	38I	A	406	2	-	5/19/46/46	0/4/3/3
5	38I	C	406	2	-	5/19/46/46	0/4/3/3
4	BME	B	405[B]	-	-	0/1/1/1	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	406	38I	B11-O30	3.53	1.55	1.47
5	B	406	38I	B11-O30	3.51	1.55	1.47
5	C	406	38I	B11-O30	3.22	1.54	1.47
5	C	406	38I	C26-CL2	2.92	1.80	1.73
5	C	406	38I	C25-CL1	2.87	1.80	1.73
5	B	406	38I	C26-CL2	2.83	1.80	1.73
5	A	406	38I	C25-CL1	2.55	1.79	1.73
5	A	406	38I	C26-CL2	2.45	1.79	1.73

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	406	38I	C22-C21-N16	-7.88	98.89	112.57
5	B	406	38I	C14-C15-N16	5.36	117.27	108.38
5	A	406	38I	C22-C21-N16	-5.25	103.46	112.57
5	A	406	38I	C14-C15-N16	4.89	116.48	108.38
5	C	406	38I	C14-C15-N16	4.87	116.46	108.38
5	B	406	38I	C18-C7-C14	4.68	115.66	108.44
5	C	406	38I	C22-C21-N16	4.40	120.20	112.57
5	A	406	38I	C15-N16-C17	4.26	104.39	100.97
5	B	406	38I	C15-N16-C17	4.24	104.36	100.97
5	A	406	38I	C18-C7-C14	4.21	114.94	108.44
5	C	406	38I	C18-C7-C14	3.88	114.42	108.44
5	B	406	38I	C20-C15-N16	-3.31	98.46	104.12
5	A	406	38I	C14-C15-C20	-3.29	104.85	113.02
5	C	406	38I	C14-C15-C20	-3.16	105.18	113.02
5	C	406	38I	O12-B11-O30	-3.15	96.87	108.94
5	B	406	38I	O3-C1-C4	3.00	121.33	113.53
5	A	406	38I	C20-C15-N16	-2.99	99.02	104.12
5	A	406	38I	O3-C1-C4	2.82	120.87	113.53
5	C	406	38I	O3-C1-C4	2.81	120.84	113.53
5	B	406	38I	C24-C25-CL1	-2.72	113.06	118.42
5	C	406	38I	C18-C17-N16	2.67	112.80	108.38
5	A	406	38I	O12-B11-O30	-2.63	98.87	108.94
5	B	406	38I	C14-C15-C20	-2.59	106.59	113.02
5	C	406	38I	C20-C15-N16	-2.58	99.71	104.12
5	B	406	38I	C23-C22-C27	2.43	121.89	118.55
5	B	406	38I	O12-B11-O30	-2.38	99.82	108.94
3	A	403	BEN	C6-C1-C2	2.34	121.55	118.57
5	C	406	38I	C15-N16-C17	2.30	102.81	100.97
5	B	406	38I	C21-N16-C15	2.27	120.04	114.10
5	C	406	38I	C18-C17-C19	-2.27	107.38	113.02
5	C	406	38I	O2-C1-C4	-2.20	116.86	122.53
5	C	406	38I	C21-N16-C15	2.17	119.77	114.10
5	B	406	38I	C19-C17-N16	-2.17	100.42	104.12
5	B	406	38I	C25-C26-CL2	2.14	125.87	120.84
5	C	406	38I	C26-C27-C22	2.12	121.83	120.46
5	A	406	38I	O12-B11-O13	2.06	116.82	108.94
5	A	406	38I	C8-C9-C10	-2.05	107.98	112.94
5	A	406	38I	O2-C1-C4	-2.01	117.34	122.53

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404[B]	BME	O1-C1-C2-S2
4	A	405[B]	BME	O1-C1-C2-S2
5	A	406	38I	C1-C4-C7-C14
5	A	406	38I	C5-C4-C7-C14
5	A	406	38I	C1-C4-C7-C18
5	A	406	38I	C5-C4-C7-C18
5	B	406	38I	C1-C4-C7-C14
5	B	406	38I	C5-C4-C7-C14
5	B	406	38I	C1-C4-C7-C18
5	B	406	38I	C5-C4-C7-C18
5	C	406	38I	C1-C4-C7-C14
5	C	406	38I	C5-C4-C7-C14
5	C	406	38I	C5-C4-C7-C18
5	A	406	38I	B11-C10-C9-C8
5	C	406	38I	C1-C4-C7-C18
5	B	406	38I	C1-C4-C5-C8
5	C	406	38I	C1-C4-C5-C8


There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	405[B]	BME	2	0
5	B	406	38I	3	0
5	A	406	38I	2	0
5	C	406	38I	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 38I B 406



Ligand 38I A 406





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/306 (100%)	-0.46	5 (1%) 70 74	6, 13, 31, 63	11 (3%)
1	B	306/306 (100%)	-0.46	6 (1%) 65 69	6, 13, 30, 60	10 (3%)
1	C	306/306 (100%)	-0.34	6 (1%) 65 69	7, 16, 35, 63	12 (3%)
All	All	918/918 (100%)	-0.42	17 (1%) 66 70	6, 14, 34, 63	33 (3%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	303	THR	3.7
1	B	24	HIS	3.4
1	A	24	HIS	3.2
1	A	87	VAL	3.2
1	B	81	LEU	3.0
1	B	80	ASP	2.9
1	C	24	HIS	2.9
1	B	105	ARG	2.8
1	A	172[A]	GLN	2.7
1	C	188	ILE	2.7
1	A	173	ASP	2.5
1	A	169	ARG	2.4
1	C	185	LYS	2.2
1	B	211	ASN	2.2
1	C	329	GLY	2.1
1	B	188	ILE	2.0
1	C	211	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BME	A	405[B]	4/4	0.45	0.54	14,26,28,43	4
4	BME	C	404[B]	4/4	0.65	0.31	23,35,40,45	4
4	BME	B	405[B]	4/4	0.69	0.20	17,34,35,63	4
4	BME	C	405[B]	4/4	0.73	0.16	23,28,35,40	4
4	BME	A	404[B]	4/4	0.74	0.31	23,27,34,34	4
4	BME	B	404[B]	4/4	0.76	0.19	34,34,45,47	4
5	38I	B	406	30/30	0.91	0.16	9,29,85,113	0
5	38I	C	406	30/30	0.91	0.16	12,25,97,110	0
5	38I	A	406	30/30	0.92	0.15	9,31,102,127	0
3	BEN	C	403	9/9	0.97	0.05	13,19,22,29	0
3	BEN	A	403	9/9	0.97	0.04	10,14,18,20	0
3	BEN	B	403	9/9	0.98	0.04	13,17,23,24	0
2	MN	A	402	1/1	0.99	0.01	8,8,8,8	0
2	MN	B	402	1/1	0.99	0.01	10,10,10,10	0
2	MN	B	401	1/1	1.00	0.01	8,8,8,8	0
2	MN	A	401	1/1	1.00	0.01	7,7,7,7	0
2	MN	C	401	1/1	1.00	0.01	9,9,9,9	0
2	MN	C	402	1/1	1.00	0.01	8,8,8,8	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 38I B 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 38I C 406:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 38I A 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.