



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:19 PM UTC

PDB ID : 6IXP / pdb_00006ixp
Title : Structure of Myo2-GTD in complex with Mmr1
Authors : Tang, K.; Wei, Z.
Deposited on : 2018-12-11
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1819 (2.76-2.72)
Clashscore	190562	1866 (2.76-2.72)
Ramachandran outliers	187476	1830 (2.76-2.72)
Sidechain outliers	187428	1831 (2.76-2.72)
RSRZ outliers	180081	1819 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	10% 79% 13% 7%
1	D	421	8% 80% 12% 8%
2	B	39	5% 31% 10% 59%
2	C	39	13% 33% 13% 54%
2	E	39	13% 41% 59%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	39	5% 33% 13% 54%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	1	0
			3149	2045	504	590	10			
1	D	386	Total	C	N	O	S	0	1	0
			3121	2027	499	585	10			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1148	GLY	-	expression tag	UNP P19524
A	1149	PRO	-	expression tag	UNP P19524
A	1150	GLY	-	expression tag	UNP P19524
A	1151	SER	-	expression tag	UNP P19524
A	?	-	PHE	deletion	UNP P19524
A	?	-	LEU	deletion	UNP P19524
A	?	-	ASN	deletion	UNP P19524
A	?	-	LYS	deletion	UNP P19524
A	?	-	ILE	deletion	UNP P19524
A	?	-	PHE	deletion	UNP P19524
D	1148	GLY	-	expression tag	UNP P19524
D	1149	PRO	-	expression tag	UNP P19524
D	1150	GLY	-	expression tag	UNP P19524
D	1151	SER	-	expression tag	UNP P19524
D	?	-	PHE	deletion	UNP P19524
D	?	-	LEU	deletion	UNP P19524
D	?	-	ASN	deletion	UNP P19524
D	?	-	LYS	deletion	UNP P19524
D	?	-	ILE	deletion	UNP P19524
D	?	-	PHE	deletion	UNP P19524

- Molecule 2 is a protein called Mitochondrial MYO2 receptor-related protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	S	0	0	0
			120	76	23	20	1			
2	C	18	Total	C	N	O		0	0	0
			146	92	27	27				
2	E	16	Total	C	N	O	S	0	0	0
			110	69	22	18	1			
2	F	18	Total	C	N	O		0	0	0
			146	92	27	27				

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	392	GLY	-	expression tag	UNP Q06324
B	393	PRO	-	expression tag	UNP Q06324
B	394	GLY	-	expression tag	UNP Q06324
B	395	SER	-	expression tag	UNP Q06324
B	396	GLU	-	expression tag	UNP Q06324
B	397	PHE	-	expression tag	UNP Q06324
C	392	GLY	-	expression tag	UNP Q06324
C	393	PRO	-	expression tag	UNP Q06324
C	394	GLY	-	expression tag	UNP Q06324
C	395	SER	-	expression tag	UNP Q06324
C	396	GLU	-	expression tag	UNP Q06324
C	397	PHE	-	expression tag	UNP Q06324
E	392	GLY	-	expression tag	UNP Q06324
E	393	PRO	-	expression tag	UNP Q06324
E	394	GLY	-	expression tag	UNP Q06324
E	395	SER	-	expression tag	UNP Q06324
E	396	GLU	-	expression tag	UNP Q06324
E	397	PHE	-	expression tag	UNP Q06324
F	392	GLY	-	expression tag	UNP Q06324
F	393	PRO	-	expression tag	UNP Q06324
F	394	GLY	-	expression tag	UNP Q06324
F	395	SER	-	expression tag	UNP Q06324
F	396	GLU	-	expression tag	UNP Q06324
F	397	PHE	-	expression tag	UNP Q06324


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Myosin-2




• Molecule 1: Myosin-2





• Molecule 2: Mitochondrial MYO2 receptor-related protein 1








● Molecule 2: Mitochondrial MYO2 receptor-related protein 1



● Molecule 2: Mitochondrial MYO2 receptor-related protein 1



● Molecule 2: Mitochondrial MYO2 receptor-related protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.39Å 63.46Å 169.08Å 90.00° 104.72° 90.00°	Depositor
Resolution (Å)	53.38 – 2.73 53.38 – 2.73	Depositor EDS
% Data completeness (in resolution range)	92.7 (53.38-2.73) 83.2 (53.38-2.73)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.73Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.242 , 0.284 0.248 , 0.289	Depositor DCC
R_{free} test set	1326 reflections (4.37%)	wwPDB-VP
Wilson B-factor (Å ²)	67.5	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 68.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.000 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6796	wwPDB-VP
Average B, all atoms (Å ²)	106.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0360e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/3215	0.74	4/4371 (0.1%)
1	D	0.32	0/3186	0.74	2/4331 (0.0%)
2	B	0.31	0/121	0.86	0/164
2	C	0.24	0/146	0.65	0/195
2	E	0.29	0/111	0.83	0/152
2	F	0.24	0/146	0.66	0/195
All	All	0.31	0/6925	0.74	6/9408 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1549	ILE	CA-C-N	7.04	128.63	119.84
1	D	1549	ILE	C-N-CA	7.04	128.63	119.84
1	A	1549	ILE	CA-C-N	5.89	127.20	119.84
1	A	1549	ILE	C-N-CA	5.89	127.20	119.84
1	A	1485	SER	CA-C-N	5.33	125.63	119.92
1	A	1485	SER	C-N-CA	5.33	125.63	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3149	0	3158	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3121	0	3129	30	0
2	B	120	0	131	2	0
2	C	146	0	161	4	0
2	E	110	0	109	0	0
2	F	146	0	161	4	0
3	A	4	0	0	0	0
All	All	6796	0	6849	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1401:LYS:HB3	1:A:1404:PHE:HB3	1.75	0.67
1:D:1401:LYS:HB3	1:D:1404:PHE:HB3	1.78	0.66
1:A:1480:VAL:HG11	1:A:1486:PRO:HA	1.84	0.59
1:A:1483:TYR:HB2	1:D:1422:GLU:OE1	2.06	0.55
1:D:1391:ASP:OD2	1:D:1438:HIS:ND1	2.30	0.53
1:D:1190:ALA:HA	1:D:1367:TRP:HB3	1.91	0.53
1:D:1480:VAL:HG11	1:D:1486:PRO:HA	1.90	0.53
1:A:1433:THR:HG22	1:A:1437:GLN:NE2	2.24	0.52
2:F:415:VAL:HG12	2:F:419:LYS:HE3	1.90	0.52
1:D:1229:LEU:HD23	1:D:1301:LEU:HD22	1.91	0.52
2:B:417:ASP:OD1	2:B:418:LEU:N	2.42	0.51
1:D:1418:THR:O	1:D:1422:GLU:HG2	2.10	0.51
2:C:415:VAL:HG12	2:C:419:LYS:HE3	1.93	0.51
1:A:1411:LEU:HD21	1:D:1411:LEU:HD11	1.93	0.50
1:A:1449:ARG:O	1:A:1455:ASP:HB3	2.12	0.50
1:A:1247:SER:O	1:A:1250:PHE:HB3	2.12	0.49
1:A:1324:ALA:O	1:A:1328:SER:HB2	2.12	0.49
1:A:1229:LEU:HD23	1:A:1301:LEU:HD22	1.95	0.49
1:A:1457:ASP:HA	1:A:1460:ARG:HD3	1.94	0.49
1:D:1565:VAL:O	1:D:1569:VAL:HG23	2.13	0.49
1:D:1247:SER:O	1:D:1250:PHE:HB3	2.13	0.48
1:A:1190:ALA:HA	1:A:1367:TRP:HB3	1.96	0.48
2:F:408:THR:OG1	2:F:409:ARG:N	2.47	0.48
1:D:1362:PHE:HB3	1:D:1423:TRP:CZ2	2.49	0.47
1:D:1276:LYS:HE3	1:D:1284:TYR:CZ	2.49	0.47
1:A:1565:VAL:O	1:A:1569:VAL:HG23	2.16	0.46
1:D:1494:TYR:CE2	1:D:1498:ILE:HD11	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1317:LEU:HD22	1:A:1321:ALA:HB2	1.99	0.45
1:A:1200:TYR:CG	1:A:1201:PRO:HD3	2.52	0.45
1:A:1433:THR:HG23	1:A:1440:ILE:HD13	1.98	0.45
2:C:408:THR:OG1	2:C:409:ARG:N	2.46	0.45
1:D:1256:ARG:HD2	1:D:1256:ARG:HA	1.70	0.45
1:D:1550:PRO:HD2	1:D:1553:LEU:HD12	1.98	0.45
1:A:1311:LYS:NZ	2:C:422:GLU:OE2	2.50	0.45
1:A:1362:PHE:HB3	1:A:1423:TRP:CZ2	2.52	0.44
1:D:1494:TYR:CZ	1:D:1498:ILE:HD11	2.52	0.44
1:A:1263:VAL:HG22	1:A:1295:LYS:HE2	2.00	0.44
1:A:1494:TYR:CZ	1:A:1498:ILE:HD11	2.52	0.44
1:A:1369:MET:HE1	1:A:1382:VAL:HG21	1.99	0.44
1:A:1391:ASP:OD2	1:A:1438:HIS:ND1	2.40	0.44
1:A:1550:PRO:HD2	1:A:1553:LEU:HD12	2.00	0.44
1:D:1246:PRO:HA	1:D:1368:CYS:SG	2.58	0.43
1:A:1195:LYS:HG2	1:A:1373:HIS:CD2	2.54	0.43
1:D:1180:LYS:HE2	1:D:1234:LYS:HE3	1.99	0.43
1:D:1200:TYR:CG	1:D:1201:PRO:HD3	2.53	0.43
1:D:1407:TRP:HD1	1:D:1447:GLN:HA	1.84	0.43
1:A:1158:GLU:O	1:A:1162:ARG:HG3	2.17	0.43
1:A:1226:ALA:HA	2:C:408:THR:HG22	2.00	0.43
1:A:1246:PRO:HA	1:A:1368:CYS:SG	2.58	0.43
1:A:1170:LEU:HD12	1:A:1170:LEU:HA	1.85	0.42
1:D:1263:VAL:HG22	1:D:1295:LYS:HE2	2.01	0.42
1:D:1311:LYS:HD3	2:F:422:GLU:HG3	2.01	0.42
1:D:1324:ALA:O	1:D:1328:SER:HB2	2.20	0.42
1:A:1524:THR:HA	1:A:1525:PRO:HD3	1.86	0.42
1:A:1158:GLU:HG2	1:A:1162:ARG:HE	1.86	0.41
1:A:1434:GLU:HA	1:A:1437:GLN:HG2	2.01	0.41
1:D:1456:ILE:HG13	1:D:1491:ILE:HG23	2.01	0.41
1:D:1226:ALA:HA	2:F:408:THR:HG22	2.02	0.41
1:D:1485:SER:HA	1:D:1486:PRO:HD3	1.90	0.41
1:A:1174:ILE:HG13	1:A:1224:PHE:HE1	1.86	0.41
1:D:1249:VAL:HA	1:D:1252:LEU:HD12	2.03	0.41
1:D:1317:LEU:HD22	1:D:1321:ALA:HB2	2.03	0.41
1:D:1240:LYS:HB3	1:D:1240:LYS:HE3	1.89	0.41
2:B:403:ILE:HA	2:B:404:PRO:HD3	1.84	0.40
1:D:1377:GLU:OE1	1:D:1377:GLU:N	2.49	0.40
1:A:1276:LYS:HE3	1:A:1284:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/421 (91%)	372 (97%)	13 (3%)	0	100	100
1	D	381/421 (90%)	369 (97%)	12 (3%)	0	100	100
2	B	14/39 (36%)	14 (100%)	0	0	100	100
2	C	16/39 (41%)	16 (100%)	0	0	100	100
2	E	14/39 (36%)	14 (100%)	0	0	100	100
2	F	16/39 (41%)	16 (100%)	0	0	100	100
All	All	826/998 (83%)	801 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/378 (92%)	346 (99%)	3 (1%)	70	82
1	D	346/378 (92%)	343 (99%)	3 (1%)	70	82
2	B	14/33 (42%)	14 (100%)	0	100	100
2	C	17/33 (52%)	17 (100%)	0	100	100
2	E	11/33 (33%)	11 (100%)	0	100	100
2	F	17/33 (52%)	17 (100%)	0	100	100
All	All	754/888 (85%)	748 (99%)	6 (1%)	73	83

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1311	LYS
1	A	1375	GLU
1	A	1483	TYR
1	D	1240	LYS
1	D	1311	LYS
1	D	1375	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1414	ASN
1	D	1447	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	390/421 (92%)	1.10	42 (10%)	11 10	42, 99, 177, 213	3 (0%)
1	D	386/421 (91%)	0.99	35 (9%)	15 14	47, 99, 152, 190	3 (0%)
2	B	16/39 (41%)	1.25	2 (12%)	8 8	87, 116, 150, 155	0
2	C	18/39 (46%)	1.62	5 (27%)	1 1	86, 122, 148, 151	0
2	E	16/39 (41%)	1.52	5 (31%)	1 1	95, 134, 153, 160	0
2	F	18/39 (46%)	1.19	2 (11%)	10 10	87, 121, 148, 158	0
All	All	844/998 (84%)	1.07	91 (10%)	11 10	42, 101, 165, 213	6 (0%)

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1334	PHE	4.2
1	A	1187	ALA	3.9
1	D	1542	PHE	3.8
1	D	1185	PRO	3.7
1	D	1549	ILE	3.6
1	D	1452	THR	3.6
1	A	1411	LEU	3.5
1	D	1330	SER	3.5
1	A	1542	PHE	3.4
2	F	408	THR	3.3
1	A	1462	ILE	3.3
1	D	1187	ALA	3.2
2	C	408	THR	3.1
1	A	1452	THR	3.1
1	A	1456	ILE	3.1
1	D	1190	ALA	3.0
2	E	417	ASP	3.0
1	A	1453	ILE	3.0
1	A	1447	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	1551	ALA	2.9
1	D	1275	PHE	2.9
1	D	1412	GLN	2.9
1	A	1152	ASN	2.9
1	D	1189	VAL	2.9
2	C	421	ILE	2.9
1	D	1568	GLN	2.9
1	A	1287	TYR	2.9
1	A	1483	TYR	2.9
1	D	1191	ILE	2.8
1	D	1407	TRP	2.8
1	A	1448	VAL	2.8
2	E	416	LEU	2.7
1	A	1184	VAL	2.7
1	D	1152	ASN	2.7
1	A	1185	PRO	2.7
1	A	1227	GLN	2.6
1	D	1227	GLN	2.6
1	D	1492	LEU	2.6
1	A	1437	GLN	2.6
1	A	1463	CYS	2.5
1	A	1271	THR	2.5
1	A	1196	ARG	2.5
1	A	1504	ALA	2.5
1	A	1281	ASP	2.5
1	D	1400	MET	2.5
1	D	1321	ALA	2.5
1	D	1451	TYR	2.4
1	A	1249	VAL	2.4
1	D	1458	ILE	2.4
2	B	418	LEU	2.4
1	A	1223	SER	2.4
1	A	1327	ILE	2.4
1	A	1459	LEU	2.4
1	D	1462	ILE	2.4
1	A	1378	VAL	2.3
1	D	1504	ALA	2.3
1	A	1278	GLY	2.3
1	D	1158	GLU	2.3
1	D	1223	SER	2.3
2	C	416	LEU	2.3
1	A	1451	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1449	ARG	2.2
1	D	1278	GLY	2.2
1	A	1331	LEU	2.2
1	A	1464	TYR	2.2
2	E	403	ILE	2.2
1	A	1300	ALA	2.2
1	D	1368	CYS	2.2
1	A	1540	ARG	2.2
1	D	1432	GLY	2.2
1	D	1287	TYR	2.2
1	A	1485	SER	2.2
2	B	415	VAL	2.2
1	A	1190	ALA	2.1
2	E	412	ARG	2.1
2	E	418	LEU	2.1
1	D	1294	LEU	2.1
1	D	1505	LEU	2.1
1	A	1460	ARG	2.1
1	A	1274	THR	2.1
2	C	424	GLN	2.1
1	A	1475	ILE	2.1
1	A	1275	PHE	2.1
1	D	1186	ASP	2.1
1	D	1272	GLU	2.1
1	A	1538	LYS	2.1
1	A	1484	GLU	2.0
1	D	1483	TYR	2.0
2	C	415	VAL	2.0
1	D	1457	ASP	2.0
2	F	425	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.