



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:11 PM UTC

PDB ID : 6IXZ / pdb_00006ixz
Title : X-ray structure of sortase C from Clostridium perfringens SM101
Authors : Kamitori, S.; Tamai, E.
Deposited on : 2018-12-12
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	6% 68% 16% • 14%
1	B	220	10% 62% 16% • 20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sortase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	0	0
			1511	963	244	303	1			
1	B	177	Total	C	N	O	S	0	0	0
			1413	902	225	285	1			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	MET	-	initiating methionine	UNP Q0SWL7
A	253	GLU	-	expression tag	UNP Q0SWL7
A	254	HIS	-	expression tag	UNP Q0SWL7
A	255	HIS	-	expression tag	UNP Q0SWL7
A	256	HIS	-	expression tag	UNP Q0SWL7
A	257	HIS	-	expression tag	UNP Q0SWL7
A	258	HIS	-	expression tag	UNP Q0SWL7
A	259	HIS	-	expression tag	UNP Q0SWL7
B	40	MET	-	initiating methionine	UNP Q0SWL7
B	253	GLU	-	expression tag	UNP Q0SWL7
B	254	HIS	-	expression tag	UNP Q0SWL7
B	255	HIS	-	expression tag	UNP Q0SWL7
B	256	HIS	-	expression tag	UNP Q0SWL7
B	257	HIS	-	expression tag	UNP Q0SWL7
B	258	HIS	-	expression tag	UNP Q0SWL7
B	259	HIS	-	expression tag	UNP Q0SWL7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	49	Total	O	0	0
			49	49		
2	B	35	Total	O	0	0
			35	35		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sortase family protein



• Molecule 1: Sortase family protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.45Å 63.41Å 76.53Å 90.00° 101.39° 90.00°	Depositor
Resolution (Å)	19.63 – 1.98 19.63 – 1.98	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.63-1.98) 98.7 (19.63-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.231 , 0.283 0.237 , 0.285	Depositor DCC
R_{free} test set	1357 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3008	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/1537	0.80	0/2086
1	B	0.61	0/1437	0.80	0/1951
All	All	0.62	0/2974	0.80	0/4037

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1511	0	1504	27	0
1	B	1413	0	1402	31	0
2	A	49	0	0	4	0
2	B	35	0	0	1	0
All	All	3008	0	2906	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:LEU:HD11	1:B:127:VAL:CG2	2.13	0.79

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:LEU:HD11	1:B:127:VAL:HG23	1.71	0.72
1:B:150:THR:CG2	1:B:214:TYR:HD2	2.04	0.70
1:A:150:THR:HG23	2:A:337:HOH:O	1.90	0.70
1:A:179:LEU:HD22	1:A:227:ARG:CZ	2.22	0.70
1:A:189:VAL:HG13	1:A:193:GLU:HB2	1.74	0.68
1:B:150:THR:HG22	1:B:214:TYR:HD2	1.59	0.67
1:B:150:THR:HG22	1:B:214:TYR:CD2	2.29	0.67
1:A:191:PRO:HA	1:A:220:ARG:HD2	1.75	0.67
1:B:108:ILE:O	1:B:110:ILE:HD13	1.97	0.65
1:B:133:THR:HG21	1:B:145:VAL:HG13	1.80	0.64
1:A:182:LYS:HB2	1:A:228:THR:HG21	1.80	0.63
1:B:194:ILE:HA	1:B:197:LEU:HD12	1.83	0.60
1:A:190:HIS:HB3	2:A:340:HOH:O	2.02	0.59
1:A:183:VAL:HG11	1:A:186:ILE:HD11	1.86	0.57
1:B:209:LEU:HD13	1:B:222:LEU:HD21	1.85	0.57
1:B:112:LEU:CD1	1:B:127:VAL:HG23	2.35	0.56
1:A:165:LYS:HD3	1:A:166:ASP:O	2.07	0.55
1:B:150:THR:C	1:B:159:THR:HG22	2.32	0.55
1:B:209:LEU:HD13	1:B:222:LEU:CD2	2.37	0.55
1:A:190:HIS:NE2	1:B:120:VAL:HA	2.23	0.54
1:B:150:THR:O	1:B:159:THR:HG22	2.09	0.52
1:A:143:HIS:HA	1:A:205:TYR:O	2.10	0.52
1:B:179:LEU:HD12	1:B:179:LEU:N	2.24	0.51
1:A:209:LEU:C	1:A:209:LEU:HD23	2.36	0.51
1:B:193:GLU:HB3	1:B:196:GLU:CG	2.41	0.50
1:B:181:TYR:CE1	1:B:206:VAL:HG13	2.47	0.49
1:A:191:PRO:HD2	2:A:340:HOH:O	2.13	0.49
1:B:181:TYR:CZ	1:B:206:VAL:HG13	2.48	0.49
1:B:114:ILE:HD12	1:B:174:ILE:HD11	1.95	0.48
1:A:182:LYS:HB2	1:A:228:THR:CG2	2.43	0.48
1:B:127:VAL:HG22	1:B:148:ALA:HB2	1.94	0.48
1:B:193:GLU:HB3	1:B:196:GLU:HG2	1.96	0.48
1:A:61:LYS:O	1:A:62:ASP:C	2.58	0.46
1:B:150:THR:CG2	1:B:214:TYR:CD2	2.90	0.46
1:B:109:ASN:ND2	2:B:302:HOH:O	2.48	0.46
1:B:119:SER:H	1:B:122:ILE:HD12	1.81	0.46
1:A:61:LYS:O	1:A:64:GLU:N	2.48	0.46
1:B:112:LEU:CD1	1:B:127:VAL:CG2	2.89	0.46
1:A:44:LYS:O	1:A:47:VAL:HG12	2.17	0.44
1:A:54:THR:O	1:A:55:GLU:C	2.59	0.44
1:B:112:LEU:HD11	1:B:127:VAL:HG21	1.93	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:THR:CG2	2:A:337:HOH:O	2.59	0.42
1:A:190:HIS:CE1	1:B:120:VAL:HG22	2.55	0.42
1:A:71:ASN:HB3	1:A:92:LEU:CD2	2.49	0.42
1:B:112:LEU:HB2	1:B:113:PRO:HD2	2.01	0.42
1:A:103:ILE:HB	1:A:114:ILE:HD11	2.01	0.42
1:B:180:ALA:HB2	1:B:230:LEU:HD12	2.02	0.42
1:A:95:HIS:O	1:A:96:GLU:C	2.62	0.41
1:B:192:ASP:O	1:B:194:ILE:N	2.53	0.41
1:A:175:LEU:O	1:A:176:LYS:HB2	2.21	0.41
1:A:116:TYR:CD2	1:A:135:LEU:HD21	2.56	0.41
1:A:140:GLU:CG	1:A:202:ASP:HA	2.51	0.41
1:B:65:VAL:HG11	1:B:176:LYS:HE2	2.03	0.41
1:A:179:LEU:CD2	1:A:227:ARG:NH2	2.84	0.40
1:A:165:LYS:C	1:A:165:LYS:CD	2.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/220 (86%)	182 (97%)	6 (3%)	0	100	100
1	B	173/220 (79%)	166 (96%)	6 (4%)	1 (1%)	21	12
All	All	361/440 (82%)	348 (96%)	12 (3%)	1 (0%)	36	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	193	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/202 (85%)	163 (95%)	9 (5%)	21	9
1	B	162/202 (80%)	147 (91%)	15 (9%)	8	2
All	All	334/404 (83%)	310 (93%)	24 (7%)	13	5

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	LYS
1	A	93	ASN
1	A	96	GLU
1	A	99	VAL
1	A	165	LYS
1	A	175	LEU
1	A	189	VAL
1	A	218	THR
1	A	219	GLU
1	B	47	VAL
1	B	63	GLU
1	B	81	ASN
1	B	85	GLU
1	B	87	GLN
1	B	110	ILE
1	B	124	LYS
1	B	127	VAL
1	B	140	GLU
1	B	145	VAL
1	B	150	THR
1	B	172	LEU
1	B	177	LYS
1	B	195	ASP
1	B	206	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	ASN
1	A	109	ASN
1	A	217	ASN
1	B	71	ASN
1	B	81	ASN
1	B	87	GLN
1	B	109	ASN
1	B	149	HIS
1	B	190	HIS
1	B	217	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/220 (86%)	0.52	13 (6%) 23 31	15, 27, 50, 62	0
1	B	177/220 (80%)	0.76	22 (12%) 8 11	16, 27, 62, 84	0
All	All	367/440 (83%)	0.64	35 (9%) 14 19	15, 27, 58, 84	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	47	VAL	7.0
1	B	120	VAL	6.3
1	B	197	LEU	5.7
1	B	194	ILE	4.4
1	B	196	GLU	3.8
1	B	150	THR	3.8
1	A	46	ALA	3.8
1	B	123	LEU	3.7
1	B	230	LEU	3.4
1	B	159	THR	2.9
1	A	190	HIS	2.9
1	B	60	VAL	2.8
1	B	122	ILE	2.8
1	B	158	PHE	2.7
1	B	119	SER	2.7
1	B	124	LYS	2.7
1	A	91	TYR	2.6
1	B	149	HIS	2.6
1	B	63	GLU	2.5
1	A	231	SER	2.4
1	B	148	ALA	2.4
1	B	57	ILE	2.3
1	B	195	ASP	2.3
1	A	45	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	93	ASN	2.3
1	A	47	VAL	2.3
1	A	74	LYS	2.3
1	A	194	ILE	2.2
1	A	230	LEU	2.2
1	B	61	LYS	2.2
1	A	43	ASN	2.1
1	B	55	GLU	2.1
1	B	214	TYR	2.1
1	A	87	GLN	2.1
1	A	86	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.