



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:35 AM UTC

PDB ID : 2IZL / pdb_00002izl
Title : STREPTAVIDIN-2-IMINOBIOTIN PH 7.3 I222 COMPLEX
Authors : Katz, B.A.
Deposited on : 1997-08-13
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7025 (1.50-1.46)
Ramachandran outliers	187476	6917 (1.50-1.46)
Sidechain outliers	187428	6914 (1.50-1.46)
RSRZ outliers	180081	6781 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	B	123	3% 72% 24% ..
1	D	123	3% 75% 21% ..

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STREPTAVIDIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	122	Total 1832	C 590	H 898	N 159	O 185	6	5	0
1	D	121	Total 1767	C 569	H 857	N 157	O 184	10	2	0

- Molecule 2 is 2-IMINOBIOTIN (CCD ID: IMI) (formula: $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total 33	C 10	H 17	N 3	O 2	S 1	0	0
2	D	1	Total 32	C 10	H 16	N 3	O 2	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	79	Total 237	H 158	O 79	0	4
3	D	119	Total 357	H 238	O 119	0	9


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: STREPTAVIDIN



• Molecule 1: STREPTAVIDIN



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	95.41Å 106.54Å 48.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.50 – 1.48 7.50 – 1.50	Depositor EDS
% Data completeness (in resolution range)	86.0 (7.50-1.48) 88.2 (7.50-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 1.50Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.185 , 0.211 0.196 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 60.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4258	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IMI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.91	11/976 (1.1%)	1.75	21/1333 (1.6%)
1	D	1.96	7/940 (0.7%)	1.77	18/1287 (1.4%)
All	All	1.94	18/1916 (0.9%)	1.76	39/2620 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	87	HIS	CG-ND1	-7.96	1.29	1.38
1	B	127	HIS	CG-ND1	-7.44	1.30	1.38
1	D	127	HIS	CG-ND1	-7.38	1.30	1.38
1	D	87	HIS	CG-ND1	-7.21	1.30	1.38
1	D	111	THR	CA-CB	6.11	1.62	1.53
1	B	79	TRP	CA-CB	6.01	1.60	1.52
1	B	98	GLY	N-CA	5.84	1.50	1.44
1	D	93	SER	N-CA	5.57	1.53	1.46
1	B	78	ALA	N-CA	5.38	1.52	1.46
1	B	87	HIS	CD2-NE2	-5.26	1.32	1.37
1	B	109	LEU	CA-C	5.15	1.58	1.52
1	B	105	ASN	N-CA	5.11	1.52	1.46
1	D	77	VAL	N-CA	5.11	1.52	1.46
1	B	45	SER	CA-CB	5.10	1.61	1.53
1	B	22[A]	TYR	N-CA	5.09	1.52	1.45
1	B	22[B]	TYR	N-CA	5.09	1.52	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	51	GLU	N-CA	5.07	1.52	1.46
1	D	79	TRP	CA-CB	5.05	1.58	1.52

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ALA	N-CA-C	-8.96	94.63	109.24
1	D	78	ALA	N-CA-C	-8.75	94.98	109.24
1	D	115	THR	N-CA-C	-7.95	99.06	110.59
1	B	115	THR	N-CA-C	-7.87	99.92	110.55
1	B	123	THR	N-CA-C	7.55	120.71	108.41
1	D	123	THR	N-CA-C	7.51	121.14	108.90
1	D	85	ASN	CA-CB-CG	-7.11	105.49	112.60
1	B	104	ILE	N-CA-C	-7.00	97.40	107.77
1	B	107	GLN	CA-CB-CG	-6.61	100.89	114.10
1	B	81	ASN	CA-CB-CG	6.46	119.06	112.60
1	D	107[A]	GLN	N-CA-C	-6.27	98.77	109.24
1	D	107[B]	GLN	N-CA-C	-6.27	98.77	109.24
1	B	107	GLN	N-CA-C	-6.18	98.92	109.24
1	D	75	TRP	CG-CD1-NE1	-6.08	102.30	110.20
1	B	128	ASP	CA-CB-CG	6.05	118.65	112.60
1	D	79	TRP	N-CA-C	6.04	120.06	110.70
1	B	21	TRP	NE1-CE2-CZ2	5.98	139.07	130.10
1	B	92	TRP	CG-CD1-NE1	-5.88	102.55	110.20
1	B	75	TRP	CG-CD1-NE1	-5.75	102.73	110.20
1	B	95	GLN	OE1-CD-NE2	-5.59	117.01	122.60
1	D	75	TRP	NE1-CE2-CZ2	5.59	138.48	130.10
1	B	75	TRP	NE1-CE2-CZ2	5.54	138.41	130.10
1	D	106	THR	N-CA-C	5.49	117.53	109.24
1	D	120	TRP	CG-CD1-NE1	-5.46	103.10	110.20
1	B	79	TRP	N-CA-C	5.39	119.05	110.70
1	B	92	TRP	CD1-NE1-CE2	5.39	118.60	108.90
1	B	85	ASN	CA-CB-CG	-5.37	107.23	112.60
1	D	92	TRP	CG-CD1-NE1	-5.33	103.27	110.20
1	D	21	TRP	NE1-CE2-CZ2	5.27	138.00	130.10
1	B	29	PHE	CA-CB-CG	5.24	119.03	113.80
1	B	21	TRP	CG-CD1-NE1	-5.20	103.44	110.20
1	D	92	TRP	CD1-NE1-CE2	5.19	118.24	108.90
1	D	21	TRP	CG-CD1-NE1	-5.15	103.51	110.20
1	D	79	TRP	CG-CD1-NE1	-5.14	103.52	110.20
1	B	108	TRP	CG-CD1-NE1	-5.13	103.53	110.20
1	B	120	TRP	CG-CD1-NE1	-5.11	103.56	110.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	95	GLN	OE1-CD-NE2	-5.10	117.50	122.60
1	B	113	GLY	N-CA-C	-5.03	102.30	111.15
1	D	31	VAL	N-CA-C	5.02	117.23	108.95

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	ARG	Sidechain

CLOSE-CONTACTS INFOmissingINFO

5.2 Torsion angles

5.2.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	124/123 (101%)	121 (98%)	2 (2%)	1 (1%)	16	3
1	D	121/123 (98%)	116 (96%)	1 (1%)	4 (3%)	3	0
All	All	245/246 (100%)	237 (97%)	3 (1%)	5 (2%)	5	1

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	14	GLU
1	D	15	ALA
1	D	16	GLY
1	D	52	SER
1	B	52	SER

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	94/90 (104%)	93 (99%)	1 (1%)	65	40
1	D	90/90 (100%)	86 (96%)	4 (4%)	25	4
All	All	184/180 (102%)	179 (97%)	5 (3%)	44	11

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	66	THR
1	D	36	ASP
1	D	107[A]	GLN
1	D	107[B]	GLN
1	D	110	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.5 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IMI	D	300	-	16,17,17	1.82	7 (43%)	19,23,23	2.21	6 (31%)
2	IMI	B	300	-	16,17,17	1.58	5 (31%)	19,23,23	2.21	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMI	D	300	-	-	3/7/28/28	0/2/2/2
2	IMI	B	300	-	-	2/7/28/28	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	IMI	C3-N1	-3.08	1.30	1.35
2	D	300	IMI	C7-C2	2.85	1.60	1.52
2	D	300	IMI	C3-N2	-2.61	1.30	1.35
2	D	300	IMI	C3-N1	-2.53	1.30	1.35
2	B	300	IMI	O12-C11	-2.42	1.22	1.30
2	D	300	IMI	C10-C11	2.41	1.56	1.50
2	B	300	IMI	C3-N2	-2.38	1.31	1.35
2	B	300	IMI	C5-N1	2.31	1.49	1.46
2	D	300	IMI	C4-N2	2.29	1.50	1.45
2	D	300	IMI	C5-N1	2.27	1.49	1.46
2	B	300	IMI	C10-C11	2.17	1.55	1.50
2	D	300	IMI	O12-C11	-2.17	1.23	1.30

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	300	IMI	C4-C2-S1	4.81	110.50	105.03
2	D	300	IMI	C5-C6-S1	4.80	112.67	106.06
2	B	300	IMI	C4-C2-S1	4.69	110.36	105.03
2	B	300	IMI	C5-C6-S1	3.95	111.50	106.06
2	B	300	IMI	C6-C5-N1	-3.90	108.15	113.18
2	D	300	IMI	C6-C5-N1	-3.83	108.24	113.18
2	B	300	IMI	N1-C3-N2	3.68	112.96	109.21
2	B	300	IMI	C2-C4-N2	-3.21	109.95	113.34
2	D	300	IMI	N1-C3-N2	2.81	112.06	109.21
2	D	300	IMI	C7-C2-S1	2.24	119.25	112.78
2	B	300	IMI	C8-C7-C2	-2.19	109.00	114.04
2	D	300	IMI	C8-C7-C2	-2.17	109.06	114.04

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	300	IMI	C4-C2-C7-C8
2	D	300	IMI	C4-C2-C7-C8
2	D	300	IMI	C9-C10-C11-O11
2	D	300	IMI	C9-C10-C11-O12
2	B	300	IMI	C9-C10-C11-O11

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Other polymers [i](#)

There are no such residues in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	B	122/123 (99%)	-0.36	4 (3%)	49	53	7, 20, 32, 67	16 (13%)
1	D	120/123 (97%)	-0.43	4 (3%)	49	53	6, 20, 36, 44	12 (10%)
All	All	242/246 (98%)	-0.40	8 (3%)	49	53	6, 20, 35, 67	28 (11%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	14	GLU	3.1
1	B	134[A]	LYS	2.8
1	B	67	ASP	2.4
1	B	100	ALA	2.3
1	D	13	ALA	2.2
1	D	68	GLY	2.2
1	D	133	VAL	2.2
1	B	15	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IMI	B	300	16/16	0.97	0.04	19,21,24,25	0
2	IMI	D	300	16/16	0.97	0.04	20,22,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.