



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 01:20 PM UTC

PDB ID : 5IZC / pdb_00005izc
Title : Trypanosoma brucei PTR1 in complex with inhibitor F032
Authors : Pozzi, C.; Landi, G.; Di Pisa, F.; Mangani, S.
Deposited on : 2016-03-25
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	
1	B	268	
1	D	268	
2	C	268	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pteridine reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	5	0
			1864	1178	325	348	13			
1	B	247	Total	C	N	O	S	0	6	0
			1868	1181	325	350	12			
1	D	248	Total	C	N	O	S	0	3	0
			1839	1160	322	345	12			

- Molecule 2 is a protein called Pteridine reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	241	Total	C	N	O	S	0	5	0
			1800	1133	312	343	12			

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is N 2 -[(thiophen-2-yl)methyl]-1,3,4-thiadiazole-2,5-diamine (CCD ID: 6F4) (formula: C₇H₈N₄S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	S	0	0
			13	7	4	2		
4	B	1	Total	C	N	S	0	0
			13	7	4	2		
4	C	1	Total	C	N	S	0	0
			13	7	4	2		
4	D	1	Total	C	N	S	0	0
			13	7	4	2		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	135	Total 135	O 135	0	0
7	B	164	Total 164	O 164	0	0
7	C	141	Total 141	O 141	0	0
7	D	133	Total 133	O 133	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pteridine reductase




- Molecule 1: Pteridine reductase




- Molecule 1: Pteridine reductase



- Molecule 2: Pteridine reductase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.63Å 89.44Å 82.64Å 90.00° 115.87° 90.00°	Depositor
Resolution (Å)	74.36 – 1.92 74.36 – 1.92	Depositor EDS
% Data completeness (in resolution range)	97.6 (74.36-1.92) 97.6 (74.36-1.92)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.161 , 0.206 0.172 , 0.216	Depositor DCC
R_{free} test set	3668 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8206	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.70 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6034e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6F4, CME, OCS, ACT, GOL, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.19	2/1895 (0.1%)	1.07	1/2572 (0.0%)
1	B	1.25	5/1902 (0.3%)	1.09	0/2581
1	D	1.17	1/1864 (0.1%)	1.08	1/2532 (0.0%)
2	C	1.23	3/1820 (0.2%)	1.10	0/2472
All	All	1.21	11/7481 (0.1%)	1.09	2/10157 (0.0%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	130	ALA	C-O	-6.06	1.18	1.24
1	B	200	ASN	CA-C	-5.68	1.45	1.52
1	B	57	VAL	N-CA	5.61	1.52	1.46
1	B	193	ALA	C-O	-5.60	1.19	1.24
2	C	164	VAL	N-CA	-5.51	1.39	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	SER	N-CA-C	5.72	118.31	110.24
1	D	250	GLN	N-CA-C	5.52	119.12	112.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1864	0	1901	11	0
1	B	1868	0	1911	14	0
1	D	1839	0	1861	14	0
2	C	1800	0	1804	16	0
3	A	48	0	25	0	0
3	B	48	0	25	2	0
3	C	48	0	25	3	0
3	D	48	0	25	1	0
4	A	13	0	0	0	0
4	B	13	0	0	0	0
4	C	13	0	0	1	0
4	D	13	0	0	0	0
5	A	8	0	6	1	0
5	C	4	0	3	0	0
6	B	6	0	8	0	0
7	A	135	0	0	0	0
7	B	164	0	0	1	0
7	C	141	0	0	5	0
7	D	133	0	0	1	0
All	All	8206	0	7594	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:ASN:OD1	1:A:141:ARG:NH1	2.30	0.62
1:D:136:MET:HE3	7:D:527:HOH:O	1.99	0.61
1:B:164:VAL:HG22	1:B:179:HIS:CD2	2.39	0.58
1:A:133:LEU:HD23	1:A:136[A]:MET:HE2	1.85	0.58
1:A:65:ASN:HA	1:A:69:LEU:HD22	1.85	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/268 (92%)	240 (97%)	7 (3%)	0	100	100
1	B	246/268 (92%)	238 (97%)	8 (3%)	0	100	100
1	D	244/268 (91%)	236 (97%)	8 (3%)	0	100	100
2	C	236/268 (88%)	227 (96%)	9 (4%)	0	100	100
All	All	973/1072 (91%)	941 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/213 (92%)	194 (98%)	3 (2%)	57	47
1	B	199/213 (93%)	193 (97%)	6 (3%)	36	20
1	D	193/213 (91%)	191 (99%)	2 (1%)	68	64
2	C	188/212 (89%)	186 (99%)	2 (1%)	65	60
All	All	777/851 (91%)	764 (98%)	13 (2%)	55	41

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	220	LYS
1	B	250	GLN
1	D	179	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	190	LEU
1	D	166	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	175	ASN
1	D	67	ASN
1	D	236	GLN
1	D	127	ASN
1	D	179	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	A	168	1	8,9,10	0.91	1 (12%)	6,9,11	1.48	1 (16%)
1	CME	B	168	1	8,9,10	0.86	0	6,9,11	1.38	1 (16%)
2	CME	C	168	2	8,9,10	0.74	0	6,9,11	1.62	1 (16%)
1	CME	D	168	1	8,9,10	0.66	0	6,9,11	1.44	1 (16%)
2	OCS	C	59	2	6,8,9	1.92	1 (16%)	7,11,13	2.16	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	168	1	-	2/5/8/10	-
1	CME	B	168	1	-	2/5/8/10	-
2	CME	C	168	2	-	0/5/8/10	-
1	CME	D	168	1	-	0/5/8/10	-
2	OCS	C	59	2	-	1/4/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	59	OCS	OD3-SG	3.91	1.56	1.45
1	A	168	CME	CB-SG	-2.09	1.74	1.81

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	59	OCS	OD3-SG-CB	4.12	112.90	106.76
2	C	168	CME	CB-SG-SD	-3.55	94.66	103.86
1	B	168	CME	CB-SG-SD	-3.24	95.47	103.86
1	D	168	CME	CB-SG-SD	-3.22	95.52	103.86
1	A	168	CME	CA-CB-SG	-2.98	102.21	114.45

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	168	CME	SD-CE-CZ-OH
2	C	59	OCS	N-CA-CB-SG
1	B	168	CME	SD-CE-CZ-OH
1	B	168	CME	CZ-CE-SD-SG
1	A	168	CME	CA-CB-SG-SD

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	168	CME	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	A	301	-	50,52,52	1.60	10 (20%)	71,80,80	1.67	16 (22%)
3	NAP	B	301	-	50,52,52	1.68	9 (18%)	71,80,80	2.02	20 (28%)
4	6F4	C	302	-	14,14,14	1.75	2 (14%)	16,18,18	2.48	6 (37%)
5	ACT	A	303	-	3,3,3	0.78	0	3,3,3	1.15	0
5	ACT	C	303	-	3,3,3	0.77	0	3,3,3	0.71	0
4	6F4	B	302	-	14,14,14	2.19	5 (35%)	16,18,18	2.90	7 (43%)
3	NAP	C	301	-	50,52,52	1.59	7 (14%)	71,80,80	1.99	21 (29%)
6	GOL	B	303	-	5,5,5	0.98	0	5,5,5	2.04	2 (40%)
4	6F4	D	302	-	14,14,14	2.10	4 (28%)	16,18,18	3.31	7 (43%)
4	6F4	A	302	-	14,14,14	1.65	4 (28%)	16,18,18	2.51	6 (37%)
5	ACT	A	304	-	3,3,3	1.05	0	3,3,3	1.20	0
3	NAP	D	301	-	50,52,52	1.74	11 (22%)	71,80,80	1.63	18 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	301	-	-	0/35/67/67	0/5/5/5
3	NAP	B	301	-	-	0/35/67/67	0/5/5/5
4	6F4	C	302	-	-	0/5/5/5	0/2/2/2
6	GOL	B	303	-	-	3/4/4/4	-
4	6F4	B	302	-	-	3/5/5/5	0/2/2/2
3	NAP	C	301	-	-	0/35/67/67	0/5/5/5
4	6F4	D	302	-	-	2/5/5/5	0/2/2/2
4	6F4	A	302	-	-	2/5/5/5	0/2/2/2
3	NAP	D	301	-	-	0/35/67/67	0/5/5/5

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	301	NAP	C5A-C4A	6.27	1.50	1.39
3	D	301	NAP	PA-O3	6.08	1.66	1.59
3	B	301	NAP	PA-O3	6.02	1.66	1.59
3	B	301	NAP	C5A-C4A	4.96	1.47	1.39
3	D	301	NAP	C5A-C4A	4.89	1.47	1.39

The worst 5 of 103 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	NAP	C2B-C1B-N9A	-6.99	102.24	113.75
4	B	302	6F4	SAJ-CAH-NAG	6.93	128.55	121.32
4	D	302	6F4	SAJ-CAK-NAL	-6.55	109.09	113.79
3	C	301	NAP	C3N-C7N-N7N	5.93	125.04	117.74
4	D	302	6F4	SAJ-CAH-NAI	-5.67	108.19	114.52

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	6F4	SAJ-CAH-NAG-CAF
4	B	302	6F4	CAB-CAC-CAF-NAG
4	B	302	6F4	SAD-CAC-CAF-NAG
6	B	303	GOL	O1-C1-C2-C3
4	A	302	6F4	NAI-CAH-NAG-CAF


There are no ring outliers.


5 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	NAP	2	0
4	C	302	6F4	1	0
5	A	303	ACT	1	0
3	C	301	NAP	3	0
3	D	301	NAP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/268 (92%)	-0.25	2 (0%) 82 88	9, 17, 37, 58	7 (2%)
1	B	246/268 (91%)	-0.35	3 (1%) 76 82	7, 16, 35, 52	13 (5%)
1	D	247/268 (92%)	-0.17	6 (2%) 59 66	7, 18, 40, 65	8 (3%)
2	C	239/268 (89%)	-0.17	8 (3%) 49 54	9, 18, 41, 59	9 (3%)
All	All	980/1072 (91%)	-0.24	19 (1%) 66 73	7, 17, 38, 65	37 (3%)

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	210	PRO	3.8
1	B	211	VAL	3.8
2	C	208	LEU	3.7
2	C	1	MET	3.1
2	C	221	TRP	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	168	10/11	0.89	0.12	20,36,58,67	0
1	CME	D	168	10/11	0.92	0.10	23,32,55,60	0
1	CME	B	168	10/11	0.93	0.10	19,33,62,70	0
2	CME	C	168	10/11	0.94	0.10	19,32,62,64	0
2	OCS	C	59	9/10	0.97	0.06	16,20,24,26	0

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	A	304	4/4	0.82	0.17	42,42,46,48	0
5	ACT	A	303	4/4	0.86	0.13	35,36,37,39	0
4	6F4	C	302	13/13	0.86	0.15	23,28,34,37	13
4	6F4	D	302	13/13	0.88	0.13	20,24,34,40	13
6	GOL	B	303	6/6	0.88	0.12	28,34,37,48	0
4	6F4	B	302	13/13	0.90	0.13	14,19,34,36	13
3	NAP	C	301	48/48	0.92	0.09	18,26,33,36	48
4	6F4	A	302	13/13	0.92	0.10	21,26,34,43	13
5	ACT	C	303	4/4	0.95	0.08	22,24,29,29	0
3	NAP	D	301	48/48	0.96	0.06	16,20,26,26	0
3	NAP	A	301	48/48	0.97	0.05	17,20,24,28	0
3	NAP	B	301	48/48	0.97	0.05	13,17,20,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.