



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:48 AM UTC

PDB ID : 9IZE / pdb_00009ize
Title : Acarbose hydrolase from human gut microbiota *K. grimontii* TD1, Apg mutant enzyme D336A, complexed with acarviosine-glucose
Authors : Zhou, J.H.; Huang, J.Y.
Deposited on : 2024-08-01
Resolution : 1.87 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	605	
2	C	2	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	605	4863	3101	868	877	17	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLU	GLY	conflict	UNP A0A7H4P971
A	336	ALA	ASP	engineered mutation	UNP A0A7H4P971

- Molecule 2 is an oligosaccharide called (2 {S},3 {R},4 {S},5 {R},6 {R})-5-[[[(1 {S},4 {R},5 {S},6 {S})-3-(hydroxymethyl)-4,5,6-tris(oxidanyl)cyclohex-2-en-1-yl]amino]-6-methyl-oxane-2,3,4-triol-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	33	19	1	13	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	426	Total	O	0	0
			426	426		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltodextrin glucosidase



- Molecule 2: (2 {S},3 {R},4 {S},5 {R},6 {R})-5-(((1 {S},4 {R},5 {S},6 {S})-3-(hydroxymethyl)-4,5,6-tris(oxidanyl)cyclohex-2-en-1-yl]amino]-6-methyl-oxane-2,3,4-triol-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	75.47Å 75.47Å 400.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.70 – 1.87 46.70 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.70-1.87) 99.8 (46.70-1.87)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.87Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.208 , 0.249 0.208 , 0.250	Depositor DCC
R_{free} test set	2896 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtrriage
Anisotropy	0.674	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5322	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EAX, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/5012	0.47	0/6824

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4863	0	4616	26	0
2	C	33	0	11	4	0
3	A	426	0	0	4	0
All	All	5322	0	4627	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:2:A1EAX:C3H	2:C:2:A1EAX:C4H	1.74	1.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:521:ARG:NH1	3:A:703:HOH:O	2.27	0.68
1:A:211:TYR:CZ	2:C:1:GLC:H2	2.28	0.67
1:A:158:ARG:HH12	1:A:222:GLN:NE2	1.94	0.65
1:A:261:ARG:HG3	1:A:270:CYS:SG	2.38	0.64
1:A:16:GLN:HB3	1:A:21:MET:HE1	1.90	0.53
1:A:158:ARG:HH12	1:A:222:GLN:HE22	1.55	0.53
2:C:2:A1EAX:C3H	2:C:2:A1EAX:C5H	2.64	0.53
1:A:260:ASP:HB3	1:A:268:GLY:H	1.75	0.51
1:A:136:SER:O	1:A:139:ARG:HB3	2.10	0.51
1:A:419:GLN:HG3	1:A:537:ALA:O	2.11	0.51
2:C:2:A1EAX:C4H	2:C:2:A1EAX:C2H	2.55	0.50
1:A:524:HIS:CE1	1:A:551:GLN:HB3	2.46	0.50
1:A:358:THR:O	1:A:362:LYS:HG2	2.14	0.47
1:A:140:GLU:CB	1:A:143:GLN:HB2	2.44	0.47
1:A:31:LEU:HB3	3:A:916:HOH:O	2.15	0.46
1:A:473:TRP:CD2	1:A:474:PRO:HD2	2.51	0.46
1:A:255:SER:O	1:A:264:ARG:NH2	2.45	0.45
1:A:256:HIS:CE1	1:A:258:TRP:CD2	3.06	0.43
1:A:524:HIS:HE1	1:A:551:GLN:HB3	1.82	0.43
1:A:143:GLN:HB3	1:A:222:GLN:HA	2.00	0.43
1:A:135:ARG:HG3	1:A:176:GLY:O	2.19	0.42
1:A:160:ALA:HB3	1:A:163:GLU:HG3	2.01	0.42
1:A:72:ASN:ND2	3:A:706:HOH:O	2.34	0.42
1:A:256:HIS:O	1:A:260:ASP:N	2.52	0.42
1:A:568:GLU:HG2	3:A:1049:HOH:O	2.20	0.42
1:A:213:THR:HG21	1:A:216:TYR:CE1	2.55	0.41
1:A:393:TYR:HB3	1:A:398:PHE:CE2	2.56	0.41
1:A:258:TRP:HA	1:A:277:TRP:CE3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	605/605 (100%)	578 (96%)	26 (4%)	1 (0%)	43 34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	492	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	497/504 (99%)	492 (99%)	5 (1%)	68 60

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	VAL
1	A	157	VAL
1	A	349	ASN
1	A	385	ASP
1	A	597	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	98	ASN
1	A	107	GLN
1	A	127	GLN
1	A	290	HIS
1	A	443	GLN
1	A	533	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	12,12,12	1.15	1 (8%)	17,17,17	1.80	3 (17%)
2	A1EAX	C	2	2	21,22,23	6.99	10 (47%)	22,32,34	2.54	11 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	A1EAX	C	2	2	-	1/6/43/46	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	A1EAX	C6H-C5H	18.95	1.59	1.32
2	C	2	A1EAX	C3H-C4H	15.58	1.74	1.53
2	C	2	A1EAX	C2H-C1H	13.71	1.70	1.53
2	C	2	A1EAX	C4H-C5H	-9.30	1.43	1.51
2	C	2	A1EAX	C3H-C2H	-7.94	1.31	1.52
2	C	2	A1EAX	C1H-C6H	-5.79	1.42	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	A1EAX	C1H-N4	-4.04	1.40	1.47
2	C	2	A1EAX	C2-C3	-3.30	1.47	1.52
2	C	2	A1EAX	O5-C5	2.82	1.49	1.43
2	C	1	GLC	O5-C1	2.32	1.48	1.42
2	C	2	A1EAX	C3-C4	-2.25	1.49	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	A1EAX	C4H-C5H-C6H	-5.39	113.38	122.23
2	C	1	GLC	O5-C1-C2	4.61	118.40	110.30
2	C	2	A1EAX	O2H-C2H-C1H	-4.40	100.42	109.08
2	C	2	A1EAX	O7H-C7H-C5H	-3.86	100.60	112.74
2	C	2	A1EAX	O4H-C4H-C3H	3.83	118.49	110.53
2	C	1	GLC	C1-O5-C5	3.41	120.25	113.65
2	C	2	A1EAX	C3-C4-N4	-3.36	101.85	111.49
2	C	2	A1EAX	O2H-C2H-C3H	3.06	117.58	110.38
2	C	2	A1EAX	O2-C2-C3	-2.85	104.24	110.15
2	C	2	A1EAX	O5-C1-C2	-2.80	104.11	110.79
2	C	2	A1EAX	O3-C3-C2	-2.63	104.68	110.05
2	C	1	GLC	O3-C3-C2	-2.40	104.71	110.38
2	C	2	A1EAX	O3H-C3H-C2H	-2.20	105.18	110.38
2	C	2	A1EAX	C6H-C1H-N4	-2.00	107.73	110.68

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	A1EAX	C6H-C1H-N4-C4

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	A1EAX	3	0
2	C	1	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	605/605 (100%)	0.34	42 (6%) 23 25	12, 29, 56, 76	2 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	ALA	4.8
1	A	288	VAL	4.7
1	A	148	TYR	4.4
1	A	287	GLY	4.3
1	A	267	GLY	3.9
1	A	145	ALA	3.8
1	A	140	GLU	3.8
1	A	153	GLY	3.8
1	A	270	CYS	3.7
1	A	143	GLN	3.6
1	A	269	ALA	3.6
1	A	151	ALA	3.5
1	A	169	ALA	3.4
1	A	1	MET	3.2
1	A	266	SER	3.2
1	A	157	VAL	3.1
1	A	283	PHE	3.1
1	A	289	ALA	3.0
1	A	286	GLU	3.0
1	A	268	GLY	3.0
1	A	265	GLY	2.9
1	A	271	HIS	2.9
1	A	273	ALA	2.9
1	A	294	GLY	2.9
1	A	285	PRO	2.8
1	A	147	TYR	2.8
1	A	261	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	259	PHE	2.7
1	A	276	PRO	2.5
1	A	144	ASP	2.4
1	A	272	ASN	2.4
1	A	154	HIS	2.4
1	A	257	ALA	2.4
1	A	166	THR	2.2
1	A	156	ILE	2.2
1	A	264	ARG	2.2
1	A	57	PRO	2.1
1	A	56	ALA	2.1
1	A	222	GLN	2.1
1	A	155	ASP	2.0
1	A	142	GLU	2.0
1	A	146	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	-	-	23,27,30,31	0
2	A1EAX	C	2	21/22	-	-	23,40,48,53	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.