



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 01:29 PM UTC

PDB ID : 1J08 / pdb_00001j08
Title : Crystal structure of glutaredoxin-like protein from *Pyrococcus horikoshii*
Authors : Tanaka, Y.; Tanabe, E.; Tsumoto, K.; Kumagai, I.; Yao, M.; Tanaka, I.
Deposited on : 2002-11-11
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	226	
1	B	226	
1	C	226	
1	D	226	
1	E	226	
1	F	226	
1	G	226	
1	H	226	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glutaredoxin-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	B	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	C	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	D	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	E	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	F	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	G	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			
1	H	225	Total	C	N	O	S	0	0	0
			1793	1151	285	347	10			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	151	Total	O	0	0
			151	151		
2	B	186	Total	O	0	0
			186	186		
2	C	182	Total	O	0	0
			182	182		
2	D	160	Total	O	0	0
			160	160		
2	E	139	Total	O	0	0
			139	139		
2	F	132	Total	O	0	0
			132	132		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	102	Total 102	O 102	0	0
2	H	118	Total 118	O 118	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: glutaredoxin-like protein




- Molecule 1: glutaredoxin-like protein




- Molecule 1: glutaredoxin-like protein



- Molecule 1: glutaredoxin-like protein





- Molecule 1: glutaredoxin-like protein

Chain E: 78% 19%



- Molecule 1: glutaredoxin-like protein

Chain F: 79% 20%



- Molecule 1: glutaredoxin-like protein

Chain G: 69% 30%



- Molecule 1: glutaredoxin-like protein

Chain H: 79% 20%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	144.95Å 257.33Å 48.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.30	Depositor
% Data completeness (in resolution range)	98.5 (10.00-2.30)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	0.08	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.188 , 0.242	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	15514	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.40	0/1827	0.83	2/2462 (0.1%)
1	B	0.45	1/1827 (0.1%)	0.89	3/2462 (0.1%)
1	C	0.40	0/1827	0.84	1/2462 (0.0%)
1	D	0.39	0/1827	0.85	1/2462 (0.0%)
1	E	0.38	0/1827	0.85	4/2462 (0.2%)
1	F	0.39	0/1827	0.85	1/2462 (0.0%)
1	G	0.35	0/1827	0.85	4/2462 (0.2%)
1	H	0.35	0/1827	0.84	2/2462 (0.1%)
All	All	0.39	1/14616 (0.0%)	0.85	18/19696 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	175	MET	SD-CE	-7.50	1.60	1.79

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	171	ILE	N-CA-C	6.70	117.76	107.99
1	H	97	GLY	N-CA-C	-6.07	105.94	112.08
1	B	38	CYS	CA-CB-SG	-6.04	100.50	114.40
1	D	174	ASP	N-CA-C	5.92	119.28	107.62
1	E	174	ASP	N-CA-C	5.76	118.18	107.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1793	0	1789	32	0
1	B	1793	0	1789	31	0
1	C	1793	0	1789	33	0
1	D	1793	0	1789	36	0
1	E	1793	0	1789	37	0
1	F	1793	0	1789	47	0
1	G	1793	0	1789	60	0
1	H	1793	0	1789	29	0
2	A	151	0	0	3	0
2	B	186	0	0	0	0
2	C	182	0	0	7	0
2	D	160	0	0	5	0
2	E	139	0	0	4	1
2	F	132	0	0	0	0
2	G	102	0	0	3	0
2	H	118	0	0	3	0
All	All	15514	0	14312	305	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 305 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:168:LYS:HZ3	1:F:170:LYS:HB2	1.18	1.05
1:F:168:LYS:NZ	1:F:170:LYS:HB2	1.75	0.98
1:G:73:LYS:HZ3	1:G:74:TYR:N	1.62	0.98
1:F:133:ASP:HA	1:F:168:LYS:HE3	1.49	0.94
1:F:168:LYS:HE2	1:F:170:LYS:HD2	1.50	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:341:HOH:O	2:E:341:HOH:O[2_655]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/226 (99%)	216 (97%)	7 (3%)	0	100	100
1	B	223/226 (99%)	219 (98%)	4 (2%)	0	100	100
1	C	223/226 (99%)	213 (96%)	10 (4%)	0	100	100
1	D	223/226 (99%)	213 (96%)	9 (4%)	1 (0%)	30	38
1	E	223/226 (99%)	215 (96%)	7 (3%)	1 (0%)	30	38
1	F	223/226 (99%)	216 (97%)	7 (3%)	0	100	100
1	G	223/226 (99%)	214 (96%)	9 (4%)	0	100	100
1	H	223/226 (99%)	216 (97%)	6 (3%)	1 (0%)	30	38
All	All	1784/1808 (99%)	1722 (96%)	59 (3%)	3 (0%)	43	55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	33	GLU
1	E	33	GLU
1	H	33	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/198 (100%)	191 (97%)	6 (3%)	36	53
1	B	197/198 (100%)	193 (98%)	4 (2%)	48	67

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	197/198 (100%)	192 (98%)	5 (2%)	42	60
1	D	197/198 (100%)	192 (98%)	5 (2%)	42	60
1	E	197/198 (100%)	193 (98%)	4 (2%)	48	67
1	F	197/198 (100%)	195 (99%)	2 (1%)	68	82
1	G	197/198 (100%)	194 (98%)	3 (2%)	57	75
1	H	197/198 (100%)	196 (100%)	1 (0%)	81	90
All	All	1576/1584 (100%)	1546 (98%)	30 (2%)	50	69

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	175	MET
1	G	46	GLN
1	D	127	GLU
1	H	205	LYS
1	F	14	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	189	ASN
1	G	207	GLN
1	H	46	GLN
1	C	201	ASN
1	C	199	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.