



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 12:46 PM UTC

PDB ID : 2J0R / pdb_00002j0r
Title : Structure of the haem-chaperone Proteobacteria-protein HemS
Authors : Schneider, S.; Sharp, K.H.; Barker, P.D.; Paoli, M.
Deposited on : 2006-08-04
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMIN TRANSPORT PROTEIN HEMS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	13	5	0
			2681	1683	483	508	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	162	ASN	ILE	conflict	UNP P31517
A	334	GLN	GLU	conflict	UNP P31517

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 6 is DODECAETHYLENE GLYCOL (CCD ID: 12P) (formula: $C_{24}H_{50}O_{13}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			19	12	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	134	Total	O	0	0
			134	134		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HEMIN TRANSPORT PROTEIN HEMS



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.02Å 68.27Å 73.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.06 – 1.90 50.06 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.06-1.90) 99.9 (50.06-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.189 , 0.233 0.201 , 0.243	Depositor DCC
R_{free} test set	1289 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2901	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEG, EDO, 12P, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.47	50/2742 (1.8%)	1.05	15/3715 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (50) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	255	ILE	C-O	-9.14	1.14	1.24
1	A	147	MET	C-O	-7.80	1.17	1.24
1	A	278	THR	C-O	-7.04	1.16	1.24
1	A	126	ILE	C-O	-6.95	1.16	1.24
1	A	254	GLN	C-O	-6.73	1.15	1.24
1	A	136	LEU	C-O	-6.69	1.14	1.23
1	A	44	ALA	C-O	-6.61	1.15	1.23
1	A	19	LYS	C-O	-6.46	1.16	1.23
1	A	52	ARG	C-O	-6.38	1.16	1.24
1	A	25	ALA	C-O	-6.38	1.16	1.24
1	A	325	GLN	C-O	-6.37	1.15	1.24
1	A	257	THR	C-O	-6.34	1.16	1.24
1	A	304	LEU	C-O	-6.25	1.16	1.24
1	A	301	VAL	C-O	-6.23	1.17	1.24
1	A	293	ARG	C-O	-6.22	1.17	1.24
1	A	111	ALA	CA-CB	-6.04	1.43	1.53
1	A	135	ALA	C-O	-5.86	1.16	1.23
1	A	299	GLY	C-O	-5.83	1.18	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	156[A]	GLN	C-O	-5.81	1.16	1.24
1	A	112	PHE	C-O	-5.78	1.16	1.23
1	A	309	ALA	CA-CB	-5.71	1.44	1.53
1	A	99	LEU	C-O	-5.67	1.17	1.24
1	A	22	ARG	C-O	-5.67	1.17	1.24
1	A	248	GLY	C-O	-5.63	1.15	1.23
1	A	40	VAL	C-O	-5.61	1.17	1.24
1	A	224	VAL	C-O	-5.61	1.17	1.23
1	A	148	PRO	C-O	-5.59	1.17	1.24
1	A	45	LYS	CE-NZ	-5.56	1.32	1.49
1	A	168	GLU	C-O	-5.53	1.17	1.24
1	A	92	LEU	C-O	-5.52	1.17	1.23
1	A	264	THR	C-O	-5.50	1.16	1.24
1	A	271	ASN	C-O	-5.47	1.17	1.23
1	A	109	ALA	C-O	-5.45	1.16	1.23
1	A	247	VAL	C-O	-5.44	1.18	1.24
1	A	114	LEU	C-O	-5.39	1.17	1.23
1	A	86	LEU	C-O	-5.34	1.17	1.23
1	A	139	VAL	C-O	-5.31	1.18	1.24
1	A	256	PHE	C-O	-5.24	1.17	1.23
1	A	42	HIS	C-O	-5.22	1.17	1.24
1	A	309	ALA	C-O	-5.18	1.17	1.24
1	A	137	HIS	C-O	-5.16	1.17	1.23
1	A	51	ALA	C-O	-5.16	1.18	1.24
1	A	279	LEU	C-O	-5.15	1.17	1.24
1	A	125	SER	C-O	-5.09	1.17	1.23
1	A	102	ARG	C-O	-5.08	1.18	1.24
1	A	100	ASP	C-O	-5.07	1.18	1.23
1	A	111	ALA	C-O	-5.06	1.17	1.23
1	A	305	GLU	C-O	-5.03	1.18	1.24
1	A	44	ALA	CA-CB	-5.01	1.45	1.54
1	A	27	LEU	C-O	-5.00	1.18	1.24

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	GLY	N-CA-C	11.41	127.38	112.77
1	A	86	LEU	CA-C-N	8.22	135.86	121.80
1	A	86	LEU	C-N-CA	8.22	135.86	121.80
1	A	248	GLY	N-CA-C	7.49	122.71	111.18
1	A	219	ASP	N-CA-C	-6.63	105.34	113.50
1	A	340	ASN	N-CA-C	6.54	120.72	112.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49	GLY	N-CA-C	6.46	128.49	113.18
1	A	19	LYS	N-CA-C	6.26	120.49	109.96
1	A	87[A]	ASN	CA-C-N	5.62	126.18	120.00
1	A	87[A]	ASN	C-N-CA	5.62	126.18	120.00
1	A	105	LEU	N-CA-C	5.54	119.14	112.38
1	A	206	ASN	N-CA-C	5.48	118.95	111.39
1	A	44	ALA	N-CA-C	5.38	117.59	109.41
1	A	207	LEU	N-CA-C	5.16	118.00	109.85
1	A	170	LEU	N-CA-C	-5.04	103.39	110.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2681	0	2620	37	1
2	A	20	0	30	1	0
3	A	21	0	30	3	0
4	A	10	0	14	2	0
5	A	16	0	22	1	0
6	A	19	0	25	3	0
7	A	134	0	0	1	0
All	All	2901	0	2741	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:PHE:HB2	3:A:1347:PEG:H41	1.65	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:GLU:HG2	1:A:204:ARG:HH22	1.54	0.71
1:A:65:LYS:HZ2	6:A:1352:12P:H241	1.56	0.70
1:A:241:ASN:HD22	1:A:241:ASN:H	1.41	0.69
1:A:65:LYS:HE2	6:A:1352:12P:H351	1.74	0.68
1:A:241:ASN:HD21	1:A:334:GLN:HE22	1.42	0.66
1:A:332:ARG:HH22	3:A:1348:PEG:H21	1.66	0.61
1:A:47:LEU:C	1:A:49:GLY:H	2.08	0.59
1:A:81:TYR:CZ	1:A:99:LEU:HD21	2.41	0.56
1:A:241:ASN:ND2	1:A:334:GLN:HE22	2.04	0.55
1:A:52:ARG:NH2	7:A:2024:HOH:O	2.42	0.53
1:A:118:THR:C	1:A:119:ARG:HG3	2.34	0.53
1:A:316:GLN:C	1:A:317:LEU:HD12	2.35	0.51
1:A:131:HIS:HB3	1:A:168:GLU:HG3	1.93	0.51
1:A:65:LYS:HD2	6:A:1352:12P:H242	1.93	0.49
1:A:170:LEU:HA	1:A:171:SER:C	2.35	0.49
1:A:118:THR:O	1:A:120:HIS:N	2.45	0.49
1:A:148:PRO:HG2	4:A:1349:PGE:H1	1.94	0.48
1:A:136:LEU:HD22	1:A:251:GLY:HA3	1.96	0.48
1:A:317:LEU:HD12	1:A:317:LEU:N	2.29	0.48
1:A:47:LEU:O	1:A:49:GLY:N	2.47	0.47
1:A:241:ASN:H	1:A:241:ASN:ND2	2.10	0.46
3:A:1347:PEG:H22	3:A:1347:PEG:H42	1.57	0.46
1:A:330:GLN:O	1:A:334:GLN:HG3	2.15	0.46
1:A:47:LEU:C	1:A:49:GLY:N	2.73	0.46
1:A:49:GLY:O	1:A:50:ASP:C	2.58	0.45
1:A:96:PRO:O	1:A:97:ARG:HB2	2.17	0.45
1:A:208:THR:O	1:A:209:ARG:C	2.56	0.45
1:A:189:TRP:CE3	1:A:216:VAL:HG11	2.53	0.44
1:A:210:GLN:HB3	2:A:1343:EDO:H11	2.01	0.43
1:A:122:VAL:O	1:A:122:VAL:HG13	2.18	0.42
1:A:19:LYS:NZ	1:A:27:LEU:HD11	2.34	0.41
1:A:318:TYR:CE1	5:A:1351:1PE:H131	2.56	0.41
1:A:200:GLN:HA	1:A:203:LYS:HG2	2.02	0.41
1:A:78:MET:HE1	1:A:142:THR:HG21	2.03	0.41
1:A:118:THR:C	1:A:119:ARG:CG	2.93	0.41
1:A:147:MET:HG2	4:A:1349:PGE:H62	2.03	0.41
1:A:9:TYR:CD1	1:A:35:LEU:HD11	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:GLU:OE2	1:A:211:GLN:NE2[2_444]	1.28	0.92

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/345 (96%)	317 (96%)	11 (3%)	2 (1%)	21 13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	LYS
1	A	49	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	284/292 (97%)	268 (94%)	16 (6%)	19 11

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	71	THR
1	A	77	GLN
1	A	93	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	117	GLU
1	A	119	ARG
1	A	143	GLU
1	A	168	GLU
1	A	171	SER
1	A	188	GLU
1	A	193	THR
1	A	207	LEU
1	A	241	ASN
1	A	267	GLN
1	A	293	ARG
1	A	320	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	85	HIS
1	A	127	GLN
1	A	196	HIS
1	A	200	GLN
1	A	226	ASN
1	A	231	GLN
1	A	237	GLN
1	A	241	ASN
1	A	266	HIS
1	A	320	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	12P	A	1352	-	18,18,36	0.48	0	17,17,35	0.32	0
2	EDO	A	1344	-	3,3,3	0.34	0	2,2,2	0.51	0
2	EDO	A	1346	-	3,3,3	0.46	0	2,2,2	0.36	0
3	PEG	A	1348	-	6,6,6	0.45	0	5,5,5	0.32	0
4	PGE	A	1349	-	9,9,9	0.61	0	8,8,8	0.43	0
3	PEG	A	1347	-	6,6,6	0.52	0	5,5,5	0.48	0
2	EDO	A	1345	-	3,3,3	0.47	0	2,2,2	0.34	0
5	1PE	A	1351	-	15,15,15	0.50	0	14,14,14	0.26	0
2	EDO	A	1342	-	3,3,3	0.42	0	2,2,2	0.41	0
3	PEG	A	1350	-	6,6,6	0.39	0	5,5,5	0.21	0
2	EDO	A	1343	-	3,3,3	0.49	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	12P	A	1352	-	-	9/16/16/34	-
2	EDO	A	1344	-	-	1/1/1/1	-
2	EDO	A	1346	-	-	1/1/1/1	-
3	PEG	A	1348	-	-	2/4/4/4	-
4	PGE	A	1349	-	-	7/7/7/7	-
3	PEG	A	1347	-	-	2/4/4/4	-
2	EDO	A	1345	-	-	1/1/1/1	-
5	1PE	A	1351	-	-	8/13/13/13	-
2	EDO	A	1342	-	-	1/1/1/1	-
3	PEG	A	1350	-	-	1/4/4/4	-
2	EDO	A	1343	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (34) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1347	PEG	C4-C3-O2-C2
3	A	1350	PEG	C4-C3-O2-C2
5	A	1351	1PE	OH4-C13-C23-OH3
6	A	1352	12P	O28-C29-C30-O31
6	A	1352	12P	O25-C26-C27-O28
4	A	1349	PGE	O3-C5-C6-O4
5	A	1351	1PE	OH2-C12-C22-OH3
5	A	1351	1PE	OH6-C15-C25-OH5
4	A	1349	PGE	O1-C1-C2-O2
4	A	1349	PGE	O2-C3-C4-O3
3	A	1348	PEG	O2-C3-C4-O4
5	A	1351	1PE	OH7-C16-C26-OH6
2	A	1345	EDO	O1-C1-C2-O2
6	A	1352	12P	O31-C32-C33-O34
2	A	1342	EDO	O1-C1-C2-O2
2	A	1344	EDO	O1-C1-C2-O2
6	A	1352	12P	C23-C24-O25-C26
4	A	1349	PGE	C3-C4-O3-C5
5	A	1351	1PE	C25-C15-OH6-C26
5	A	1351	1PE	C24-C14-OH5-C25
3	A	1347	PEG	C1-C2-O2-C3
2	A	1346	EDO	O1-C1-C2-O2
4	A	1349	PGE	C6-C5-O3-C4
2	A	1343	EDO	O1-C1-C2-O2
6	A	1352	12P	C33-C32-O31-C30
6	A	1352	12P	C27-C26-O25-C24
3	A	1348	PEG	O1-C1-C2-O2
6	A	1352	12P	C36-C35-O34-C33
5	A	1351	1PE	C13-C23-OH3-C22
5	A	1351	1PE	C23-C13-OH4-C24
6	A	1352	12P	C24-C23-O22-C21
4	A	1349	PGE	C1-C2-O2-C3
4	A	1349	PGE	C4-C3-O2-C2
6	A	1352	12P	O22-C23-C24-O25

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1352	12P	3	0
3	A	1348	PEG	1	0
4	A	1349	PGE	2	0
3	A	1347	PEG	2	0
5	A	1351	1PE	1	0
2	A	1343	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/345 (95%)	0.55	38 (11%) 9 10	10, 19, 44, 60	6 (1%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	GLY	10.1
1	A	118	THR	4.8
1	A	199	PHE	4.4
1	A	180	ALA	4.0
1	A	341	LYS	3.9
1	A	170	LEU	3.8
1	A	181	THR	3.8
1	A	120	HIS	3.7
1	A	171	SER	3.6
1	A	119	ARG	3.5
1	A	193	THR	3.3
1	A	198	PHE	3.2
1	A	187	ALA	3.1
1	A	196	HIS	3.0
1	A	184	ALA	2.9
1	A	194	ASP	2.9
1	A	202	LEU	2.8
1	A	203	LYS	2.8
1	A	297	LYS	2.8
1	A	50	ASP	2.7
1	A	339	ASN	2.7
1	A	195	VAL	2.5
1	A	3	LYS	2.5
1	A	333	ASP	2.4
1	A	14	ALA	2.4
1	A	300	PHE	2.3
1	A	200	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	205	ASN	2.3
1	A	201	LEU	2.3
1	A	169	PRO	2.2
1	A	20	TYR	2.1
1	A	183	GLU	2.1
1	A	87[A]	ASN	2.1
1	A	340	ASN	2.1
1	A	207	LEU	2.1
1	A	215	ALA	2.1
1	A	208	THR	2.0
1	A	191	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGE	A	1349	10/10	0.55	0.21	30,41,44,44	0
3	PEG	A	1348	7/7	0.69	0.20	59,59,60,61	0
2	EDO	A	1342	4/4	0.74	0.17	41,42,43,44	0
3	PEG	A	1347	7/7	0.74	0.17	35,37,40,42	0
3	PEG	A	1350	7/7	0.76	0.16	30,33,39,42	0
6	12P	A	1352	19/37	0.76	0.16	37,45,55,55	0
5	1PE	A	1351	16/16	0.79	0.15	48,49,52,52	0
2	EDO	A	1345	4/4	0.80	0.14	43,43,44,44	0
2	EDO	A	1344	4/4	0.81	0.15	34,37,39,41	0
2	EDO	A	1343	4/4	0.81	0.24	40,40,40,41	0
2	EDO	A	1346	4/4	0.82	0.13	44,44,44,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.