



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 01:13 AM UTC

PDB ID : 6J0L / pdb_00006j0l
Title : Crystal structure of intracellular B30.2 domain of BTN3A3 mutant in complex with sulfate ion
Authors : Yang, Y.Y.; Liu, W.D.; Cai, N.N.; Chen, C.C.; Guo, R.T.; Zhang, Y.H.
Deposited on : 2018-12-24
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Butyrophilin subfamily 3 member A3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1532	984	258	283	7			
1	B	187	Total	C	N	O	S	0	0	0
			1530	982	258	283	7			

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	MET	-	expression tag	UNP O00478
A	266	GLY	-	expression tag	UNP O00478
A	267	SER	-	expression tag	UNP O00478
A	268	SER	-	expression tag	UNP O00478
A	269	HIS	-	expression tag	UNP O00478
A	270	HIS	-	expression tag	UNP O00478
A	271	HIS	-	expression tag	UNP O00478
A	272	HIS	-	expression tag	UNP O00478
A	273	HIS	-	expression tag	UNP O00478
A	274	HIS	-	expression tag	UNP O00478
A	275	SER	-	expression tag	UNP O00478
A	276	SER	-	expression tag	UNP O00478
A	277	GLY	-	expression tag	UNP O00478
A	278	LEU	-	expression tag	UNP O00478
A	279	VAL	-	expression tag	UNP O00478
A	280	PRO	-	expression tag	UNP O00478
A	281	ARG	-	expression tag	UNP O00478
A	282	GLY	-	expression tag	UNP O00478
A	283	SER	-	expression tag	UNP O00478
A	284	HIS	-	expression tag	UNP O00478
A	285	MET	-	expression tag	UNP O00478
A	286	GLU	-	expression tag	UNP O00478
A	287	ASN	-	expression tag	UNP O00478
A	288	LEU	-	expression tag	UNP O00478
A	289	TYR	-	expression tag	UNP O00478

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	290	PHE	-	expression tag	UNP O00478
A	291	GLN	-	expression tag	UNP O00478
A	292	GLY	-	expression tag	UNP O00478
A	293	ALA	-	expression tag	UNP O00478
A	294	GLY	-	expression tag	UNP O00478
A	295	ALA	-	expression tag	UNP O00478
A	296	GLY	-	expression tag	UNP O00478
A	297	ALA	-	expression tag	UNP O00478
A	351	HIS	ARG	engineered mutation	UNP O00478
B	265	MET	-	expression tag	UNP O00478
B	266	GLY	-	expression tag	UNP O00478
B	267	SER	-	expression tag	UNP O00478
B	268	SER	-	expression tag	UNP O00478
B	269	HIS	-	expression tag	UNP O00478
B	270	HIS	-	expression tag	UNP O00478
B	271	HIS	-	expression tag	UNP O00478
B	272	HIS	-	expression tag	UNP O00478
B	273	HIS	-	expression tag	UNP O00478
B	274	HIS	-	expression tag	UNP O00478
B	275	SER	-	expression tag	UNP O00478
B	276	SER	-	expression tag	UNP O00478
B	277	GLY	-	expression tag	UNP O00478
B	278	LEU	-	expression tag	UNP O00478
B	279	VAL	-	expression tag	UNP O00478
B	280	PRO	-	expression tag	UNP O00478
B	281	ARG	-	expression tag	UNP O00478
B	282	GLY	-	expression tag	UNP O00478
B	283	SER	-	expression tag	UNP O00478
B	284	HIS	-	expression tag	UNP O00478
B	285	MET	-	expression tag	UNP O00478
B	286	GLU	-	expression tag	UNP O00478
B	287	ASN	-	expression tag	UNP O00478
B	288	LEU	-	expression tag	UNP O00478
B	289	TYR	-	expression tag	UNP O00478
B	290	PHE	-	expression tag	UNP O00478
B	291	GLN	-	expression tag	UNP O00478
B	292	GLY	-	expression tag	UNP O00478
B	293	ALA	-	expression tag	UNP O00478
B	294	GLY	-	expression tag	UNP O00478
B	295	ALA	-	expression tag	UNP O00478
B	296	GLY	-	expression tag	UNP O00478
B	297	ALA	-	expression tag	UNP O00478

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	351	HIS	ARG	engineered mutation	UNP O00478

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	104	Total	O	0	0
			104	104		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	118	Total 118	O 118	0	0

i

- Molecule 1: Butyrophilin subfamily 3 member A3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.07Å 73.47Å 74.90Å 90.00° 111.39° 90.00°	Depositor
Resolution (Å)	25.00 – 1.95 25.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.4 (25.00-1.95) 97.4 (25.00-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.48 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.176 , 0.234 0.186 , 0.239	Depositor DCC
R_{free} test set	1443 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3324	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/1582	0.79	0/2157
1	B	0.74	0/1579	0.84	1/2152 (0.0%)
All	All	0.72	0/3161	0.81	1/4309 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	475	GLU	N-CA-C	5.01	112.53	108.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1532	0	1470	12	0
1	B	1530	0	1468	10	0
2	A	20	0	0	0	0
2	B	20	0	0	1	0
3	A	104	0	0	1	1
3	B	118	0	0	0	0
All	All	3324	0	2938	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:390:LYS:HA	2:B:503:SO4:O1	1.82	0.79
1:A:344:ASN:HB2	3:A:646:HOH:O	1.99	0.62
1:A:304:MET:O	1:A:305:ALA:CB	2.48	0.61
1:A:404:MET:HE1	1:A:453:ILE:HG21	1.82	0.60
1:A:475:GLU:HB2	1:A:476:PRO:CD	2.34	0.57

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:694:HOH:O	3:A:694:HOH:O[2_557]	1.97	0.23

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/221 (84%)	179 (97%)	5 (3%)	1 (0%)	24	16
1	B	185/221 (84%)	178 (96%)	7 (4%)	0	100	100
All	All	370/442 (84%)	357 (96%)	12 (3%)	1 (0%)	36	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	305	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/191 (87%)	161 (96%)	6 (4%)	31	21
1	B	166/191 (87%)	164 (99%)	2 (1%)	63	61
All	All	333/382 (87%)	325 (98%)	8 (2%)	43	36

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	442	ILE
1	B	300	HIS
1	A	479	LEU
1	A	442	ILE
1	A	484	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	344	ASN
1	B	344	ASN
1	B	421	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	502	-	4,4,4	0.50	0	6,6,6	0.37	0
2	SO4	B	501	-	4,4,4	0.32	0	6,6,6	0.70	0
2	SO4	A	501	-	4,4,4	0.71	0	6,6,6	0.63	0
2	SO4	B	504	-	4,4,4	0.47	0	6,6,6	0.15	0
2	SO4	A	502	-	4,4,4	0.44	0	6,6,6	0.51	0
2	SO4	B	503	-	4,4,4	0.40	0	6,6,6	0.56	0
2	SO4	A	503	-	4,4,4	0.42	0	6,6,6	0.17	0
2	SO4	A	504	-	4,4,4	0.50	0	6,6,6	0.17	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	503	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	187/221 (84%)	0.25	3 (1%) 70 77	22, 33, 64, 100	0
1	B	187/221 (84%)	0.19	4 (2%) 63 70	22, 31, 50, 63	0
All	All	374/442 (84%)	0.22	7 (1%) 66 74	22, 32, 55, 100	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	302	TRP	5.6
1	B	299	TYR	4.1
1	B	484	ILE	3.8
1	B	298	ALA	3.5
1	A	299	TYR	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	504	5/5	0.78	0.08	59,67,80,87	0
2	SO4	B	502	5/5	0.85	0.08	45,57,64,73	0
2	SO4	B	504	5/5	0.86	0.08	56,64,72,80	0
2	SO4	B	503	5/5	0.90	0.07	61,62,72,76	0
2	SO4	A	503	5/5	0.91	0.07	53,60,76,77	0
2	SO4	A	501	5/5	0.96	0.12	28,28,33,38	0
2	SO4	B	501	5/5	0.97	0.08	34,34,40,49	0
2	SO4	A	502	5/5	0.99	0.04	26,28,29,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.