



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:52 AM UTC

PDB ID : 6J0U / pdb_00006j0u
Title : Crystal Structure of the acyltransferase domain from the third module of the ansamitocin polyketide synthase
Authors : Zhang, F.; Zheng, J.
Deposited on : 2018-12-26
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	 2% 83% 6% 11%
1	B	469	 2% 82% 7% 10%
1	C	469	 2% 82% 7% 11%
1	D	469	 2% 83% 6% 10%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 13754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type 1 modular polyketide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	2	0
			3115	1952	591	567	5			
1	B	420	Total	C	N	O	S	0	1	0
			3116	1953	592	566	5			
1	C	419	Total	C	N	O	S	0	0	0
			3103	1944	590	564	5			
1	D	420	Total	C	N	O	S	0	1	0
			3116	1953	592	566	5			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	287	Total	O	0	0
			287	287		
3	B	324	Total	O	0	0
			324	324		
3	C	348	Total	O	0	0
			348	348		
3	D	315	Total	O	0	0
			315	315		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type 1 modular polyketide synthase




- Molecule 1: Type 1 modular polyketide synthase



- Molecule 1: Type 1 modular polyketide synthase



● Molecule 1: Type 1 modular polyketide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	214.73Å 100.30Å 103.41Å 90.00° 113.07° 90.00°	Depositor
Resolution (Å)	50.00 – 1.79 50.00 – 1.79	Depositor EDS
% Data completeness (in resolution range)	98.6 (50.00-1.79) 98.6 (50.00-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, R_{free}	0.187 , 0.219 0.202 , 0.228	Depositor DCC
R_{free} test set	8630 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13754	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 82.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9615e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	0/3176	0.79	0/4338
1	B	0.92	0/3175	0.80	0/4337
1	C	0.92	0/3158	0.78	0/4313
1	D	0.92	1/3175 (0.0%)	0.81	0/4337
All	All	0.92	1/12684 (0.0%)	0.79	0/17325

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	151	LEU	C-O	5.04	1.29	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3115	0	3184	16	0
1	B	3116	0	3185	27	0
1	C	3103	0	3170	20	0
1	D	3116	0	3185	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	10	0	0	0	0
2	D	10	0	0	1	0
3	A	287	0	0	2	0
3	B	324	0	0	7	0
3	C	348	0	0	6	0
3	D	315	0	0	7	0
All	All	13754	0	12724	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:90:ARG:NH2	3:C:601:HOH:O	1.99	0.94
1:B:98[A]:GLN:NE2	3:B:601:HOH:O	2.10	0.85
1:A:205:ARG:HH22	1:A:332:ASN:HD22	1.28	0.81
1:B:205:ARG:HH22	1:B:332:ASN:HD22	1.26	0.81
1:C:205:ARG:HH22	1:C:332:ASN:HD22	1.28	0.80
1:D:98[B]:GLN:NE2	3:D:601:HOH:O	2.13	0.80
1:D:205:ARG:HH22	1:D:332:ASN:HD22	1.27	0.80
1:B:271:ARG:HH11	1:B:271:ARG:CG	1.99	0.76
1:B:208:LEU:HG	1:B:292:LEU:HD13	1.69	0.75
1:C:239:ARG:NH1	1:C:262:GLU:OE1	2.20	0.75
1:B:47:ARG:NH2	1:B:419:VAL:O	2.24	0.71
1:B:216:ARG:NH2	3:B:604:HOH:O	2.25	0.68
1:C:148:VAL:HG22	3:C:810:HOH:O	1.93	0.67
1:A:216:ARG:NH2	3:A:601:HOH:O	2.27	0.67
1:B:98[B]:GLN:NE2	1:B:279:GLY:O	2.29	0.66
1:A:239:ARG:NH1	1:A:262:GLU:OE1	2.30	0.64
1:A:41:ARG:HB3	1:A:42:PRO:HD3	1.81	0.62
1:B:271:ARG:HH11	1:B:271:ARG:HG2	1.65	0.61
1:B:93:LEU:HD22	1:B:355:VAL:HB	1.82	0.60
1:C:216:ARG:HD2	3:C:760:HOH:O	2.01	0.60
1:D:208:LEU:HG	1:D:292:LEU:HD13	1.83	0.60
1:C:41:ARG:HB3	1:C:42:PRO:HD3	1.84	0.59
1:D:93:LEU:HD13	1:D:357:VAL:HG21	1.85	0.57
1:B:257:ASP:OD2	1:B:261:ARG:NH1	2.37	0.56
1:B:369:ASP:OD2	3:B:602:HOH:O	2.17	0.56
1:D:239:ARG:HD2	3:D:793:HOH:O	2.05	0.56
1:A:98[A]:GLN:NE2	1:A:279:GLY:O	2.39	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:LEU:HD22	1:D:355:VAL:HB	1.90	0.54
1:C:257:ASP:OD2	1:C:261:ARG:NH1	2.42	0.53
1:B:271:ARG:HG2	1:B:271:ARG:NH1	2.24	0.52
1:D:264:GLU:OE1	3:D:602:HOH:O	2.19	0.52
1:B:181:GLN:NE2	3:B:608:HOH:O	2.43	0.51
1:C:384:ARG:NH1	3:C:607:HOH:O	2.43	0.51
1:C:148:VAL:HG23	3:C:861:HOH:O	2.11	0.51
1:D:216:ARG:NH2	3:D:604:HOH:O	2.43	0.51
1:B:271:ARG:HH11	1:B:271:ARG:HG3	1.74	0.51
1:B:41:ARG:HB3	1:B:42:PRO:HD3	1.92	0.51
1:B:271:ARG:CG	1:B:271:ARG:NH1	2.66	0.50
1:A:3:VAL:HG13	1:A:4:PRO:HD2	1.93	0.49
1:B:105:VAL:HG21	1:B:359:PRO:HB3	1.94	0.49
1:A:138:VAL:CG1	1:A:154:VAL:HG21	2.42	0.49
1:A:205:ARG:HH22	1:A:332:ASN:ND2	2.05	0.49
2:D:501:PO4:P	3:D:609:HOH:O	2.71	0.48
1:A:93:LEU:HD22	1:A:355:VAL:HB	1.96	0.47
1:A:98[B]:GLN:NE2	3:A:604:HOH:O	2.40	0.47
1:C:93:LEU:HD22	1:C:355:VAL:HB	1.95	0.47
1:C:208:LEU:HG	1:C:292:LEU:HD13	1.96	0.47
1:D:140:VAL:O	1:D:140:VAL:HG22	2.12	0.47
1:C:232:LEU:CD1	1:C:272:MET:HE1	2.43	0.47
1:C:351:HIS:HE1	3:C:601:HOH:O	1.97	0.47
1:D:105:VAL:HG21	1:D:359:PRO:HB3	1.95	0.47
1:D:351:HIS:HE1	3:D:745:HOH:O	1.97	0.47
1:D:41:ARG:HB3	1:D:42:PRO:HD3	1.95	0.47
1:D:138:VAL:HG11	1:D:150:VAL:CG1	2.45	0.46
1:B:181:GLN:HG3	3:B:752:HOH:O	2.14	0.46
1:A:93:LEU:HG	1:A:173:PRO:HB3	1.98	0.46
1:C:105:VAL:O	1:C:106:ALA:HB3	2.16	0.46
1:D:181:GLN:NE2	3:D:611:HOH:O	2.47	0.46
1:A:308:PRO:HB2	1:A:310:ILE:O	2.15	0.46
1:B:239:ARG:NH1	1:B:262:GLU:OE1	2.43	0.46
1:D:327:GLU:HG3	1:D:331:ARG:NH2	2.31	0.46
1:A:105:VAL:O	1:A:106:ALA:HB3	2.16	0.46
1:B:205:ARG:HH22	1:B:332:ASN:ND2	2.06	0.45
1:C:3:VAL:HG22	1:C:4:PRO:HD2	1.99	0.45
1:B:239:ARG:HD3	3:B:670:HOH:O	2.17	0.45
1:B:184:VAL:HG11	1:B:202:VAL:HG23	1.99	0.44
1:C:140:VAL:O	1:C:142:PRO:HD3	2.18	0.43
1:C:3:VAL:CG2	1:C:4:PRO:HD2	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:3:VAL:HG23	1:D:4:PRO:HD2	2.01	0.42
1:C:371:ALA:HB1	1:C:376:THR:HG21	2.01	0.42
1:D:98[A]:GLN:NE2	1:D:279:GLY:O	2.52	0.42
1:B:3:VAL:HG13	1:B:4:PRO:HD2	2.02	0.42
1:A:105:VAL:HG21	1:A:359:PRO:HB3	2.01	0.42
1:A:138:VAL:HG12	1:A:154:VAL:HG21	2.02	0.41
1:D:300:LEU:HB3	1:D:303:ILE:HD11	2.01	0.41
1:B:93:LEU:HD13	1:B:357:VAL:HG21	2.02	0.41
1:C:93:LEU:HD13	1:C:357:VAL:HG21	2.03	0.41
1:C:149:ASP:HB3	1:C:207:ALA:HA	2.03	0.41
1:D:274:ARG:HD3	1:D:274:ARG:N	2.36	0.40
1:A:152:GLN:HB2	1:A:153:PRO:HD3	2.03	0.40
1:B:176:VAL:HG21	1:B:190:ALA:HB2	2.02	0.40
1:B:49:GLN:NE2	3:B:620:HOH:O	2.54	0.40
1:B:306:ARG:NH2	1:B:309:GLU:OE2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/469 (89%)	410 (98%)	9 (2%)	0	100	100
1	B	419/469 (89%)	410 (98%)	9 (2%)	0	100	100
1	C	417/469 (89%)	406 (97%)	11 (3%)	0	100	100
1	D	419/469 (89%)	409 (98%)	10 (2%)	0	100	100
All	All	1674/1876 (89%)	1635 (98%)	39 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/352 (89%)	305 (97%)	8 (3%)	40	28
1	B	313/352 (89%)	302 (96%)	11 (4%)	32	19
1	C	311/352 (88%)	306 (98%)	5 (2%)	55	47
1	D	313/352 (89%)	310 (99%)	3 (1%)	68	64
All	All	1250/1408 (89%)	1223 (98%)	27 (2%)	47	34

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	VAL
1	A	93	LEU
1	A	122	ARG
1	A	138	VAL
1	A	309	GLU
1	A	375	ARG
1	A	406	VAL
1	A	419	VAL
1	B	61	ARG
1	B	93	LEU
1	B	98[A]	GLN
1	B	98[B]	GLN
1	B	122	ARG
1	B	126	VAL
1	B	271	ARG
1	B	292	LEU
1	B	306	ARG
1	B	406	VAL
1	B	410	ARG
1	C	30	GLU
1	C	292	LEU
1	C	306	ARG
1	C	309	GLU
1	C	419	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	93	LEU
1	D	122	ARG
1	D	292	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	GLN
1	A	332	ASN
1	A	351	HIS
1	B	27	HIS
1	B	49	GLN
1	B	152	GLN
1	B	181	GLN
1	B	332	ASN
1	B	351	HIS
1	C	181	GLN
1	C	332	ASN
1	C	351	HIS
1	D	181	GLN
1	D	332	ASN
1	D	351	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	D	502	-	4,4,4	1.03	0	6,6,6	0.53	0
2	PO4	C	501	-	4,4,4	1.48	1 (25%)	6,6,6	0.22	0
2	PO4	D	501	-	4,4,4	1.00	0	6,6,6	0.72	0
2	PO4	A	501	-	4,4,4	0.94	0	6,6,6	0.53	0
2	PO4	C	502	-	4,4,4	0.62	0	6,6,6	0.75	0
2	PO4	B	501	-	4,4,4	0.85	0	6,6,6	0.80	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	PO4	P-O1	2.74	1.57	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/469 (89%)	0.12	8 (1%) 66 66	13, 20, 41, 64	2 (0%)
1	B	420/469 (89%)	0.03	9 (2%) 63 64	12, 21, 40, 62	1 (0%)
1	C	419/469 (89%)	0.08	8 (1%) 66 66	13, 21, 39, 70	0
1	D	420/469 (89%)	0.08	9 (2%) 63 64	13, 21, 37, 63	1 (0%)
All	All	1678/1876 (89%)	0.08	34 (2%) 65 65	12, 21, 39, 70	4 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	VAL	7.3
1	A	1	VAL	6.1
1	C	419	VAL	6.0
1	B	420	PRO	4.8
1	C	1	VAL	4.6
1	B	1	VAL	4.5
1	C	323	ARG	4.3
1	D	1	VAL	3.8
1	B	132	GLY	3.5
1	D	323	ARG	3.5
1	A	132	GLY	3.3
1	D	420	PRO	3.3
1	C	132	GLY	3.3
1	D	129	ASP	2.9
1	C	131	ALA	2.8
1	D	419	VAL	2.6
1	B	129	ASP	2.6
1	B	419	VAL	2.6
1	D	279	GLY	2.6
1	A	2	PRO	2.5
1	B	75	GLU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	279	GLY	2.4
1	A	271	ARG	2.4
1	D	132	GLY	2.3
1	B	271	ARG	2.2
1	A	261	ARG	2.1
1	C	31	ARG	2.1
1	A	323	ARG	2.1
1	C	47	ARG	2.1
1	D	138	VAL	2.1
1	B	323	ARG	2.1
1	C	2	PRO	2.1
1	B	306	ARG	2.0
1	D	140	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	501	5/5	0.75	0.15	66,72,77,98	0
2	PO4	B	501	5/5	0.79	0.12	44,50,65,67	0
2	PO4	D	502	5/5	0.81	0.14	34,57,65,75	0
2	PO4	C	502	5/5	0.82	0.13	31,53,66,72	0
2	PO4	C	501	5/5	0.82	0.13	31,43,53,71	0
2	PO4	D	501	5/5	0.86	0.12	47,51,67,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.