



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:13 AM UTC

PDB ID : 2J1G / pdb_00002j1g
Title : L-ficolin complexed to N-acetyl-cystein
Authors : Garlatti, V.; Gaboriaud, C.
Deposited on : 2006-08-11
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	13% 67% 22% 10%
1	B	218	89% 9%
1	C	218	81% 16%
1	D	218	% 82% 16%
1	E	218	85% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	218	
2	G	5	
3	H	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	P4C	D	302	-	X	-	-
5	ACT	C	302	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FICOLIN-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	3	0
			1617	1025	280	305	7			
1	B	217	Total	C	N	O	S	0	0	0
			1736	1092	305	330	9			
1	C	216	Total	C	N	O	S	0	0	0
			1729	1087	304	329	9			
1	D	214	Total	C	N	O	S	0	0	0
			1715	1078	302	327	8			
1	E	218	Total	C	N	O	S	0	1	0
			1755	1102	311	333	9			
1	F	217	Total	C	N	O	S	0	0	0
			1736	1092	305	330	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	THR	VAL	conflict	UNP Q15485
A	247	THR	VAL	conflict	UNP Q15485
B	168	THR	VAL	conflict	UNP Q15485
B	247	THR	VAL	conflict	UNP Q15485
C	168	THR	VAL	conflict	UNP Q15485
C	247	THR	VAL	conflict	UNP Q15485
D	168	THR	VAL	conflict	UNP Q15485
D	247	THR	VAL	conflict	UNP Q15485
E	168	THR	VAL	conflict	UNP Q15485
E	247	THR	VAL	conflict	UNP Q15485
F	168	THR	VAL	conflict	UNP Q15485
F	247	THR	VAL	conflict	UNP Q15485

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	5	Total	C	N	O	0	0	0
			60	34	2	24			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is O-ACETALDEHYDYL-HEXAETHYLENE GLYCOL (CCD ID: P4C) (formula: C₁₄H₂₈O₈).



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	5	0
			5	3	2		
4	E	1	Total	C	O	0	0
			17	11	6		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	C	1	Total	C	O	0	0
			4	2	2		
5	E	1	Total	C	O	0	0
			4	2	2		
5	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		
6	C	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Ca	0	0
			1	1		
6	E	1	Total	Ca	0	0
			1	1		
6	F	1	Total	Ca	0	0
			1	1		

- Molecule 7 is N-ACETYL-L-CYSTEINE (CCD ID: SC2) (formula: $C_5H_9NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	F	1	Total	C	N	O	S	2	0
			10	5	1	3	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	77	Total	O	0	0
			77	77		
8	B	206	Total	O	0	0
			206	206		
8	C	168	Total	O	0	0
			168	168		
8	D	91	Total	O	0	0
			91	91		
8	E	185	Total	O	0	0
			185	185		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	F	107	Total 107	O 107	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FICOLIN-2




• Molecule 1: FICOLIN-2



• Molecule 1: FICOLIN-2



• Molecule 1: FICOLIN-2





• Molecule 1: FICOLIN-2

Chain E: 85% 13% .



• Molecule 1: FICOLIN-2

Chain F: 11% 78% 18% .



• Molecule 2: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 20% 80%



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	99.04Å 99.04Å 142.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 1.95 15.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-1.95) 99.9 (15.00-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.87 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.176 , 0.210 0.178 , 0.212	Depositor DCC
R_{free} test set	5667 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l 0.049 for h,-h-k,-l 0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11289	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CA, MAN, ACT, NAG, P4C, SC2, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.39	0/1660	0.67	0/2242
1	B	0.56	0/1784	0.77	0/2413
1	C	0.53	0/1776	0.78	2/2402 (0.1%)
1	D	0.41	0/1762	0.74	0/2383
1	E	0.53	0/1803	0.78	2/2439 (0.1%)
1	F	1.10	4/1784 (0.2%)	1.13	9/2413 (0.4%)
All	All	0.64	4/10569 (0.0%)	0.83	13/14292 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	1	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	72	PRO	N-CA	-30.92	1.00	1.47
1	F	73	CYS	CA-C	-25.17	1.19	1.52
1	F	72	PRO	CA-CB	11.14	1.75	1.53
1	F	73	CYS	CA-CB	-8.65	1.38	1.53

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	73	CYS	CB-CA-C	21.22	152.65	110.42
1	F	72	PRO	CB-CA-C	-20.78	70.62	110.10
1	F	72	PRO	N-CA-C	18.25	157.72	112.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	73	CYS	N-CA-CB	-15.52	84.27	110.49
1	F	72	PRO	N-CA-CB	9.32	113.25	103.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	F	73	CYS	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	73	CYS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1617	0	1493	33	0
1	B	1736	0	1608	21	0
1	C	1729	0	1603	25	0
1	D	1715	0	1585	25	0
1	E	1755	0	1623	24	0
1	F	1736	0	1609	39	0
2	G	60	0	52	1	0
3	H	28	0	25	0	0
4	A	18	0	22	0	0
4	B	4	0	4	0	0
4	D	5	0	4	0	0
4	E	17	0	20	1	0
5	B	4	0	3	0	0
5	C	8	0	6	4	0
5	E	4	0	3	0	0
5	F	4	0	3	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	E	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	F	1	0	0	0	0
7	F	10	0	8	1	0
8	A	77	0	0	4	0
8	B	206	0	0	4	0
8	C	168	0	0	7	0
8	D	91	0	0	4	0
8	E	185	0	0	3	0
8	F	107	0	0	10	0
All	All	11289	0	9671	163	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 163 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:GLN:HE21	1:B:243:LYS:CE	1.53	1.22
1:B:216:GLN:NE2	1:B:243:LYS:HE3	1.57	1.19
1:B:216:GLN:HE21	1:B:243:LYS:HE3	1.03	1.17
1:F:188:ALA:HB3	8:F:404:HOH:O	1.58	1.03
1:A:227:LEU:O	1:A:228:ASN:HB2	1.59	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	192/218 (88%)	174 (91%)	13 (7%)	5 (3%)	4 1
1	B	215/218 (99%)	203 (94%)	11 (5%)	1 (0%)	24 16
1	C	214/218 (98%)	200 (94%)	13 (6%)	1 (0%)	24 16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	212/218 (97%)	201 (95%)	10 (5%)	1 (0%)	24	16
1	E	217/218 (100%)	206 (95%)	9 (4%)	2 (1%)	14	6
1	F	215/218 (99%)	201 (94%)	12 (6%)	2 (1%)	14	6
All	All	1265/1308 (97%)	1185 (94%)	68 (5%)	12 (1%)	12	6

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	ASN
1	F	73	CYS
1	F	160	ALA
1	A	229	THR
1	A	198	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/183 (92%)	159 (95%)	9 (5%)	20	9
1	B	182/183 (100%)	177 (97%)	5 (3%)	39	32
1	C	181/183 (99%)	170 (94%)	11 (6%)	17	7
1	D	179/183 (98%)	172 (96%)	7 (4%)	28	18
1	E	184/183 (100%)	176 (96%)	8 (4%)	26	15
1	F	182/183 (100%)	170 (93%)	12 (7%)	15	5
All	All	1076/1098 (98%)	1024 (95%)	52 (5%)	22	12

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	104	LEU
1	E	150	LEU
1	F	202	VAL
1	D	150	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	90	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	139	GLN
1	F	94	HIS
1	E	156	HIS
1	E	246	HIS
1	F	195	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	G	1	1,2	14,14,15	0.70	0	17,19,21	1.17	2 (11%)
2	NAG	G	2	2	14,14,15	0.57	0	17,19,21	0.71	0
2	BMA	G	3	2	11,11,12	0.80	0	15,15,17	1.02	1 (6%)
2	MAN	G	4	2	11,11,12	0.60	0	15,15,17	0.91	1 (6%)
2	FUC	G	5	2	10,10,11	0.66	0	14,14,16	1.10	0
3	NAG	H	1	1,3	14,14,15	0.52	0	17,19,21	0.82	0
3	NAG	H	2	3	14,14,15	0.56	0	17,19,21	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	BMA	G	3	2	-	0/2/19/22	0/1/1/1
2	MAN	G	4	2	-	0/2/19/22	0/1/1/1
2	FUC	G	5	2	-	-	0/1/1/1
3	NAG	H	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	3	BMA	C1-C2-C3	2.36	113.08	109.64
2	G	1	NAG	C3-C4-C5	-2.34	105.99	110.23
2	G	1	NAG	O5-C1-C2	-2.19	107.90	111.29
2	G	4	MAN	C2-C3-C4	-2.03	107.28	110.86

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	5	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	P4C	A	301	-	17,17,21	4.97	2 (11%)	16,16,20	4.22	3 (18%)
5	ACT	B	301	-	3,3,3	0.80	0	3,3,3	1.38	0
4	P4C	E	303	-	16,16,21	1.05	1 (6%)	15,15,20	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	C	302	-	3,3,3	0.77	0	3,3,3	1.11	0
5	ACT	F	301	-	3,3,3	0.76	0	3,3,3	1.36	0
5	ACT	C	301	-	3,3,3	0.82	0	3,3,3	1.29	0
5	ACT	E	301	-	3,3,3	0.80	0	3,3,3	1.47	0
4	P4C	D	302	-	4,4,21	1.50	1 (25%)	3,3,20	1.89	1 (33%)
7	SC2	F	302	-	9,9,9	1.61	1 (11%)	11,11,11	3.32	3 (27%)
4	P4C	B	303	-	3,3,21	2.21	1 (33%)	2,2,20	1.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	P4C	A	301	-	-	3/15/15/19	-
4	P4C	E	303	-	-	7/14/14/19	-
7	SC2	F	302	-	-	4/10/10/10	-
4	P4C	D	302	-	-	2/2/2/19	-
4	P4C	B	303	-	-	0/1/1/19	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	P4C	C15-C14	-20.10	0.48	1.49
7	F	302	SC2	OT-CT	-4.28	1.13	1.23
4	E	303	P4C	O1-C2	-3.71	1.23	1.42
4	B	303	P4C	O1-C2	-3.61	1.23	1.42
4	A	301	P4C	O16-C17	3.45	1.57	1.42

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	P4C	O13-C14-C15	12.91	169.21	110.35
4	A	301	P4C	O16-C15-C14	10.35	157.51	110.35
7	F	302	SC2	OT-CT-CM	-9.49	105.16	122.05
7	F	302	SC2	OT-CT-N	4.30	129.58	121.98
4	D	302	P4C	O7-C6-C5	3.27	131.09	111.82

There are no chirality outliers.

5 of 16 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	F	302	SC2	OT-CT-N-CA
7	F	302	SC2	CM-CT-N-CA
4	E	303	P4C	C15-C14-O13-C12
4	E	303	P4C	O13-C14-C15-O16
4	D	302	P4C	O4-C5-C6-O7


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	303	P4C	1	0
5	C	302	ACT	4	0
7	F	302	SC2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/218 (90%)	0.55	29 (14%) 6 6	24, 51, 84, 95	4 (2%)
1	B	217/218 (99%)	-1.15	0 100 100	24, 28, 38, 49	0
1	C	216/218 (99%)	-0.94	1 (0%) 87 90	21, 28, 39, 60	2 (0%)
1	D	214/218 (98%)	-0.40	3 (1%) 73 79	37, 43, 55, 63	0
1	E	218/218 (100%)	-1.00	0 100 100	16, 32, 40, 53	2 (0%)
1	F	217/218 (99%)	0.46	25 (11%) 9 11	22, 36, 47, 53	5 (2%)
All	All	1279/1308 (97%)	-0.43	58 (4%) 38 44	16, 35, 58, 95	13 (1%)

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	257	GLY	4.8
1	F	268	TRP	4.2
1	F	206	ALA	3.5
1	A	247	THR	3.1
1	F	179	PHE	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	H	2	14/15	0.97	0.05	47,48,49,49	0
2	MAN	G	4	11/12	0.98	0.05	46,48,49,50	0
2	FUC	G	5	10/11	0.98	0.04	38,41,42,44	0
3	NAG	H	1	14/15	0.98	0.04	42,46,47,48	0
2	BMA	G	3	11/12	0.98	0.04	44,47,48,48	0
2	NAG	G	2	14/15	0.99	0.03	33,34,39,41	0
2	NAG	G	1	14/15	0.99	0.04	27,28,33,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	P4C	A	301	18/22	0.98	0.05	20,45,48,48	5
4	P4C	B	303	4/22	-	-	72,74,74,74	4
4	P4C	D	302	5/22	-	-	20,20,20,20	5
4	P4C	E	303	17/22	0.98	0.05	54,58,66,67	0
5	ACT	B	301	4/4	0.98	0.08	38,39,39,40	0
5	ACT	F	301	4/4	0.98	0.04	44,44,45,45	0
7	SC2	F	302	10/10	0.98	0.06	48,54,54,54	5
5	ACT	E	301	4/4	0.99	0.04	41,41,41,41	0
5	ACT	C	301	4/4	0.99	0.04	30,30,31,32	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	CA	D	301	1/1	0.99	0.05	53,53,53,53	0
6	CA	F	303	1/1	0.99	0.04	52,52,52,52	0
5	ACT	C	302	4/4	0.99	0.03	38,39,39,39	4
6	CA	E	302	1/1	1.00	0.05	25,25,25,25	0
6	CA	C	303	1/1	1.00	0.05	29,29,29,29	0
6	CA	B	302	1/1	1.00	0.04	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around P4C A 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around P4C B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around P4C D 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P4C E 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.