



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:35 PM UTC

PDB ID : 4J14 / pdb_00004j14
Title : Crystal Structure of Human Cytochrome P450 CYP46A1 with Posaconazole Bound
Authors : Stout, C.D.; Mast, N.; Pikuleva, I.A.
Deposited on : 2013-01-31
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cholesterol 24-hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total 3507	C 2241	N 615	O 634	S 17	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	49	MET	-	expression tag	UNP Q9Y6A2
A	50	ALA	-	expression tag	UNP Q9Y6A2
A	501	HIS	-	expression tag	UNP Q9Y6A2
A	502	HIS	-	expression tag	UNP Q9Y6A2
A	503	HIS	-	expression tag	UNP Q9Y6A2
A	504	HIS	-	expression tag	UNP Q9Y6A2

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	
			43	34	1	4	4	
							0	0

- Molecule 3 is POSACONAZOLE (CCD ID: X2N) (formula: $C_{37}H_{42}F_2N_8O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	
			51	37	2	8	4	
							0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	82	Total	O	0	0
			82	82		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cholesterol 24-hydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	121.64Å 121.64Å 143.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.78 – 2.50 92.76 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (92.78-2.50) 99.8 (92.76-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.08 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.213 , 0.263 0.186 , 0.231	Depositor DCC
R_{free} test set	1003 reflections (5.29%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3689	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: X2N, HEM, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	1/3579 (0.0%)	0.99	7/4828 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	385	VAL	CA-CB	5.53	1.58	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	481	PRO	N-CA-C	-8.09	95.81	112.47
1	A	236	LYS	N-CA-C	5.80	117.81	110.33
1	A	80	PHE	N-CA-C	5.68	117.30	109.29
1	A	194	LYS	N-CA-C	-5.48	104.95	111.03
1	A	167	ILE	CB-CA-C	-5.29	105.12	112.04
1	A	228	THR	N-CA-C	5.27	117.34	109.59
1	A	388	ASN	N-CA-C	5.07	118.59	111.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	227	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3507	0	3558	57	0
2	A	43	0	30	4	0
3	A	51	0	42	7	0
4	A	6	0	8	2	0
5	A	82	0	0	4	0
All	All	3689	0	3638	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:THR:HA	1:A:229:LEU:CB	1.82	1.07
1:A:228:THR:HA	1:A:229:LEU:HB2	1.52	0.88
1:A:351:GLN:HE21	1:A:355:GLN:HE21	1.22	0.84
1:A:273:ALA:HB3	4:A:603:GOL:H12	1.61	0.83
1:A:228:THR:HA	1:A:229:LEU:HB3	1.63	0.79
1:A:228:THR:CG2	1:A:231:LYS:HB2	2.13	0.79
1:A:251:ARG:NH1	1:A:294:ASP:OD1	2.17	0.77
2:A:601:HEM:NA	3:A:602:X2N:HAS	1.98	0.76
1:A:480:ASP:C	1:A:481:PRO:O	2.24	0.74
2:A:601:HEM:C1A	3:A:602:X2N:HAS	2.24	0.72
1:A:306:THR:HG22	1:A:475:THR:HG21	1.72	0.71
1:A:228:THR:HG22	1:A:231:LYS:HB2	1.72	0.71
1:A:211:LEU:HD21	1:A:300:PHE:HZ	1.55	0.71
1:A:108:MET:HA	1:A:108:MET:HE2	1.76	0.68
1:A:362:ARG:NH1	1:A:408:PRO:O	2.27	0.67
1:A:273:ALA:HB3	4:A:603:GOL:C1	2.27	0.65
1:A:339:LYS:HE2	1:A:341:TYR:O	1.96	0.65
1:A:228:THR:CA	1:A:229:LEU:HB2	2.24	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ILE:HG22	3:A:602:X2N:HAG	1.80	0.64
1:A:306:THR:HG21	3:A:602:X2N:NBE	2.14	0.62
1:A:351:GLN:NE2	1:A:355:GLN:HE21	1.99	0.57
1:A:324:ILE:HG12	5:A:724:HOH:O	2.05	0.57
1:A:310:HIS:CE1	1:A:361:LEU:HG	2.40	0.56
1:A:209:LYS:N	1:A:210:PRO:HD2	2.20	0.56
1:A:274:ASP:HB3	1:A:277:THR:H	1.69	0.55
1:A:163:GLN:O	1:A:167:ILE:HG12	2.08	0.54
1:A:80:PHE:HE1	3:A:602:X2N:HAZ2	1.73	0.53
1:A:122:GLY:HA3	1:A:294:ASP:CG	2.33	0.53
1:A:370:THR:OG1	2:A:601:HEM:O2A	2.24	0.53
1:A:332:VAL:HG22	1:A:336:ILE:HD11	1.90	0.53
1:A:235:GLY:HA2	5:A:744:HOH:O	2.08	0.52
1:A:296:PHE:CD1	1:A:296:PHE:C	2.89	0.51
1:A:380:ILE:HD12	1:A:385:VAL:HG21	1.92	0.51
1:A:189:MET:HG2	1:A:203:MET:SD	2.50	0.51
1:A:276:LEU:HA	1:A:279:ILE:HG12	1.93	0.50
1:A:491:TRP:C	1:A:491:TRP:CD1	2.90	0.50
1:A:281:LYS:O	1:A:284:GLU:HG2	2.12	0.50
1:A:126:VAL:HG11	3:A:602:X2N:HBJ	1.94	0.50
1:A:123:GLN:HB2	5:A:746:HOH:O	2.13	0.48
1:A:233:LEU:N	1:A:234:PRO:CD	2.78	0.47
1:A:456:GLN:HB3	1:A:457:ARG:HG2	1.97	0.47
1:A:480:ASP:O	1:A:481:PRO:O	2.32	0.47
1:A:201:THR:O	1:A:208:GLN:HG3	2.14	0.46
1:A:231:LYS:C	1:A:234:PRO:HD2	2.40	0.46
1:A:283:GLU:HG3	1:A:292:LEU:HB2	1.98	0.45
1:A:313:PHE:CD2	1:A:364:TYR:HB3	2.52	0.45
2:A:601:HEM:C4A	3:A:602:X2N:HAS	2.52	0.45
1:A:489:ARG:HG2	5:A:764:HOH:O	2.17	0.45
1:A:412:ASN:O	1:A:415:ARG:HG3	2.17	0.44
1:A:466:GLN:CD	1:A:483:LEU:HB2	2.43	0.43
1:A:240:LEU:O	1:A:244:ARG:HG3	2.19	0.43
1:A:165:VAL:HG22	1:A:454:LEU:HD23	2.01	0.43
1:A:227:ASN:C	1:A:229:LEU:HB2	2.43	0.42
1:A:473:GLN:OE1	1:A:477:LYS:HE3	2.19	0.42
1:A:374:LEU:HD11	1:A:378:THR:OG1	2.19	0.42
1:A:402:ASP:HA	1:A:405:PHE:O	2.20	0.42
1:A:218:MET:HE1	1:A:301:ILE:HG12	2.02	0.42
1:A:359:GLU:CD	1:A:362:ARG:HH21	2.29	0.41
1:A:369:GLY:HA3	1:A:393:PHE:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:PHE:CD1	1:A:442:PHE:C	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	434/456 (95%)	413 (95%)	18 (4%)	3 (1%)	18 34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	LEU
1	A	234	PRO
1	A	481	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	379/397 (96%)	364 (96%)	15 (4%)	28 54

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	VAL
1	A	112	LEU
1	A	136	LYS
1	A	159	GLU
1	A	181	GLN
1	A	226	ARG
1	A	229	LEU
1	A	232	PHE
1	A	237	ARG
1	A	239	GLN
1	A	245	GLU
1	A	287	GLN
1	A	292	LEU
1	A	332	VAL
1	A	339	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN
1	A	123	GLN
1	A	227	ASN
1	A	351	GLN
1	A	355	GLN
1	A	440	GLN
1	A	444	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	X2N	A	602	2	55,57,57	1.58	7 (12%)	68,82,82	2.17	24 (35%)
4	GOL	A	603	-	5,5,5	0.39	0	5,5,5	0.36	0
2	HEM	A	601	3,1	50,50,50	3.32	30 (60%)	67,82,82	1.98	17 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	X2N	A	602	2	-	9/38/59/59	0/7/7/7
4	GOL	A	603	-	-	4/4/4/4	-
2	HEM	A	601	3,1	-	6/14/54/54	-

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	FE-NB	8.12	2.19	1.94
3	A	602	X2N	OAC-CBP	6.31	1.34	1.22
2	A	601	HEM	C3C-C4C	-6.15	1.35	1.46
2	A	601	HEM	C3C-C2C	6.09	1.49	1.37
2	A	601	HEM	C3B-C2B	5.96	1.49	1.37
2	A	601	HEM	CHB-C1B	5.74	1.50	1.38
2	A	601	HEM	C1C-C2C	-5.71	1.34	1.45
2	A	601	HEM	CHD-C4C	5.65	1.49	1.38
2	A	601	HEM	CAB-C3B	-5.43	1.32	1.47
3	A	602	X2N	CBN-NBW	-5.41	1.34	1.44
2	A	601	HEM	CHC-C1C	4.97	1.48	1.38
2	A	601	HEM	C1B-C2B	-4.61	1.35	1.44
2	A	601	HEM	CBB-CAB	4.50	1.52	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	HEM	CHA-C1A	4.49	1.49	1.39
2	A	601	HEM	C4D-ND	-3.75	1.33	1.40
2	A	601	HEM	C1B-NB	-3.74	1.33	1.40
2	A	601	HEM	C1A-C2A	3.63	1.51	1.44
2	A	601	HEM	C4A-C3A	3.36	1.51	1.43
2	A	601	HEM	CHD-C1D	-3.27	1.31	1.39
3	A	602	X2N	CAT-NBF	3.23	1.35	1.30
3	A	602	X2N	CAA-CAU	-3.17	1.32	1.51
2	A	601	HEM	FE-ND	3.08	2.04	1.94
2	A	601	HEM	CHA-C4D	-3.00	1.32	1.38
2	A	601	HEM	FE-NC	2.89	2.04	1.95
3	A	602	X2N	CBP-NBW	-2.75	1.35	1.39
2	A	601	HEM	C4A-NA	-2.53	1.34	1.39
2	A	601	HEM	C4D-C3D	2.52	1.49	1.45
2	A	601	HEM	C1D-ND	-2.31	1.34	1.38
3	A	602	X2N	OBG-CBK	-2.31	1.32	1.37
2	A	601	HEM	FE-NA	2.28	2.02	1.95
2	A	601	HEM	C1D-C2D	2.18	1.48	1.44
3	A	602	X2N	CAQ-NBE	2.13	1.35	1.32
2	A	601	HEM	C2A-C3A	-2.09	1.33	1.38
2	A	601	HEM	C4C-NC	-2.09	1.35	1.39
2	A	601	HEM	CHC-C4B	-2.08	1.34	1.39
2	A	601	HEM	C1C-NC	-2.04	1.35	1.39
2	A	601	HEM	C3D-C2D	-2.00	1.32	1.36

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	X2N	CBP-NBX-NBF	-7.01	107.36	112.87
3	A	602	X2N	CAT-NBF-NBX	6.08	109.69	103.68
2	A	601	HEM	C4B-C3B-C2B	-6.01	101.76	107.28
2	A	601	HEM	C1B-NB-C4B	4.52	110.56	105.21
2	A	601	HEM	CAA-C2A-C3A	-4.39	117.22	127.07
3	A	602	X2N	CAY-NBU-CBM	-4.14	106.82	118.11
3	A	602	X2N	NBW-CBP-NBX	4.00	107.39	102.88
3	A	602	X2N	CBC-NBV-CAS	3.64	135.96	128.82
2	A	601	HEM	CHC-C4B-NB	3.58	128.28	124.42
2	A	601	HEM	CHC-C1C-NC	-3.54	120.60	124.45
2	A	601	HEM	CHD-C4C-NC	3.46	128.22	124.45
3	A	602	X2N	CBC-NBV-NBE	-3.42	116.28	121.25
3	A	602	X2N	CBS-NBX-CBP	3.29	134.03	125.26
2	A	601	HEM	C4A-C3A-C2A	3.24	110.52	106.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	HEM	CHA-C4D-ND	3.17	128.29	124.37
2	A	601	HEM	CHD-C4C-C3C	-3.14	119.92	125.21
3	A	602	X2N	CAA-CAU-CBS	3.11	118.05	113.43
3	A	602	X2N	CAW-CAY-NBU	-3.07	104.33	110.78
2	A	601	HEM	CAA-C2A-C1A	-3.05	118.97	124.94
3	A	602	X2N	CAN-CBN-NBW	3.03	123.57	119.62
3	A	602	X2N	FAE-CBI-CAG	-3.00	112.63	118.64
2	A	601	HEM	C4D-ND-C1D	2.86	108.59	105.21
2	A	601	HEM	C3B-C4B-NB	-2.84	107.43	109.47
3	A	602	X2N	CAM-CBM-NBU	-2.83	117.46	121.39
3	A	602	X2N	CAL-CBM-NBU	2.80	125.28	121.39
3	A	602	X2N	CAX-NBT-CAW	2.66	117.56	111.57
2	A	601	HEM	C3C-C2C-C1C	-2.66	104.53	107.05
3	A	602	X2N	NBW-CAT-NBF	-2.65	106.42	111.42
3	A	602	X2N	CBO-CAP-CAG	-2.62	119.77	123.23
3	A	602	X2N	C2-CBC-NBV	2.58	115.86	112.38
3	A	602	X2N	CAT-NBW-CBP	2.49	109.11	107.27
2	A	601	HEM	CMC-C2C-C3C	-2.44	122.52	128.43
2	A	601	HEM	CHC-C1C-C2C	-2.37	120.55	125.49
2	A	601	HEM	C4C-NC-C1C	2.37	109.68	105.82
3	A	602	X2N	FAE-CBI-CAR	2.33	121.57	119.01
3	A	602	X2N	CBS-NBX-NBF	-2.32	118.61	121.28
3	A	602	X2N	CAM-CAO-CBN	2.27	123.17	120.30
3	A	602	X2N	CAQ-NBE-NBV	2.20	107.68	102.60
3	A	602	X2N	FAF-CAP-CBO	2.18	122.03	118.55
2	A	601	HEM	CBD-CAD-C3D	-2.07	106.82	112.53
3	A	602	X2N	CAO-CBN-CAN	-2.01	115.34	119.18

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	X2N	CAB-CBQ-CBS-NBX
3	A	602	X2N	OAD-CBQ-CBS-CAU
3	A	602	X2N	OAD-CBQ-CBS-NBX
3	A	602	X2N	CAU-CBS-NBX-CBP
4	A	603	GOL	O1-C1-C2-C3
4	A	603	GOL	C1-C2-C3-O3
4	A	603	GOL	O2-C2-C3-O3
2	A	601	HEM	C3A-C2A-CAA-CBA
3	A	602	X2N	CAN-CBN-NBW-CAT
4	A	603	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	HEM	C2B-C3B-CAB-CBB
3	A	602	X2N	CAO-CBN-NBW-CAT
3	A	602	X2N	CBQ-CBS-NBX-CBP
3	A	602	X2N	CBQ-CBS-NBX-NBF
2	A	601	HEM	CAA-CBA-CGA-O1A
2	A	601	HEM	CAA-CBA-CGA-O2A
3	A	602	X2N	C7-CBB-OBG-CBK
2	A	601	HEM	CAD-CBD-CGD-O1D
2	A	601	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.

3 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	X2N	7	0
4	A	603	GOL	2	0
2	A	601	HEM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand X2N A 602



Ligand HEM A 601



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/456 (95%)	-0.19	10 (2%) 61 57	14, 36, 71, 113	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	233	LEU	5.2
1	A	229	LEU	4.7
1	A	235	GLY	4.0
1	A	230	ALA	3.8
1	A	81	HIS	3.7
1	A	232	PHE	3.2
1	A	491	TRP	3.0
1	A	236	LYS	2.5
1	A	132	GLU	2.3
1	A	237	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	603	6/6	0.80	0.15	79,80,81,81	0
3	X2N	A	602	51/51	0.88	0.15	28,57,99,100	0
2	HEM	A	601	43/43	0.99	0.05	11,17,25,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around HEM A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.