



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 10:31 PM UTC

PDB ID : 4J1U / pdb_00004j1u
Title : Crystal structure of antibody 93F3 unstable variant
Authors : Wang, F.
Deposited on : 2013-02-02
Resolution : 2.58 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	66% 26% ...
1	C	219	61% 33% ..
1	E	219	3% 60% 32% 5% ..
2	B	238	63% 19% 5% 13%
2	D	238	3% 45% 37% 6% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	238	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '62%', a yellow segment labeled '22%', and a small grey segment at the end labeled '13%'. There is a small black dot on the yellow segment.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody 93F3 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1610	1011	264	329	6			
1	C	213	Total	C	N	O	S	0	0	0
			1630	1017	273	334	6			
1	E	214	Total	C	N	O	S	0	0	0
			1644	1028	278	332	6			

- Molecule 2 is a protein called antibody 93F3 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	0	0	0
			1530	968	251	305	6			
2	D	210	Total	C	N	O	S	0	0	0
			1547	981	254	306	6			
2	F	207	Total	C	N	O	S	0	0	0
			1531	970	252	303	6			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	O	0	0
			10	10		
3	B	8	Total	O	0	0
			8	8		
3	C	25	Total	O	0	0
			25	25		
3	D	8	Total	O	0	0
			8	8		
3	E	11	Total	O	0	0
			11	11		
3	F	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody 93F3 Light chain




- Molecule 1: antibody 93F3 Light chain




- Molecule 1: antibody 93F3 Light chain




- Molecule 2: antibody 93F3 Heavy chain



- Molecule 2: antibody 93F3 Heavy chain



- Molecule 2: antibody 93F3 Heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.03Å 112.57Å 86.51Å 90.00° 115.28° 90.00°	Depositor
Resolution (Å)	78.33 – 2.58 78.22 – 2.58	Depositor EDS
% Data completeness (in resolution range)	94.8 (78.33-2.58) 94.8 (78.22-2.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.210 , 0.276 0.210 , 0.276	Depositor DCC
R_{free} test set	2121 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	56.9	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9570	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1643	1.07	5/2228 (0.2%)
1	C	0.85	0/1662	1.08	4/2253 (0.2%)
1	E	1.01	3/1678 (0.2%)	1.28	15/2275 (0.7%)
2	B	0.85	0/1566	1.05	4/2133 (0.2%)
2	D	0.90	0/1586	1.13	9/2163 (0.4%)
2	F	0.94	2/1567 (0.1%)	1.08	6/2134 (0.3%)
All	All	0.90	5/9702 (0.1%)	1.12	43/13186 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	E	0	2
2	D	0	1
All	All	0	5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	35	ARG	NE-CZ	11.92	1.46	1.33
1	E	35	ARG	CZ-NH1	-9.57	1.19	1.32
2	F	168	HIS	CE1-NE2	5.43	1.38	1.32
2	F	168	HIS	CG-CD2	5.23	1.41	1.35
1	E	41	TRP	CA-C	-5.01	1.46	1.52

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	147	ARG	CA-C-N	12.80	144.74	121.70
1	E	147	ARG	C-N-CA	12.80	144.74	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	35	ARG	NE-CZ-NH2	11.71	129.74	119.20
1	E	147	ARG	N-CA-C	10.97	124.28	111.11
1	A	147	ARG	N-CA-C	8.86	120.55	111.07

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	46	PRO	Peptide
1	A	48	GLN	Peptide
2	D	111	THR	Peptide
1	E	46	PRO	Peptide
1	E	48	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1610	0	1573	48	0
1	C	1630	0	1588	73	0
1	E	1644	0	1609	64	0
2	B	1530	0	1509	35	0
2	D	1547	0	1519	90	0
2	F	1531	0	1511	35	0
3	A	10	0	0	0	0
3	B	8	0	0	0	0
3	C	25	0	0	1	0
3	D	8	0	0	0	0
3	E	11	0	0	0	0
3	F	16	0	0	1	0
All	All	9570	0	9309	331	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 331 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:ARG:N	1:A:148:GLU:HB2	1.50	1.23
1:E:147:ARG:N	1:E:148:GLU:HG2	1.59	1.15
1:E:4:MET:HE3	1:E:96:GLN:HB3	1.32	1.11
2:B:154:VAL:HA	2:B:155:THR:HG22	1.20	1.11
2:B:104:GLY:HA3	2:B:105:ASP:HB2	1.09	1.07

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/219 (94%)	188 (91%)	15 (7%)	3 (2%)	8	16
1	C	209/219 (95%)	193 (92%)	11 (5%)	5 (2%)	4	8
1	E	212/219 (97%)	186 (88%)	18 (8%)	8 (4%)	2	3
2	B	202/238 (85%)	179 (89%)	16 (8%)	7 (4%)	3	4
2	D	206/238 (87%)	176 (85%)	16 (8%)	14 (7%)	1	1
2	F	201/238 (84%)	184 (92%)	13 (6%)	4 (2%)	6	11
All	All	1236/1371 (90%)	1106 (90%)	89 (7%)	41 (3%)	3	4

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	148	GLU
2	B	105	ASP
2	B	153	PRO
2	B	162	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/193 (96%)	170 (92%)	15 (8%)	11	23
1	C	187/193 (97%)	172 (92%)	15 (8%)	11	24
1	E	188/193 (97%)	171 (91%)	17 (9%)	9	19
2	B	176/205 (86%)	156 (89%)	20 (11%)	5	11
2	D	176/205 (86%)	165 (94%)	11 (6%)	16	34
2	F	176/205 (86%)	158 (90%)	18 (10%)	7	14
All	All	1088/1194 (91%)	992 (91%)	96 (9%)	9	20

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	146	VAL
1	E	96	GLN
2	D	190	SER
1	E	51	THR
1	E	165	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	168	HIS
1	E	143	ASN
2	F	203	ASN
1	E	160	GLN
1	E	96	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/219 (95%)	0.06	1 (0%) 87 86	37, 66, 89, 110	0
1	C	213/219 (97%)	0.04	1 (0%) 87 86	34, 59, 89, 112	0
1	E	214/219 (97%)	0.04	2 (0%) 81 79	31, 59, 95, 127	0
2	B	208/238 (87%)	0.18	0 100 100	37, 67, 97, 113	0
2	D	210/238 (88%)	0.56	6 (2%) 53 49	49, 85, 116, 131	0
2	F	207/238 (86%)	0.04	5 (2%) 59 55	35, 57, 84, 127	0
All	All	1262/1371 (92%)	0.15	15 (1%) 76 74	31, 65, 101, 131	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	141	ALA	3.9
2	F	52	TRP	3.8
2	D	98	HIS	2.8
1	A	31	ASN	2.7
2	F	105	ASP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.